

論文 / 著書情報
Article / Book Information

Title	Global stability of traveling curved fronts in the Allen-Cahn equations
Authors	Hirokazu Ninomiya, Masaharu Taniguchi
Citation	Discrete and Continuous Dynamical Systems, Vol. 15, No. 3, pp. 819—832
Pub. date	2006,
URL	http://www.aims sciences.org/journals/displayArticles.jsp?paperID=1806
Copyright	Copyright (c) 2006 American Institute of Mathematical Sciences,AIMS

GLOBAL STABILITY OF TRAVELING CURVED FRONTS IN THE ALLEN-CAHN EQUATIONS

HIROKAZU NINOMIYA

Department of Applied Mathematics and Informatics,
Ryukoku University, Seta Otsu 520-2194, Japan

MASAHARU TANIGUCHI

Department of Mathematical and Computing Sciences,
Tokyo Institute of Technology,
O-okayama 2-12-1-W8-38, Tokyo 152-8552, Japan

(Communicated by Eiji Yanagida)

Abstract. This paper is concerned with the global stability of a traveling curved front in the Allen-Cahn equation. The existence of such a front is recently proved by constructing supersolutions and subsolutions. In this paper, we introduce a method to construct new subsolutions and prove the asymptotic stability of traveling curved fronts globally in space.

1. Introduction. Front propagation phenomena are often observed in various fields (e.g. [16]). We study a traveling curved front with V-formed contour lines in $\mathbb{R}^2$ in bistable media as one of the simplest non-planar multidimensional traveling waves. Such a front propagates due to the curvature effect and the difference of energy densities between two states.

In this paper we consider a solution $u(\cdot, \cdot, t) \in L^\infty(\mathbb{R}^2)$ of the Allen-Cahn equation:

$$\begin{aligned} u_t &= \Delta u + f(u) & (x, y) \in \mathbb{R}^2, t > 0, \\ u(x, y, 0) &= u_0(x, y) & (x, y) \in \mathbb{R}^2. \end{aligned} \quad (1.1)$$

Here the nonlinear term f is of bistable type and a given initial function u_0 is bounded and of class C^1 .

We consider the case where the constant states -1 and 1 are asymptotically stable and where the region of the state 1 is getting larger and larger and finally it covers the whole space. When the state 1 propagates, we can observe the characteristic profiles. In the one-dimensional space, one of the typical solutions is a traveling wave solution which always keeps its shape. Setting $u(x, t) = \Phi(\eta)$ and $\eta = x - kt$, we have

$$\begin{aligned} -\Phi''(\eta) - k\Phi'(\eta) - f(\Phi) &= 0, & \Phi'(\eta) < 0 & \quad (-\infty < \eta < \infty), \\ \Phi(-\infty) &= 1, & \Phi(+\infty) &= -1. \end{aligned} \quad (1.2)$$

The following is the standing assumptions on f .

(A1) f is of class C^1 with $f(1) = 0$, $f(-1) = 0$, $f'(1) < 0$ and $f'(-1) < 0$.

(A2) $\int_{-1}^1 f(s) ds > 0$.

2000 *Mathematics Subject Classification.* 35K57, 35B35.

Key words and phrases. traveling wave, Allen-Cahn equation, curved front, global stability.

(A3) $f(s) < 0$ and $f'(s) < 0$ for $s > 1$. $f(s) > 0$ and $f'(s) < 0$ for $s < -1$.

(A4) There exist $k > 0$ and a function $\Phi(\eta)$ satisfying (1.2).

In (1.2), the speed k is uniquely determined and the traveling wave Φ is unique up to phase shift under the assumptions (A1), (A2) and (A3) as in [5] and [3].

Example 1. If f satisfies

$$\begin{aligned} f(s) &> 0 && \text{for } s \in (\beta, 1), \\ f(s) &< 0 && \text{for } s \in (-1, \beta), \\ f'(\beta) &> 0 \end{aligned}$$

for some $\beta \in (-1, 1)$ in addition to (A1)-(A3), then it satisfies (A4).

A typical example of f is

$$f(u) = -(u+1)(u+a)(u-1), \quad (1.3)$$

where $a \in (0, 1)$ is a given number. For the nonlinearity (1.3), one has $k = \sqrt{2}a$ and $\Phi(\eta) = -\tanh(\eta/\sqrt{2})$.

There are many works on one-dimensional traveling waves. One can see [5], [1], [10] and [11].

Example 2. Another example is the following nonlinear term

$$f(u) = -u \left(u^2 - \frac{1}{4} \right) (u^2 - 1) + a(1 - u^2) \sqrt{8u^2 + 1},$$

where $0 < a < 1/4$. This f satisfies $f(\pm 1) = 0$, and it has three zero points in $(-1, 1)$ if $|a|$ is small enough. To solve (1.2) it suffices to solve the following two relations

$$\begin{aligned} \Phi''(\eta) &= \Phi \left(\Phi^2 - \frac{1}{4} \right) (\Phi^2 - 1) \\ -k\Phi'(\eta) &= a(1 - \Phi^2) \sqrt{8\Phi^2 + 1} \end{aligned} \quad -\infty < \eta < \infty.$$

From the first relation Φ is a unique solution of

$$\int_0^{\Phi(\eta)} \frac{dv}{(1-v^2)\sqrt{8v^2+1}} = -\frac{\sqrt{6}}{12}\eta$$

if one assumes $\Phi(0) = 0$. This $\Phi(\eta)$ satisfies the second relation if $k = 2\sqrt{6}a$. Terman [19] studied the existence of traveling waves when the nonlinear function has multiple zeros.

For multi-dimensional traveling wave solutions under the bistable condition (A1), Fife [4] first studied traveling curved fronts as a singular limit problem by putting a small coefficient to the diffusion term. The existence and stability of traveling curved fronts in (1.1) are recently studied by [14], [8] and [9]. For the existence and the stability of two-dimensional traveling curved fronts, see [14]. For the existence and properties two-dimensional traveling curved fronts and higher dimensional conical waves, see [8] and [9].

Related works in the interface equation are studied in [13] and [12]. See [13] and the references therein for the detail.

For nonlinearity appearing in combustion problems, the stability of the traveling curved fronts is studied in [7] for exponentially decaying perturbations that belongs to $[-1, 1]$.




FIGURE 1. The profiles of f and a traveling wave in Example 2 for $a = 1/50$.

For the bistable nonlinearity including the Allen-Cahn equations, one can expect a wider class of initial perturbations which leads to the asymptotic stability of traveling curved fronts, since the two rest states $(+1$ and $-1)$ are asymptotically stable. This is the main issue of this paper.

We study traveling wave solutions. We assume that the solutions travel towards y -direction without loss of generality. We put

$$u(x, y, t) = w(x, y - ct, t), \quad z = y - ct.$$

Then we obtain

$$\begin{aligned} w_t - w_{xx} - w_{zz} - cw_z - f(w) &= 0, & (x, z) \in \mathbb{R}^2, t > 0. \\ w|_{t=0} &= u_0 & \text{in } \mathbb{R}^2. \end{aligned} \quad (1.4)$$

We denote the solution of (1.4) with $w(x, z, 0; u_0) = u_0(x, z)$ by $w(x, z, t; u_0)$.

The traveling wave v with speed c in y -direction becomes a stationary solution of (1.4), namely, v satisfies

$$\mathcal{L}[v] = 0 \quad (1.5)$$

where

$$\mathcal{L}[v] \stackrel{\text{def}}{=} -v_{xx} - v_{zz} - cv_z - f(v).$$

Hereafter the speed c is an arbitrary constant that is bigger than k . We see that $\Phi(k(z \pm m_*x)/c)$ satisfy (1.5), where

$$m_* \stackrel{\text{def}}{=} \frac{\sqrt{c^2 - k^2}}{k} > 0.$$

They are so called *planar traveling fronts*.

Since the maximum of subsolutions is also a subsolution, it turns out that

$$\begin{aligned} v^-(x, z) &\stackrel{\text{def}}{=} \max \left\{ \Phi \left(\frac{k}{c} (z - m_*x) \right), \Phi \left(\frac{k}{c} (z + m_*x) \right) \right\} \\ &= \Phi \left(\frac{k}{c} (z - m_*|x|) \right) \end{aligned}$$

is a subsolution of (1.4). This $v^-(x, z)$ is strictly monotone decreasing in z .

If v^- is given as an initial condition, it converges to a traveling curved front. The existence of such traveling waves is shown in [14].

Theorem 1.1 ([14, Theorems 1.2 and 1.3]). *There exists a traveling wave solution $u(x, y, t) = v_*(x, y - ct)$ of (1.1) with*

$$\lim_{R \rightarrow \infty} \sup_{x^2 + z^2 > R^2} |v_*(x, z) - v^-(x, z)| = 0, \quad (1.6)$$

$$v^-(x, z) < v_*(x, z).$$

This traveling curved front v^ also satisfies the following property: for $0 < \delta < 1$, there exists a positive constant γ such that*

$$(v_*)_z \geq \gamma > 0 \quad \text{if } -1 + \delta \leq v_* \leq 1 - \delta.$$

Moreover, if $u_0(x, y)$ satisfies

$$\lim_{R \rightarrow \infty} \sup_{x^2 + y^2 > R^2} |u_0(x, y) - v^-(x, y)| = 0, \quad (1.7)$$

$$v^-(x, y) \leq u_0(x, y), \quad (1.8)$$

then the solution $u(x, y, t; u_0)$ of (1.1) satisfies

$$\lim_{t \rightarrow \infty} \sup_{(x, y) \in \mathbb{R}^2} |u(x, y, t; u_0) - v_*(x, y - ct)| = 0.$$

If $v(x, z)$ is a solution of (1.5) and satisfies

$$\lim_{R \rightarrow \infty} \sup_{x^2 + z^2 > R^2} |v(x, z) - v^-(x, z)| = 0, \quad (1.9)$$

then $v(x, y - ct)$ is called a *traveling curved front*. Actually the contours of the traveling curved front are seen as in Figure 2. Theorem 1.1 shows that the traveling curved front with the speed c exists for each $c > k$. Moreover, it says the convergence of the solutions satisfying (1.7) and (1.8) to the traveling curved front. The assumption (1.7) is essential to the speed of the traveling curved front. We can expect that the solution starting from v^- converges to v_* because the solution for $t > 0$ must satisfy (1.8). Due to the continuity of the initial functions we can also expect that the solution of (1.1) converges to v_* under the assumption (1.7) even if an initial state close to v^- is less than v^- . Actually we will show that the theorem holds true without the assumption (1.8). For this purpose we use supersolutions and subsolutions. In [14] a supersolution is constructed so that it is larger than any given initial state u_0 with (1.7). Now we need to introduce a subsolution that is smaller than any given u_0 with (1.7).

We construct such a subsolution in Section 2 and prove the following theorem.

Theorem 1.2 (global stability). *If $u_0(x, y)$ satisfies (1.7), then the solution $u(x, y, t; u_0)$ of (1.1) satisfies*

$$\lim_{t \rightarrow \infty} \sup_{(x, y) \in \mathbb{R}^2} |u(x, y, t; u_0) - v_*(x, y - ct)| = 0.$$

This theorem implies that v_* is the unique traveling curved front.

2. Construction of subsolutions. We introduce several constants for preparation. There exists a positive constant δ_1 ($0 < \delta_1 < 1/4$) with

$$-f'(s) > \kappa_1 \quad \text{if } |s + 1| < 2\delta_1 \text{ or } |s - 1| < 2\delta_1,$$

where

$$\kappa_1 \stackrel{\text{def}}{=} \frac{1}{2} \min \{-f'(-1), -f'(1)\} > 0.$$




FIGURE 2. The contours of the numerical traveling curved front with the nonlinear function f as in (1.3), $a = 1/2$ and $m_* = \sqrt{3}$.

Since $\Phi(\mu)$ is monotone decreasing in μ , we can define positive constants A and B by

$$\Phi(-A) = 1 - \frac{\delta_1}{2}, \quad \Phi(B) = -1 + \frac{\delta_1}{2},$$

respectively. Then

$$-1 + \frac{\delta_1}{2} < \Phi(\mu) < 1 - \frac{\delta_1}{2}$$

is equivalent to $-A < \mu < B$.

There exist positive constants γ_1 and K_1 with

$$\max \{ |\Phi'(\mu)|, |\Phi''(\mu)| \} \leq K_1 \operatorname{sech}(\gamma_1 |\mu|) \quad (2.10)$$

for any $\mu \in \mathbb{R}$.

Take

$$\psi(\zeta) \stackrel{\text{def}}{=} \frac{1}{m_* \gamma_2} \log \left(1 + \exp(\gamma_2 \zeta) \right). \quad (2.11)$$

The right-hand side equals

$$\frac{1}{2m_*} \int_{-\infty}^{\zeta} \left(1 + \tanh \left(\frac{1}{2} \gamma_2 \zeta' \right) \right) d\zeta'.$$

There exist some constants $K_i > 0$ ($i = 2, 3, 4$) and $\gamma_2 > 0$ so that $\psi(\zeta)$ satisfies

$$\max \left\{ \left| \psi(\zeta) - \frac{\zeta}{m_*} \right|, \left| \psi'(\zeta) - \frac{1}{m_*} \right| \right\} \leq K_2 \operatorname{sech}(\gamma_2 \zeta) \quad \text{for } \zeta \geq 0, \quad (2.12)$$

$$\max \{ |\psi(\zeta)|, |\psi'(\zeta)| \} \leq K_2 \operatorname{sech}(\gamma_2 \zeta) \quad \text{for } \zeta \leq 0, \quad (2.13)$$

$$\max \{ |\psi''(\zeta)|, |\psi'''(\zeta)| \} \leq K_2 \operatorname{sech}(\gamma_2 \zeta) \quad \text{for } \zeta \in \mathbb{R}, \quad (2.14)$$

$$k - \frac{c\psi'(\zeta)}{\sqrt{1 + \psi'(\zeta)^2}} \geq K_3 \min \{ 1, \exp(-\gamma_2 \zeta) \} \quad \text{for } \zeta \in \mathbb{R}, \quad (2.15)$$

$$0 \leq \frac{c}{\sqrt{1 + \psi'(\zeta)^2}} - km_* \leq K_4 \min \{ 1, \exp(-\gamma_2 \zeta) \} \quad \text{for } \zeta \in \mathbb{R}. \quad (2.16)$$

Then we construct a subsolution to (1.5) as follows.

Theorem 2.1 (subsolution). *There exist a positive constant ε_0 and a positive function $\alpha_0(\varepsilon)$ so that, for $0 < \varepsilon < \varepsilon_0$ and $0 < \alpha < \alpha_0(\varepsilon)$,*

$$v_1(x, z; \varepsilon, \alpha) \stackrel{\text{def}}{=} \Phi \left(\frac{\psi(\alpha z) - \alpha x}{\alpha \sqrt{1 + \psi'(\alpha z)^2}} \right) - \varepsilon \operatorname{sech}(\gamma_2 \alpha z) \quad (2.17)$$

is a subsolution to (1.5). Moreover, there exists a positive constant γ_3 with

$$(v_1)_x \geq \gamma_3 \quad \text{if } -1 + \delta_1 \leq v_1 \leq 1 - \delta_1. \quad (2.18)$$

Proof. We set

$$\begin{aligned} \xi &\stackrel{\text{def}}{=} \alpha x, & \zeta &\stackrel{\text{def}}{=} \alpha z, \\ \mu &\stackrel{\text{def}}{=} \frac{1}{\sqrt{1 + \psi'(\alpha z)^2}} \left(\frac{1}{\alpha} \psi(\alpha z) - x \right), \\ \sigma(\zeta) &\stackrel{\text{def}}{=} \varepsilon \operatorname{sech}(\gamma_2 \zeta). \end{aligned} \quad (2.19)$$

Then we have $v_1 = \Phi(\mu) - \sigma(\zeta)$. The chain rule gives

$$\mu_x = -\frac{1}{\sqrt{1 + \psi'(\zeta)^2}}, \quad \mu_{xx} = 0, \quad (2.20)$$

$$\mu_z = \frac{\psi'(\zeta)}{\sqrt{1 + \psi'(\zeta)^2}} - \frac{\alpha \psi'(\zeta) \psi''(\zeta)}{1 + \psi'(\zeta)^2} \mu, \quad (2.21)$$

$$\mu_{zz} = \frac{\alpha \psi''(\zeta) (1 - \psi'(\zeta)^2)}{(1 + \psi'(\zeta)^2)^{\frac{3}{2}}} + \frac{\alpha^2 g(\zeta)}{(1 + \psi'(\zeta)^2)^2} \mu, \quad (2.22)$$

where

$$g(\zeta) = 3\psi'(\zeta)^2 \psi''(\zeta)^2 - \left(1 + \psi'(\zeta)^2 \right) \left(\psi''(\zeta)^2 + \psi'(\zeta) \psi'''(\zeta) \right).$$

Using these equalities and (1.2), we get

$$\begin{aligned} \mathcal{L}[v_1] &= -\frac{\Phi''(\mu)}{1 + \psi'(\zeta)^2} - (\Phi'(\mu) \mu_z)_z - c \Phi'(\mu) \mu_z - f(\Phi(\mu) - \sigma(\zeta)) \\ &\quad + \alpha^2 \sigma''(\zeta) + c \alpha \sigma'(\zeta) + \Phi''(\mu) + k \Phi'(\mu) + f(\Phi) \end{aligned}$$

and thus

$$\begin{aligned} \mathcal{L}[v_1] &= \left(1 - \frac{1}{1 + \psi'(\zeta)^2} - \mu_z^2 \right) \Phi''(\mu) - \mu_{zz} \Phi'(\mu) + (k - c \mu_z) \Phi'(\mu) \\ &\quad + f(\Phi(\mu)) - f(\Phi(\mu) - \sigma(\zeta)) + \alpha^2 \sigma''(\zeta) + c \alpha \sigma'(\zeta) \\ &= I_1 + I_2 + I_3, \\ I_1 &\stackrel{\text{def}}{=} \left(1 - \frac{1}{1 + \psi'(\zeta)^2} - \mu_z^2 \right) \Phi''(\mu) + \left(-\mu_{zz} + \frac{c \alpha \psi'(\zeta) \psi''(\zeta)}{1 + \psi'(\zeta)^2} \mu \right) \Phi'(\mu), \\ I_2 &\stackrel{\text{def}}{=} \left(k - \frac{c \psi'(\zeta)}{\sqrt{1 + \psi'(\zeta)^2}} \right) \Phi'(\mu), \\ I_3 &\stackrel{\text{def}}{=} f(\Phi(\mu)) - f(\Phi(\mu) - \sigma(\zeta)) + \alpha^2 \sigma''(\zeta) + c \alpha \sigma'(\zeta). \end{aligned}$$

By (2.21) and (2.22), we have

$$\begin{aligned} I_1 = & \alpha \left(- \left(\frac{\psi'(\zeta)\psi''(\zeta)}{1+\psi'(\zeta)^2} \right)^2 \alpha \mu^2 + \frac{2\psi'(\zeta)^2\psi''(\zeta)}{(1+\psi'(\zeta)^2)^{\frac{3}{2}}} \mu \right) \Phi''(\mu) \\ & + \alpha \left(\frac{\psi''(\zeta)^2 + \psi'(\zeta)\psi'''(\zeta)}{1+\psi'(\zeta)^2} \alpha \mu - \frac{3\psi'(\zeta)^2\psi''(\zeta)^2}{(1+\psi'(\zeta)^2)^2} \alpha \mu \right. \\ & \left. - \frac{(1-\psi'(\zeta)^2)\psi''(\zeta)}{(1+\psi'(\zeta)^2)^{\frac{3}{2}}} + \frac{c\psi'(\zeta)\psi''(\zeta)}{1+\psi'(\zeta)^2} \mu \right) \Phi'(\mu) \end{aligned}$$

From (2.10)–(2.16), we have

$$|I_1| \leq K_5 \alpha \operatorname{sech}(\gamma_2 \zeta)$$

with positive constant K_5 that is independent of ε and α . Now we assume

$$0 < \varepsilon < \varepsilon_0 \leq \frac{\delta_1}{2}. \quad (2.23)$$

We have

$$I_2 \leq -K_3 |\Phi'(\mu)| \min\{1, \exp(-\gamma_2 \zeta)\}. \quad (2.24)$$

Here we have

$$I_3 = -\sigma(\zeta) \int_0^1 (-f'(\Phi(\mu) - s\sigma(\zeta))) ds + \alpha \sigma''(\zeta) + c\alpha \sigma'(\zeta).$$

If α is small enough to satisfy

$$0 < \alpha < \min \left\{ 1, \frac{\kappa_1}{2\gamma_2} \inf_{\zeta \in \mathbb{R}} \frac{\operatorname{sech} \zeta}{\gamma_2 |\operatorname{sech}'' \zeta| + |c| |\operatorname{sech}' \zeta|} \right\}, \quad (2.25)$$

we get

$$|I_3| \leq K_6 \varepsilon \operatorname{sech}(\gamma_2 \zeta) \quad \text{if } -A < \mu < B, \quad (2.26)$$

$$I_3 \leq -\frac{1}{2} \kappa_1 \sigma(\zeta) \quad \text{if } \mu \leq -A \text{ or } \mu \geq B. \quad (2.27)$$

If $\mu \leq -A$ or $\mu \geq B$, we have

$$\mathcal{L}[v_1] \leq -\frac{1}{2} \kappa_1 \sigma(\zeta) + K_5 \alpha \operatorname{sech}(\gamma_2 \zeta) < 0$$

assuming

$$\alpha < \frac{\kappa_1}{2K_5} \varepsilon. \quad (2.28)$$

If $-A \leq \mu \leq B$, we have

$$\mathcal{L}[v_1] \leq -K_3 |\Phi'(\mu)| \min\{1, \exp(-\gamma_2 \zeta)\} + (K_5 \alpha + K_6 \varepsilon) \operatorname{sech}(\gamma_2 \zeta).$$

The right-hand side is negative if

$$K_5 \alpha + K_6 \varepsilon \leq K_3 \min_{-A \leq \mu \leq B} |\Phi'(\mu)| \inf_{\zeta \in \mathbb{R}} \frac{\min\{1, \exp(-\zeta)\}}{\operatorname{sech} \zeta}. \quad (2.29)$$

Thus we obtain

$$\mathcal{L}[v_1] < 0 \quad \text{in } \mathbb{R}^2$$

under (2.23), (2.25), (2.28) and (2.29).

Finally we prove (2.18). We have

$$(v_1)_x = \frac{-\Phi'(\mu)}{\sqrt{1+\psi'(\zeta)^2}} > 0.$$

Since

$$\sup_{\zeta \in \mathbb{R}} |\psi'(\zeta)| \leq \frac{1}{m_*},$$

it suffices to show that $|\mu|$ is bounded if $-1 + \delta_1 \leq v_1 \leq 1 - \delta_1$. Recalling that $v_1 = \Phi(\mu) - \varepsilon \operatorname{sech}(\gamma_2 \zeta)$ and $0 < \varepsilon < \delta_1/2$, we obtain

$$-1 + \frac{\delta_1}{2} \leq \Phi(\mu) \leq 1 - \frac{\delta_1}{2}$$

and thus $-A \leq \mu \leq B$ from $-1 + \delta_1 \leq v_1 \leq 1 - \delta_1$. Putting

$$\gamma_3 \stackrel{\text{def}}{=} \frac{1}{\sqrt{1 + (1/m_*)^2}} \min_{-A \leq \mu \leq B} |\Phi'(\mu)|,$$

we obtain (2.18). $\square$

This lemma implies that

$$v_2(x, z) \stackrel{\text{def}}{=} v_1(-x, z) \quad (2.30)$$

is also subsolution of (1.5) with

$$-(v_2)_x \geq \gamma_3 \quad \text{if } -1 + \delta_1 \leq v_2 \leq 1 - \delta_1.$$

Lemma 2.2. *Let $w_j(x, z, t)$ ($j = 1, 2$) be defined by*

$$\begin{aligned} w_1(x, z, t) &\stackrel{\text{def}}{=} v_1(x - \rho\delta(1 - e^{-\beta t}), z) - \delta e^{-\beta t}, \\ w_2(x, z, t) &\stackrel{\text{def}}{=} v_2(x + \rho\delta(1 - e^{-\beta t}), z) - \delta e^{-\beta t}. \end{aligned}$$

Then there exists a large positive constant ρ such that, for any $\delta \in (0, \delta_1/2]$, w_1 and w_2 are also subsolutions of (1.4).

Proof. First we show for w_1 . Theorem 2.1 and simple calculation imply that

$$\begin{aligned} (w_1)_t + \mathcal{L}[w_1] &= -(v_1)_x \rho \delta \beta e^{-\beta t} + \delta \beta e^{-\beta t} - (v_1)_{xx} - (v_1)_{zz} \\ &\quad - c(v_1)_z - f(v_1 - \delta e^{-\beta t}) \\ &\leq -\delta e^{-\beta t} \left(\rho \beta (v_1)_x - \beta - \int_0^1 f'(v_1 - s\delta e^{-\beta t}) ds \right). \end{aligned}$$

We get

$$\rho \beta (v_1)_x - \beta - \int_0^1 f'(v_1 - s\delta e^{-\beta t}) ds \geq \beta \left(\gamma_3 \rho - 1 - \frac{K_7}{\beta} \right)$$

when $-1 + \delta_1 \leq v_1 \leq 1 - \delta_1$ where

$$K_7 \stackrel{\text{def}}{=} \sup_{|s| \leq 2} |f'(s)|. \quad (2.31)$$

For $v_1 < -1 + \delta_1$ or $v_1 > 1 - \delta_1$, we have an estimate

$$\rho \beta (v_1)_x - \beta - \int_0^1 f'(v_1 \pm s\delta e^{-\beta t}) ds \geq \kappa_1 - \beta.$$


We choose a small positive constant $\beta > 0$ and a large positive constant ρ satisfying

$$0 < \beta < \kappa_1, \quad \rho > \frac{\beta + K_7}{\beta \gamma_3}. \quad (2.32)$$

This immediately implies

$$(w_1)_t + \mathcal{L}[w_1] \leq 0.$$

Thus w_1 is a subsolution.

FIGURE 3. The graph of v_4 and its contours.

Since $v_2(x, z) = v_1(-x, z)$, we have

$$\begin{aligned} \mathcal{L}[w_2] &= (v_2)_x \rho \delta \beta e^{-\beta t} + \delta \beta e^{-\beta t} - (v_2)_{xx} - (v_2)_{zz} - c(v_2)_z - f(v_2 - \delta e^{-\beta t}) \\ &\leq -\delta e^{-\beta t} \left(-\rho \beta (v_2)_x - \beta - \int_0^1 f'(v_2 - s \delta e^{-\beta t}) ds \right). \end{aligned}$$

Similarly we can see that w_2 is also a subsolution. $\square$

3. Proof of Theorem 1.2. For simplicity, we denote $w(x, z, t; u_0)$ by $w(x, z, t)$ hereafter. Define

$$\begin{aligned} v_3(x, z) &\stackrel{\text{def}}{=} v_*(x, z + M), \\ v_4(x, z) &\stackrel{\text{def}}{=} \max\{v_1(x, z; \varepsilon, \alpha), v_2(x, z; \varepsilon, \alpha), v_3(x, z)\}, \end{aligned} \quad (3.33)$$

where M is a positive constant specified later. Recall that w_1 and w_2 are defined in Lemma 2.2. We also set

$$\begin{aligned} w_3(x, z, t) &\stackrel{\text{def}}{=} v_3(x, z + \rho \delta (1 - e^{-\beta t})) - \delta e^{-\beta t}, \\ w_4(x, z, t) &\stackrel{\text{def}}{=} \max\{w_1(x, z, t), w_2(x, z, t), w_3(x, z, t)\}. \end{aligned}$$


Note that $w_4(x, z, 0) = v_4(x, z) - \delta$.

Lemma 3.1. *For $0 < \varepsilon < \varepsilon_0$, $0 < \alpha < \alpha_0(\varepsilon)$ and $M > 0$, the limit of $w(x, z, t; v_4)$ as $t \rightarrow \infty$ exists in $L^\infty(\mathbb{R}^2)$ and the limit function*

$$v_{**}(x, z) \stackrel{\text{def}}{=} \lim_{t \rightarrow \infty} w(x, z, t; v_4)$$

satisfies

$$\begin{aligned} \mathcal{L}[v_{**}] &= 0, \\ v_{**} &\leq v_*, \\ \lim_{R \rightarrow \infty} \sup_{x^2 + z^2 > R^2} |v_{**} - v^-(x, z)| &= 0. \end{aligned}$$

FIGURE 4. A bird's-eye view of v_4 .

Proof. From (2.17), (2.30), (3.33) and the properties of ψ , we obtain

$$\lim_{R \rightarrow \infty} \sup_{x^2+z^2 > R^2} |v_1(x, z) - v^-(x, z)| = 0. \quad (3.34)$$

$$\lim_{R \rightarrow \infty} \sup_{x^2+z^2 > R^2} |v_4(x, z) - v^-(x, z)| = 0. \quad (3.35)$$

We note that

$$v_4 \leq \max\{v_4, v^-\}.$$

Since v_4 is a subsolution of (1.5), we have

$$v_4 \leq w(x, z, t; v_4) \leq w(x, z, t; \max\{v_4, v^-\})$$

and thus

$$v_4 \leq w(x, z, t; v_4) \leq w(x, z, t; \max\{v_4, v^-\}). \quad (3.36)$$

From Theorem 1.1 we have

$$\lim_{t \rightarrow \infty} \sup_{(x, z) \in \mathbb{R}^2} |w(x, z, t; \max\{v_4, v^-\}) - v_*(x, z)| = 0.$$

Since v_4 is a subsolution with (3.36), $w(x, z, t; v_4)$ is monotone increasing in t and the limiting function v_{**} exists with

$$\mathcal{L}[v_{**}] = 0, \quad (3.37)$$

$$v_4 \leq v_{**} \leq v_*. \quad (3.38)$$

See Sattinger [18] for instance. Thus

$$\lim_{R \rightarrow \infty} \sup_{x^2+z^2 > R^2} |v_{**}(x, z) - v^-(x, z)| = 0$$

follows from (3.38), (3.35) and

$$\lim_{R \rightarrow \infty} \sup_{x^2+z^2 > R^2} |v_*(x, z) - v^-(x, z)| = 0.$$

□

Lemma 3.2. *For any initial function u_0 satisfying (1.7) and any positive constant δ , there exist positive constants α, ε, T and M satisfying*

$$v_4(x, z) - \delta \leq w(x, z, T) \quad \text{for } (x, z) \in \mathbb{R}^2.$$

Proof. We define $W_i(t)$ ($i = 1, 2$) by

$$W'_i(t) = f(W_i(t)),$$

with initial conditions

$$\begin{aligned} W_1(0) &= \inf_{(x,z) \in \mathbb{R}^2} u_0(x, z), \\ W_2(0) &= \sup_{(x,z) \in \mathbb{R}^2} u_0(x, z). \end{aligned}$$

Then we have

$$W_1(t) \leq w(x, z, t) \leq W_2(t) \quad \text{for } (x, z) \in \mathbb{R}^2, t > 0$$

and thus

$$-1 \leq \liminf_{t \rightarrow \infty} w(x, z, t) \leq \limsup_{t \rightarrow \infty} w(x, z, t) \leq 1 \quad \text{for } (x, z) \in \mathbb{R}^2.$$

Namely, for any $\delta > 0$, there exists $T > 0$ with

$$-1 - \frac{\delta}{2} < w(x, z, t) \quad \text{for } (x, z) \in \mathbb{R}^2, t \geq T. \quad (3.39)$$

By Lemma 4.5 in [14], we have

$$\lim_{R \rightarrow \infty} \sup_{x^2 + z^2 > R^2} |w(x, z, T) - v^-(x, z)| = 0. \quad (3.40)$$

By (1.6), (3.39) and (3.40) we can choose a large constant M with

$$v_3(x, z) - \delta = v_*(x, z + M) - \delta < w(x, z, T) \quad \text{for } (x, z) \in \mathbb{R}^2.$$

By (3.40) there exists a positive constant R with

$$v^-(x, z) - \frac{\delta}{2} \leq w(x, z, T) \quad \text{for } x^2 + z^2 \geq R^2.$$

Using (3.34) we have

$$v_1(x, z; \varepsilon, \alpha) - \delta \leq v^-(x, z) - \frac{\delta}{2} \leq w(x, z, T) \quad \text{for } x^2 + z^2 \geq R^2.$$

For $x^2 + z^2 \leq R^2$, there exists a small positive constant α with

$$\begin{aligned} \frac{1}{\sqrt{1 + \psi'(\alpha z)^2}} \left(\frac{1}{\alpha} \psi(\alpha z) - x \right) &\geq \frac{1}{\sqrt{1 + \max_{|z| \leq R} \psi'(\alpha z)^2}} \left(\frac{1}{\alpha} \min_{|z| \leq R} \psi(\alpha z) - x \right) \\ &\geq \Phi^{-1} \left(-1 + \frac{\delta}{2} \right). \end{aligned}$$

Thus we have

$$v_1(x, z) - \delta \leq w(x, z, T) \quad \text{for } (x, z) \in \mathbb{R}^2. \quad (3.41)$$

Similarly we also have (3.41) with replacing v_1 by v_2 . $\square$

We use $\varepsilon > 0$ and $\alpha > 0$ as in Lemma 3.2. Hereafter we denote $v_j(x, z; \varepsilon, \alpha)$ simply by $v_j(x, z)$ for $j = 1, 2$ respectively.

Proposition 3.3. *The following holds true:*

$$v_{**}(x, z) = v_*(x, z)$$

for all x and z .

Proof. By the definition of v_{**} and Lemma 3.1, we have

$$v_*(x, z + M) \leq v_{**}(x, z) \leq v_*(x, z) \quad \text{in } \mathbb{R}^2.$$

Define

$$\Lambda \stackrel{\text{def}}{=} \inf\{\lambda : v_*(x, z + \lambda) \leq v_{**}(x, z)\}$$

Thus we have $0 \leq \Lambda \leq M$. We show $\Lambda = 0$ by contradiction. Assume $\Lambda > 0$. By (1.7), $v_*(\cdot, \cdot + \Lambda) \not\equiv v_{**}(\cdot, \cdot)$. By the strong maximum principle, we have

$$v_*(x, z + \Lambda) < v_{**}(x, z) \quad \text{in } \mathbb{R}^2. \quad (3.42)$$

As in [14, (5.4)], we can take R_* so large that

$$2\rho \sup_{|z - m_*|x| > R_* - \rho\delta_1} |(v_*)_z(x, z + \Lambda)| < 1.$$

By (3.42) and $\max\{v_1(x, z), v_2(x, z)\} \leq v_{**}(x, z) \leq v_*(x, z)$, there is a small positive constant $h(< \delta_1/2)$ with

$$v_*(x, z + \Lambda - 2\rho h) < v_{**}(x, z) \quad \text{in } D, \quad (3.43)$$

where

$$D \stackrel{\text{def}}{=} \{(x, z) : |z - m_*|x| \leq R_*\}.$$

In $\mathbb{R}^2 \setminus D$, we have

$$v_*(x, z + \Lambda - 2\rho h) - v_*(x, z + \Lambda) = -2\rho h \int_0^1 (v_*)_z(x, z + \Lambda - 2s\rho h) ds \leq h.$$

This implies

$$v_*(x, z + \Lambda - 2\rho h) - h \leq v_*(x, z + \Lambda) \quad \text{in } \mathbb{R}^2 \setminus D. \quad (3.44)$$

Combining (3.42) – (3.44), we have

$$v_*(x, z + \Lambda - 2\rho h) - h \leq v_{**}(x, z) \quad \text{in } \mathbb{R}^2.$$

Now [14, Lemma 4.4] says that $v_*(x, z + \Lambda - 2\rho h + \rho h(1 - e^{-\beta t})) - he^{-\beta t}$ is a subsolution of (1.4). Thus, for any $t > 0$,

$$v_*(x, z + \Lambda - 2\rho h + \rho h(1 - e^{-\beta t})) - he^{-\beta t} \leq v_{**}(x, z) \quad \text{in } \mathbb{R}^2.$$

Letting t tend to infinity, we have

$$v_*(x, z + \Lambda - \rho h) \leq v_{**}(x, z) \quad \text{in } \mathbb{R}^2.$$

This contradicts the definition of Λ . Thus we have $\Lambda = 0$ and $v_* \equiv v_{**}$. $\square$

Proof of Theorem 1.2. We denote $w(x, z, t)$ by $w(x, z, t; u_0)$ for simplicity. Note that

$$\min\{u_0, v^-\} \leq u_0 \leq \max\{u_0, v^-\}.$$

Theorem 1.1 says that, for any given $\delta > 0$, there exists $T > 0$ with

$$\sup_{(x, z) \in \mathbb{R}^2} |w(x, z, t; \max\{u_0, v^-\}) - v_*(x, z)| < \delta$$

for $t > T$. Using this inequality and the comparison principle, we have

$$w(x, z, t) < w(x, z, t; \max\{u_0, v^-\}) < v_*(x, z) + \delta$$

for all $(x, z) \in \mathbb{R}^2$, $t > T$. Thus we get

$$\limsup_{t \rightarrow \infty} w(x, z, t) \leq v_*(x, z)$$

for $(x, z) \in \mathbb{R}^2$. Then it suffices to show that

$$\liminf_{t \rightarrow \infty} w(x, z, t; \min\{u_0, v^-\}) = v_*(x, z)$$

for $(x, z) \in \mathbb{R}^2$ to prove the theorem. We use $\min\{u_0, v^-\}$ instead of u_0 . Hereafter we assume $u_0 \leq v^-$ without loss of generality. Lemma 3.2 immediately shows

$$w_4(x, z, t) \leq w(x, z, T + t) \leq v_*(x, z) \quad \text{for } (x, z) \in \mathbb{R}^2$$

and then

$$w(x, z, s; w^t) \leq w(x, z, T + t + s) \leq v_*(x, z) \quad \text{for } (x, z) \in \mathbb{R}^2$$

where

$$w^t(x, z) \stackrel{\text{def}}{=} w_4(x, z, t).$$

By this inequality and Lemma 2.2, [14, Lemma 4.1] and by letting t tend to infinity, we obtain

$$w(x, z, s; v_5^\delta) \leq \liminf_{t \rightarrow \infty} w(x, z, T + t + s) \leq \limsup_{t \rightarrow \infty} w(x, z, T + t + s) \leq v_*(x, z)$$

for $(x, z) \in \mathbb{R}^2$, where

$$v_5^\delta(x, z) \stackrel{\text{def}}{=} \max\{v_1(x - \rho\delta, z), v_2(x + \rho\delta, z), v_*(x, z + M + \rho\delta)\}.$$

Since δ can be chosen arbitrarily small, we have

$$w(x, z, s; v_5^0) \leq \liminf_{t \rightarrow \infty} w(x, z, T + t) \leq \limsup_{t \rightarrow \infty} w(x, z, T + t) \leq v_*(x, z)$$

for $(x, z) \in \mathbb{R}^2$. Thus letting $s \rightarrow \infty$ again, we have

$$v_{**}(x, z) \leq \liminf_{t \rightarrow \infty} w(x, z, t) \leq \limsup_{t \rightarrow \infty} w(x, z, t) \leq v_*(x, z).$$

Proposition 3.3 implies that $w(x, z, t)$ converges to $v_*(x, z)$ as $t \rightarrow \infty$. $\square$

REFERENCES

- [1] D. G. Aronson and H. F. Weinberger, Nonlinear diffusion in population genetics, Partial Differential Equations and Related Topics, ed. J. A. Goldstein, Lecture Notes in Mathematics, 446 (1975) 5–49.
- [2] X. Chen, Generation and propagation of interfaces for reaction-diffusion equations, J. Differential Equations, 96, No 1 (1992) 116–141.
- [3] X. Chen, Existence, uniqueness, and asymptotic stability of traveling waves in nonlocal evolution equations, Advances in Differential Equations, 2, No 1 (1997) 125–160.
- [4] P. C. Fife, Dynamics of Internal Layers and Diffusive Interfaces (CBMS-NSF Reg. Conf. Ser. Appl. Math. 53), SIAM, Philadelphia, 1988.
- [5] P. C. Fife and J. B. McLeod, The approach of solutions of nonlinear diffusion equations to travelling front solutions, Arch. Rat. Mech. Anal., 65 (1977) 335–361.
- [6] D. Gilbarg and N.S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer-Verlag, Berlin, 1983.
- [7] F. Hamel, R. Monneau and J.-M. Roquejoffre, Stability of travelling waves in a model for conical flames in two space dimensions, Ann. Scient. Ec. Norm. Sup. 4ème série, t.37 (2004) 469–506.
- [8] F. Hamel, R. Monneau and J.-M. Roquejoffre, Existence and qualitative properties of multi-dimensional conical bistable fronts, Discrete and Continuous Dynamical Systems, 13, No. 4 (2005) 1069–1096.
- [9] F. Hamel, R. Monneau and J.-M. Roquejoffre, Asymptotic properties and classification of bistable fronts with Lipschitz level sets, Discrete and Continuous Dynamical Systems, 14, No. 1 (2006) 75–92.
- [10] Y. I. Kanel', Certain problems on equations in the theory of burning, Soviet. Math. Dokl., 2, (1961) 48–51.

- [11] Y. I. Kanel', Stabilization of solutions of the Cauchy problem for equations encountered in combustion theory, *Mat. Sb. (N.S.)*, 59 (101) (1962) 245–288.
- [12] M. Nara and M. Taniguchi, Stability of a traveling wave in curvature flows for spatially non-decaying initial perturbations, *Discrete and Continuous Dynamical Systems*, 14, No. 1 (2006) 203–220.
- [13] H. Ninomiya and M. Taniguchi, Stability of traveling curved fronts in a curvature flow with driving force, *Methods and Application of Analysis*, 8, No 2 (2001) 429–450.
- [14] H. Ninomiya and M. Taniguchi, Existence and global stability of traveling curved fronts in the Allen-Cahn equations, *J. Differential Equations*, 213, No 1 (2005) 204–233.
- [15] T. Ogiwara and H. Matano, Monotonicity and convergence results in order-preserving systems in the presence of symmetry, *Discrete and Continuous Dynamical Systems*, 5, No 1 (1999) 1–34.
- [16] P. Pelcé, *Dynamics of Curved Fronts*, Academic Press Inc., San Diego, 1988.
- [17] M. H. Protter and H. F. Weinberger, *Maximum Principles in Differential Equations*, Springer-Verlag, Berlin, 1984.
- [18] D. H. Sattinger, Monotone methods in nonlinear elliptic and parabolic boundary value problems, *Indiana University Mathematics Journal*, 21, No 11 (1972) 979–1000.
- [19] D. Terman, Directed graphs and traveling waves, *Trans. Amer. Math. Soc.*, 289, No. 2 (1985) 809–847.

Received May 2005; revised December 2005.

E-mail address: `ninomiya@math.ryukoku.ac.jp`

E-mail address: `masaharu.taniguchi@is.titech.ac.jp`