

論文 / 著書情報
Article / Book Information

Title	Production of spin-controlled RI beams
Author	Yuichi Ichikawa, Hideki Ueno, Yuji Ishii, Takeshi Furukawa, Akihiro Yoshimi, Daisuke Kameda, Hiroshi Watanabe, Nori Aoi, KOICHIRO ASAMI, Dimiter Balabanski, Raphael Chevrier, Jean-Michel Daugas, Naoki Fukuda, Georgi Georgiev, Hironori Hayashi, Hiroaki Iijima, Naoto Inabe, Takeshi Inoue, Masayasu Ishihara, Toshiyuki Kubo, Tsubasa Nanao, Tetsuya Ohnishi, Kunifumi Suzuki, Masato Tsuchiya, Hiroyuki Takeda, Mustafa Rajabali
Journal/Book name	Nature Physics, Vol. 8, No. 12, pp. 918-922
発行日 / Issue date	2012, 12
DOI	http://dx.doi.org/10.1038/NPHYS2457
URL	http://nature.com/nphys/journal/v8/n12/full/nphys2457.html
Note	このファイルは査読前原稿です。 This file is pre-print (pre-refereeing).

Production of spin-controlled rare isotope beams

Y. Ichikawa,^{1,*} H. Ueno,¹ Y. Ishii,² T. Furukawa,³ A. Yoshimi,⁴ D. Kameda,¹ H. Watanabe,¹
N. Aoi,¹ K. Asahi,² D. L. Balabanski,⁵ R. Chevrier,⁶ J. M. Daugas,⁶ N. Fukuda,¹
G. Georgiev,⁷ H. Hayashi,² H. Iijima,² N. Inabe,¹ T. Inoue,² M. Ishihara,¹ T. Kubo,¹
T. Nanao,² T. Ohnishi,¹ K. Suzuki,² M. Tsuchiya,² H. Takeda,¹ and M. M. Rajabali⁸

¹*RIKEN Nishina Center, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan*

²*Department of Physics, Tokyo Institute of Technology,
2-12-1 Oh-okayama, Meguro, Tokyo 152-8551, Japan*

³*Department of Physics, Tokyo Metropolitan University,
1-1 Minami-Ohsawa, Hachioji, Tokyo 192-0397, Japan*

⁴*Research Core for Extreme Quantum World, Okayama University,
3-1-1 Tsushimanaka, Kita, Okayama 700-8530, Japan*

⁵*Institute for Nuclear Research and Nuclear Energy,
Bulgarian Academy of Sciences, BG-1784, Sofia, Bulgaria*

⁶*CEA, DAM, DIF, F-91297 Arpajon, France*

⁷*CSNSM, IN2P3-CNRS, Université Paris-sud, F-91404 Orsay, France*

⁸*Instituut voor Kern- en Stralingsfysica, KU Leuven,
Celestijnenlaan 200D, B-3001 Leuven, Belgium*

(Dated: June 29, 2012)

Abstract

The degree of freedom of spin in quantum systems serves as an unparalleled laboratory where intriguing quantum physical properties can be observed, and the ability to control spin is a powerful tool in physics research. We propose a novel method for controlling spin in a system of rare isotopes which takes advantage of the mechanism of the projectile fragmentation reaction combined with the momentum-dispersion matching technique. The present method was verified in an experiment at the RIKEN RI Beam Factory, in which a degree of alignment of 8% was achieved for the spin of a rare isotope ^{32}Al . The figure of merit for the present method was found to be greater than that of the conventional method by a factor of more than 50.

PACS numbers:

*yuichikawa@phys.titech.ac.jp; Present address: Department of Physics, Tokyo Institute of Technology, 2-12-1 Oh-okayama, Meguro, Tokyo 152-8551, Japan

The immense efforts expended to fully comprehend and control *quantum systems* since their discovery are now entering an intriguing stage, namely the controlling the degree of freedom of spin [1–5]. The case of nuclear systems is not an exception. In recent years, nuclear physicists have been focusing their efforts on expanding the domain of known species in the nuclear chart, which is a two-dimensional map spanned by the axes of N (number of neutrons) in the east direction and Z (number of protons) in the north direction. The key technique used to explore the south eastern (neutron-rich, or negative in isospin T_z) and north western (proton-rich, or positive in T_z) fronts of the map has been the projectile fragmentation (PF) reaction, in which an accelerated stable nucleus is transmuted into an unstable one through abrasion upon collision with a target. Several new facilities for providing rare-isotope (RI) beams by this technique, such as RIBF [6] in Japan, FRIB [7–9] in the United States, and FAIR [10–12] in Europe, have been completed or designed for exploration of the frontiers of the nuclear chart. Beyond such efforts toward exploring the N and Z axes, nuclear spin may be a “third axis” to be pursued. The study reported in the present article concerns the control of the spin orientation of an unstable nucleus produced in an RI beam at such the fragmentation-based RI beam facilities. The ability to control spin, when applied to state-of-the-art RI beams, is expected to provide unprecedented opportunities for research on nuclear structure of species situated outside the traditional region of the nuclear chart, as well as for application in materials sciences, where spin-controlled radioactive nuclei implanted in a sample could serve as probes for investigating the structure and dynamics of condensed matter.

The fragmentation of a projectile nucleus in high-energy nucleus-nucleus collisions is described remarkably well by a simple model that assumes the projectile fragment produced in the PF reaction to be a mere “spectator” of the projectile nucleus; as a spectator, this fragment survives frequent nucleon-nucleon interactions, and the other nucleons (“participants”) are abraded off through the reaction [13], as illustrated in Fig. 1. In the model, the projectile fragment acquires an angular momentum (in other words, a nuclear spin), whose orientation is determined simply as a function of the momentum of the outgoing fragment. Here the degrees of spin orientation of rank one and two, in particular, are referred to as *spin polarization* and *spin alignment*, respectively. This implies a unique relation between the spin orientation and the direction of the removed momentum \mathbf{p}_n , as illustrated in Fig. 1, which can be utilized as an obvious means for producing spin-oriented RI beams. One ad-




FIG. 1: Principle of producing spin orientation in PF reaction. A projectile with an initial momentum \mathbf{p}_0 is incident to the target. In a “participant-spectator” model, the nuclear spin \mathbf{I}_{PF} of the fragment arises from the angular momentum $\mathbf{R}_0 \times (-\mathbf{p}_n)$ with respect to the center of mass of the projectile nucleus, owing to the removal of nucleons that, before collision, were in internal motion in the projectile nucleus (Fermi motion). Here, \mathbf{p}_n is the sum of momenta of the removed nucleons (participants), and \mathbf{R}_0 is the position vector of the participants in the projectile rest frame. Furthermore, the linear momentum \mathbf{p}_{PF} of the fragment is given as $\mathbf{p}_{\text{PF}} = \mathbf{p}_0 - \mathbf{p}_n$. Thus the orientation of the spin $\mathbf{I}_{\text{PF}} = \mathbf{R}_0 \times (\mathbf{p}_0 - \mathbf{p}_{\text{PF}})$ is determined plainly as a function of the momentum \mathbf{p}_{PF} of outgoing fragment. In the figure, the axis of polarization is perpendicular to the reaction plane, while the axis of alignment is parallel to the beam axis. Insets (a), (b) and (c) illustrate cases which produce spin orientation for the fragments in the left wing, center and right wing of the momentum distribution, respectively.

vantage of this method of orienting the fragment spin is that the resulting spin orientation does not depend on the chemical or atomic properties of the RI. However, the method also shows a drawback in the sense that the spin orientation thus produced in the PF reaction tends to be partially or completely attenuated since the fragmentation generally involves the removal of a large number of nucleons from the projectile. This is quite a non-negligible flaw with respect to the yields attainable for spin-oriented beams since high-intensity primary

beams are only available for a limited set of nuclear species, and consequently in most cases RIs of interest must be produced through the removal of a large number of nucleons from the projectile. Accordingly, there has been high demand for a new technique for preventing the attenuation in spin orientation caused by large differences in mass between the projectile and the fragment. In this paper, we present a method for producing highly spin-aligned RI beams by employing a two-step PF process in combination with the momentum-dispersion matching technique.

Figure 2 illustrates three different schemes for producing spin-aligned RI beams, where each scheme uses a different configuration of elements, namely primary and secondary targets and slits for selection. The most basic scheme employs the configuration in (a), in which a nucleus of interest is directly produced from a primary beam through a single occurrence of the PF (a single-step PF reaction). As stated earlier, this scheme suffers from the drawback that the degree of spin alignment tends to be attenuated when the PF involves a large number of nucleons. With the aim to overcome this problem, configuration (b) adopts a two-step PF reaction, where a beam of nuclei produced in the first PF reaction (secondary beam) is used to obtain a beam of the nuclide of interest through a second PF reaction. In particular, using a slit installed at a momentum-dispersive focal plane, the particles forming the secondary beam are chosen to be of a nuclide containing one proton or neutron more than the nuclide of interest. Thus, the target RI beam is produced via a PF reaction in which only one proton or neutron is removed. For the RI beam obtained with this scheme, the spin alignment is expected to be high due to the simplicity of the reaction [14, 15]. We also note that a significant increase in the total production yield is suggested by experiments [16] in in-beam γ -ray spectroscopy. In the scheme presented above, however, the production yield is typically reduced by a factor of $\sim 1/1000$ because the production of target nuclei requires the successive occurrence of two highly particular reactions.

A hint on how to eliminate the disadvantage of the scheme in (b) emerged from the recognition that the quantity that determines the spin alignment is solely the momentum change δp in the process of fragmentation that produces the nuclei of interest, and that the spin alignment is not sensitive to the momentum of the incident nucleus. In scheme (b), δp is selected with the aid of two momentum slits, the first of which is used to select the momentum of the secondary beam and the second slit determines the outgoing momentum of the secondary PF. The tremendous and unnecessary drop in yield is avoided by discarding




FIG. 2: Comparison of three schemes for producing a spin-aligned RI beam of ^{32}Al from a primary beam of ^{48}Ca . The graphs below each scheme represent the typical momentum distribution and the corresponding alignment, with abscissas representing the momentum p of ^{32}Al . (a) Single-step PF method. The ^{32}Al beam is directly produced from ^{48}Ca . Since PF involves a large number of nucleons, the expected spin alignment is small. (b) Two-step PF method. ^{32}Al is produced via an intermediate nucleus ^{33}Al . The expected spin alignment is high, whereas the production yield is low due to the two fold selection with momentum slits. (c) Two-step PF method with dispersion matching. Direct selection of the change in momentum δp in the second PF can be achieved by placing a secondary target in the momentum-dispersive focal plane and a slit in the double-achromatic focal plane, because the momentum spread Δp of the incident beam is compensated for by fulfilling the condition of momentum-dispersion matching. This method yields an intensive spin-aligned RI beam while avoiding cancellation between the opposite signs of spin alignment caused by the momentum spread Δp .

the two-fold selection and introducing a single direct selection of the target δp itself. This is realized by placing a secondary target in the momentum-dispersive focal plane and a slit in the double-achromatic focal plane, as illustrated in Fig. 2 (c). The concept of realizing maximum spectral resolution in momentum loss by compensating for the beam momentum

spread of the incident beam, as executed here, is known as *dispersion matching* in ion optics [17, 18]. The important point of this technique is that the reaction products that acquire equal amounts of momentum change upon the second fragmentation are focused onto a single physical location. The application of this technique to PF-induced spin alignment can prevent the cancellation of opposite signs of spin alignment caused by momentum spread, which secondary beams unavoidably undergo.

The validity of scheme (c) was first tested with the in-flight superconducting RI separator BigRIPS [19] at the RIKEN RIBF facility [6]. The arrangement for the production of spin-aligned RI beams with the present method is shown in Fig. 3. In the reaction at the primary target position F0, ^{33}Al was produced by a PF reaction of a 345-MeV/nucleon ^{48}Ca beam on a ^9Be target with a thickness of 1.85 g/cm^2 , chosen to provide a maximum production yield for the secondary ^{33}Al beam. A wedge-shaped aluminium degrader with a mean thickness of 4.05 g/cm^2 was placed at the first momentum-dispersive focal plane F1, where the momentum acceptance at F1 was $\pm 3\%$. The secondary ^{33}Al beam was introduced to a second wedge-shaped aluminium target with a mean thickness of 2.70 g/cm^2 , placed at the second momentum-dispersive focal plane F5. The ^{32}Al nuclei (including those in isomeric state ^{32m}Al) were produced through a PF reaction involving the removal of one neutron from ^{33}Al . The thickness of the secondary target was chosen such that the energy loss from the target was comparable with the theoretical estimate for the width of the momentum distribution [20] for single-nucleon removal. In the present case, $\sigma_{\text{Goldhaber}} = 90\text{ MeV}/c$, and the momentum width for ^{32}Al was measured to be $\sigma = 80\text{ MeV}/c$ or 0.4% . The ^{32}Al beam was subsequently transported to focal plane F7 whereby the momentum dispersion between F5 to F7 was tuned to be with the same magnitude and opposite sign as that from F0 to F5 (momentum matching), effectively canceling out the momentum dispersion from the site of the first PF reaction to F7. The slit at F7 was used to select a region of momentum change at the second PF as $\delta p/p = \pm 0.15\%$ about the center of relative momentum distribution.

The ^{32}Al beam was stopped in a Cu crystal stopper mounted on the experimental apparatus for time-differential perturbed angular distribution (TDPAD) measurements, which was placed in a focal plane after the achromatic focal plane F7. This apparatus, shown in the inset of Fig. 3, enabled us to determine the spin alignment as well as the g -factor of ^{32m}Al by observing the changes in anisotropy of the de-excitation γ rays emitted from spin-aligned ^{32m}Al in synchronization with the spin precession in the presence of an external magnetic

field.

The degree of spin alignment A was determined from a ratio $R(t)$ defined as

$$R(t) = \frac{N_{13}(t) - \epsilon N_{24}(t)}{N_{13}(t) + \epsilon N_{24}(t)}, \quad (1)$$

where N_{13} (N_{24}) is the sum of the photo-peak count rates at Ge 1 and Ge 3 (Ge 2 and Ge 4), which are two pairs of Ge detectors placed diagonally to each other, as depicted in the inset of Fig. 3, and ϵ denotes a correction factor for the detection efficiency. Theoretically, the $R(t)$ ratio is expressed as a function of t as

$$R(t) = \frac{3A_{22}}{4 + A_{22}} \cos 2(\omega_L t + \alpha), \quad (2)$$

in terms of the rank-two anisotropy parameter A_{22} , which is defined as $A_{22} = AB_2F_2$. Terms with higher ranks were evaluated to be negligible in the present case of ^{32m}Al . Here, A denotes the degree of spin alignment

$$A = \sum_m \frac{3m^2 - I(I+1)}{I(2I-1)} a(m), \quad (3)$$

where $a(m)$ is the occupation probability for magnetic sublevel m , and I the nuclear spin. B_2 is the statistical tensor for complete alignment, and F_2 is the radiation parameter [21]. The parameter ω_L (Larmor frequency) is given by $\omega_L = g\mu_N B_0/\hbar$, where g is the g -factor of ^{32}Al in units of the nuclear magneton μ_N , and α is the initial phase of $R(t)$.

The ^{32}Al nucleus is known to exhibit an isomeric state ^{32m}Al [22] at 957 keV with a half-life of 200(20) ns. The spin and parity of ^{32m}Al have not been fixed among the 4^+ and 2^+ candidates. It is known that ^{32m}Al undergoes de-excitation by E2 transition [23] with emission of γ rays with an energy of 222 keV and subsequently decays in cascade to the ground state by emitting 735-keV γ rays. Figure 4 (a) shows a γ -ray energy spectrum measured with the Ge detectors, where 222-keV de-excitation γ rays are clearly observed as a peak. The time variations $N_{13}(t)$ and $N_{24}(t)$ of the intensities for this peak obtained with detectors pairs Ge 1 - 3 and Ge 2 - 4, respectively, are presented in Fig. 4 (b), in which the corresponding abscissas represent the time difference of the signals at either of the Ge detector pairs relative to the beam particle signal at a plastic scintillator placed in front of the stopper crystal. The $R(t)$ ratio evaluated according to Eq. 1 is shown in Fig. 4 (c).

From the least χ^2 fitting of the theoretical function of Eq. 2 to the experimental $R(t)$ ratio of Eq. 1, we obtained the degree of spin alignment as $A = 8(1)\%$, and the g -factor of

^{32m}Al was determined for the first time to be $g = 1.32(1)$. Also, the spin and parity were assigned to be $I^\pi = 4^+$ through comparison of the g -factor with theoretical calculations. Detailed analysis and extended discussion regarding the ^{32}Al nuclear structure based on the obtained g -factor and spin-parity will be presented elsewhere [24].

A remeasurement of the degree of spin alignment was also performed during the experiment, in which the momentum acceptance in the F5 focal plane was narrowed to be $\pm 0.5\%$, while maintaining other conditions unchanged. This measurement corresponded to the two-step PF reaction without dispersion matching (case (b) in Fig. 2). The degree of spin alignment derived from this measurement, $9(2)\%$, is consistent with the above value obtained with the proposed method, $8(1)\%$, thus confirming that the present method of producing spin-aligned RI beams is valid and performs well.

A supplementary experiment was carried out in order to compare the performance of the present method with that of the single-step method. ^{32}Al was directly produced in a PF reaction of a ^{48}Ca beam on a 4-mm thick Be target. The thickness of the production target was chosen such that the energy loss in the target was comparable with the Goldhaber width [20] (4% in this case), where the momentum acceptance at F1 was set to be $\pm 0.5\%$. As a result, the spin alignment was measured to be less than 0.8% (2σ confidence level). A comparison of the two methods is summarized in Table I. The figure of merit (FOM) for the production of such spin-aligned RI beams should be defined to be proportional to the yield and the square of the degree of alignment. In the measurement with the single-step PF reaction, a primary beam whose intensity was deliberately attenuated by a factor of 1/100 was used in order to avoid saturation in the counting rate at the data acquisition system. Here, the FOM was compared on the basis of actual effectiveness without correction for the attenuation, in which the resulting FOM for the new method was found to be improved by a factor of more than 50. Note that the degree of spin alignment in the single-step PF reaction could not be determined within a measurement time comparable with that of the two-step PF reaction. The superiority in FOM of the new method over the single-step PF reaction method should be even more pronounced for nuclei located farther from the primary beam.

Theoretically, the maximum of the spin alignment for the case of single-nucleon removal from ^{33}Al with a momentum acceptance of $\pm 0.15\%$ is estimated to be 30% in a way similar to that described in [25, 26]. The estimation is based on a model proposed by Hüfner and Nemes [13], where the cross-section for the abrasion of one nucleon leading to a fragment

TABLE I: Comparison of the two-step and single-step PF methods. The isomer to ground-state ratio for the production of ^{32}Al was derived from the γ -ray count, detection efficiency, and number of ^{32}Al beam particles at the final focal plane for each method. The in-flight decay of ^{32m}Al from the production target to the final focal plane was also taken into account.

	Two-step method	Single-step method
Reaction	$^{48}\text{Ca} \rightarrow ^{33}\text{Al} \rightarrow ^{32}\text{Al}$	$^{48}\text{Ca} \rightarrow ^{32}\text{Al}$
Energy	200 MeV/nucleon	345 MeV/nucleon
Target	10-mm thick Be	4-mm thick Be
$\Delta p/p$	$\pm 0.15\%$	$\pm 0.5\%$
Yield of ^{32}Al	2.3 kcps	8.6 kcps (1/100 Att.)
Yield of ^{32m}Al	0.5 kcps	0.9 kcps (1/100 Att.)
Isomer ratio	50(6)%	59(5)%
Alignment	8(1)%	$< 0.8\%$ (2σ)
Measurement time	11.9 h	9.3 h

of substate m with momentum p is proportional to the probability of finding a particle of substate $-m$ with momentum $-p$ at the surface of the target nucleus. The maximum evaluated in this way is in fact four times greater than that obtained experimentally, which may result from de-excitation from higher states populated through the PF reaction, such as the (4^-) [27] and 1^+ [23] states. This suggests that the ability to select the reaction path in populating the state of interest is key to achieving augmented spin alignment. Thus, spin alignment via PF reactions depends strongly on both the reaction mechanism and the nuclear structure. Under these circumstances, the achieved degree of alignment, 1/4 of the theoretical maximum which was obtained despite the situation that the reaction path to the isomeric state was not unique, is rather satisfactory. If we choose a nucleus produced by a unique reaction path, a degree of spin alignment closer to the theoretical maximum might be possible to achieve.

Figure 5 shows the result of simulating the accessibility of unstable nuclei via the two-step PF method (red region) and the conventional method (blue region). Clearly, the adoption of the two-step method drastically expands the set of accessible nuclei in the nuclear chart.

The present method for producing spin-aligned RI beams is applicable to ground and long-lived excited (i.e., isomeric) states of nuclei. Lastly, we discuss the prospects for application of the proposed method, taking as an example research on the nuclear structure of isomers through measurements of their electromagnetic moments. The angular momentum plays a vital role in a nuclear system. Recent studies using RI beams have revealed an unexpectedly abundant occurrence of high-spin isomers in newly explored regions of the nuclear chart [29–31]. Among others, one important mechanism for such isomerism arises from the short-range nature of the attractive two-body interactions in the nucleus, as discussed below. Assuming that an odd- Z , odd- N nucleus consists of an active proton and an active neutron in their respective orbits \mathbf{j}_p and \mathbf{j}_n around a core of spin-parity $J^\pi = 0^+$, there arises a specific order in terms of energy among the individual states formed by the two valence nucleons, owing to the short-ranged attractive interaction between the proton and neutron. Those states, with angular momenta \mathbf{j}_p and \mathbf{j}_n , exhibiting parallel and antiparallel coupling, undergo the most pronounced decrease in energy [32, 33], since the spatial overlap between the proton and neutron wave functions is largest for such a state (coplanar proton and neutron). Thus, the energy $E(J)$ of the state with spin J (where $\mathbf{J} = \mathbf{j}_p + \mathbf{j}_n$ and $\mathbf{J}^2 = J(J + 1)$) varies with J such that $E(J)$ takes a minimum at $J = |j_p - j_n|$, rises as J increases, reaching a maximum at a certain value of J , and finally reaches the second minimum at $J = |j_p + j_n|$ (the stretched spin), as depicted in Fig. 5. In such a situation the spin-stretched state is likely to be an isomer, since it can decay to the lower states (necessarily with considerably lower J) only by hindered γ transitions with high multipolarities (spin-gap isomerism). Spin-gap isomerism is thus a generic phenomenon stemming from the intrinsic nature of nucleon-nucleon interaction, and is considered to occur systematically in the region of medium and heavy nuclei, where there are single-particle orbitals with large angular momenta. The present method can selectively populate and even more importantly spin-align such states via a unique path of removing a proton and a neutron from an even-even nucleus (as the secondary projectile) in the secondary reaction. A spin-stretched state is a pure state in principle, its configuration being interpretable transparently, and therefore the determination of its magnetic moment and/or electric quadrupole moment should provide detailed and reliable information on its nuclear structure. A similar mechanism for producing spin-gap isomers is valid also for some even-even nuclei, such as ^{96}Cd nuclei [34] in which two protons and two neutrons play an active role. Another mechanism by which one can obtain

a significant spin difference between neighboring states is in the regions where unique-parity intruder orbitals come close to the Fermi surface. As an example one can mention the importance of the neutron $g_{9/2}$ orbital for the observed island of isomers around ^{68}Ni [35] or similarly the role played by the neutron $h_{11/2}$ orbital in the Sn isotopes [36]. A number of those isomeric states would have a single-particle character which makes them easy to access by the present two-step PF method using a single-nucleon removal, and also could serve as a very good test ground for the nuclear structure.

In summary, we developed a novel method for RI beam production that allows for concurrent control of the spin of the produced RI. The validity of the method was demonstrated in a test experiment performed at RIBF, RIKEN, in which an RI beam of ^{32}Al with an alignment of 8% was produced from a primary beam of ^{48}Ca , with ^{33}Al as an intermediate product (i.e., the secondary beam particle). The FOM of the method was found to be more than 50 times greater than that of the conventional single-step PF reaction in this particular case. A simulation study conducted for performance evaluation indicated that the present method dramatically broadens the domain of accessible nuclei in the nuclear chart. Such an ability to control spin, when applied to state-of-the-art RI beams, is expected to provide unprecedented opportunities for research on the nuclear structure of species situated outside the traditional region of the nuclear chart, as well as for applications in material research where spin-controlled radioactive nuclei implanted in a sample serve as probes into the structure and dynamics of condensed matter.

Acknowledgments

Experiments were performed under Program No. NP0702-RIBF018 at RIBF, operated by RIKEN Nishina Center and CNS, The University of Tokyo. We thank the RIKEN Ring Cyclotron staff for their cooperation during experiments. Y.I. is grateful to the Special Postdoctoral Researchers Program, RIKEN. This work was partly supported by the JSPS KAKENHI (22340071 and 20532089), and by the JSPS and MAEE under the Japan-France Integrated Action Program (SAKURA).

[1] S. A. Wolf *et al.*, Science **294**, 1488 (2001).

- [2] P. Benioff, Phys. Rev. Lett. **48**, 1581 (1982).
- [3] P. Neumann *et al.*, Science **320**, 1326 (2008).
- [4] Y. K. Kato *et al.*, Phys. Rev. Lett. **93**, 176601 (2004).
- [5] N. Kumada *et al.*, Phys. Rev. Lett. **101**, 137602 (2008).
- [6] Y. Yano *et al.*, Nucl. Instrum. Meth. **B261**, 1009 (2007).
- [7] D. J. Morrissey *et al.*, Nucl. Phys. **A616**, 45 (1997).
- [8] <http://science.energy.gov/np/facilities/frib>
- [9] <http://www.frib.msu.edu>
- [10] W. Henning *et al.*, Nucl. Phys. **A734**, 654 (2004).
- [11] H. Geissel *et al.*, Nucl. Instrum. Meth. **B204**, 71 (2003).
- [12] http://www.gsi.de/portrait/fair_e.html
- [13] J. Hüfner and M. C. Nemes, Phys. Rev. C **23**, 2538 (1981).
- [14] K. Asahi *et al.*, Phys. Lett. **B251**, 488 (1990).
- [15] M. Ishihara *et al.*, Progress in Research, Texas A&M University, 23 (1979).
- [16] K. Yoneda *et al.*, Phys. Lett. **B499**, 233 (2001).
- [17] B. L. Cohen *et al.*, Rev. Sci. Instr. **30**, 415 (1959).
- [18] H. G. Blosser *et al.*, Nucl. Instrum. Meth. **91**, 61 (1971).
- [19] T. Kubo *et al.*, Nucl. Instrum. Meth. **B204**, 97 (2003).
- [20] A. -S. Goldhaber, Phys. Lett. **B53**, 306 (1974).
- [21] H. Morinaga and T. Yamazaki, *In-Beam Gamma-Ray Spectroscopy* (North-Holland, Amsterdam, 1976).
- [22] M. Robinson *et al.*, Phys. Rev. C **52**, R1465 (1996).
- [23] S. Grévy *et al.*, Nucl. Phys. **A734**, 369 (2004).
- [24] Y. Ichikawa *et al.*, in preparation.
- [25] K. Asahi *et al.*, Phys. Rev. C **43**, 456 (1991).
- [26] W. -D. Schmidt-Ott *et al.*, Z. Phys. A **350**, 215 (1994).
- [27] B. Fornal *et al.*, Phys. Rev. C **55**, 762 (1997).
- [28] K. Sümmerer and B. Blank, Phys. Rev. C **61**, 034607 (2000).
- [29] S. Pietri *et al.*, Nucl. Instrum. Meth. **B261**, 1079 (2007).
- [30] C .M. Folden III *et al.*, Phys. Rev. C **79**, 064318 (2009).
- [31] D. Kameda *et al.*, 3rd Joint Meeting of the Nuclear Physics Divisions of the American Physical

Society and the Physical Society of Japan, Hawaii, United States, October 13-17 (2009).

- [32] J. P. Schiffer and W. W. True, *Rev. Mod. Phys.* **48**, 191 (1976).
- [33] R. F. Casten, *Nuclear Structure from a Simple Perspective* (Oxford, New York, 1990.)
- [34] B. S. Nara Singh *et al.*, *Phys. Rev. Lett.* **107**, 172502 (2011).
- [35] R. Grzywacz *et al.*, *Phys. Rev. Lett.* **81**, 766 (1998).
- [36] R. L. Lozeva *et al.*, *Phys. Rev. C* **77**, 064313 (2008).




FIG. 3: Experimental setup. A primary beam of ^{48}Ca ions, preaccelerated to 114 MeV/nucleon, is introduced into SRC (superconducting ring cyclotron), where it is further accelerated to 345 MeV/nucleon. The beam is subsequently adjusted to be incident to a primary target located in the focal plane F0. The projectile fragments are analyzed and selected in the BigRIPS beamline, where F1, F4, F5 and F6 are the momentum-dispersive focal planes and F2, F3 and F7 are the double-achromatic focal planes. In the present experiment, a second PF producing ^{32}Al from ^{33}Al takes place in F5. The TDPAD shown in the inset consists of a Cu crystal stopper, a dipole magnet, Ge detectors, a plastic scintillator and a collimator. The Cu stopper is 3.0 mm in thickness and $30 \times 30 \text{ mm}^2$ in area, and the dipole magnet provides a static magnetic field $B_0 = 0.259 \text{ T}$. ^{32m}Al are implanted into the Cu crystal, and de-excitation γ rays are detected with four Ge detectors located at a distance of 7.0 cm from the stopper and at angles of $\pm 45^\circ$ and $\pm 135^\circ$ with respect to the beam axis. The relative detection efficiency was 35% for one and 15–20% for the other three. A plastic scintillator of 0.1 mm in thickness was placed upstream of the stopper, the signal from which provided the time-zero trigger for the TDPAD measurement.




FIG. 4: Experimental results. (a) Energy spectrum. De-excitation γ rays emitted from ^{32m}Al at 222 keV and 735 keV were observed. (b) Time variations of $N_{13}(t)$ and $N_{24}(t)$ for 222-keV γ rays. Events around $t = 0$ originate from prompt γ rays. (c) $R(t)$ ratio deduced from $N_{13}(t)$ and $N_{24}(t)$, according to Eq. 1 The solid line represents the theoretical $R(t)$ function in Eq. 2 after fitting to the experimental $R(t)$.




FIG. 5: Nuclear chart of “accessible” nuclei. Black boxes indicate stable nuclei, while colored boxes indicate unstable nuclei. Among the latter, red boxes represent those “accessible” with the single-step PF method, and blue boxes represent nuclei which are only accessible with the two-step PF method., where “accessible” here means that the nucleus of interest is producible with its spin aligned and with a production yield sufficiently large to determine the g -factor of its isomeric state with a 5σ confidence level in a one-day beam time. Here, the following conditions are assumed: The degree of spin alignment is 10% for single-nucleon removal from the beam particle, and reduces exponentially to 1% down to 10-nucleon removal, as has been determined empirically; the intensity of the primary beam (rare-gas nuclei or ^{48}Ca) is assumed to be $1\ \mu\text{A}$, which is the designed maximum intensity at RIBF; the cross-sections for the PF reactions are estimated based on parameter sets known as EPAX2 [28], and the cross-section for the secondary PF reaction is assumed to be $1/1000$, as usual; the isomeric to ground state population ratio for the nucleus of interest in the PF reaction is 50%; and the external magnetic field up to 1 Tesla is available for the TDPAD measurement. The inset depicts the mechanism of spin-gap isomerism for an odd- Z , odd- N nucleus consisting of an active proton and an active neutron in their respective orbits j_p and j_n around a core of spin-parity $J^\pi = 0^+$.