

論文 / 著書情報
Article / Book Information

題目(和文)	両親媒性 α ヘリックス細胞膜透過性ペプチドとヘパラン硫酸との相互作用
Title(English)	Interaction of Amphiphilic α -Helical Cell-Penetrating Peptides with Heparan Sulfate
著者(和文)	楊 継
Author(English)	Ji Yang
出典(和文)	学位:博士(工学), 学位授与機関:東京工業大学, 報告番号:甲第9640号, 授与年月日:2014年9月25日, 学位の種別:課程博士, 審査員:三原 久和,和地 正明,櫻井 実,蒲池 利章,小倉 俊一郎
Citation(English)	Degree:Doctor (Engineering), Conferring organization: Tokyo Institute of Technology, Report number:甲第9640号, Conferred date:2014/9/25, Degree Type:Course doctor, Examiner:,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	論文要旨
Type(English)	Summary

(博士課程)
Doctoral Program

論文要旨

THESIS SUMMARY

専攻:	生物プロセス	専攻	申請学位(専攻分野):	博士	(工学)
Department of			Academic Degree Requested	Doctor of	
学生氏名:	Yang Ji		指導教員(主):	三原久和	
Student's Name			Academic Advisor(main)		
			指導教員(副):		
			Academic Advisor(sub)		

要旨(英文 800 語程度)
Thesis Summary (approx.800 English Words)

Cell-penetrating peptides (CPPs) are able to be taken up by cells and can deliver macromolecular cargos. However, the mechanism of this internalization is not yet fully understood. Recent studies have suggested that negatively charged extracellular glycosaminoglycans (GAGs), such as heparan sulfates (HS), can bind cationic CPPs and form cluster upon these peptides binding to them, thereby affecting the efficiency of a wide range of CPPs. Previously Mihara group has screened the cellular uptake (CU) activities of 54 systematically designed amphiphilic α -helical peptides in HeLa cells and the CU activities of 24 designed peptides have been examined in four cell lines. Notably, even a mutation in a single residue significantly alters the CU ability of a peptide. To determine the structure-CU activity relationship of CPPs, four peptides (RF: GLKKLARLFHKLLKLGC-NH₂; EF: GLKKLAELFHKLLKLGC-NH₂; RA: GLKKLARLAHKLLKLGC-NH₂; EA: GLKKLAELAHKLLKLGC-NH₂), which contain a difference in one or two amino acid(s) (i.e., Arg/Glu and Ala/Phe), were chosen from the CPP library to examine their interactions with HS in this study.

First, the interactions of the four CPPs with HS were investigated experimentally by means of circular dichroism (CD), fluorescence spectroscopy, isothermal titration calorimetry (ITC), and dynamic light scattering (DLS). Fluorescence spectroscopy and ITC analysis indicated that the HS-binding affinities abilities of the four CPPs correlated well with their CU activities in HeLa and A549 cells (RF $\approx$ EF > RA > EA), showing that their HS-binding affinities were RF > EF $\approx$ RA >> EA. The stoichiometry of the binding of CPP to HS led to charge neutralization as a molecular principle for binding. The α -helical content of CPP was enhanced when the amphiphilic CPP bound to HS, and the CD analysis yielded an identical order of secondary structure variation of the four designed CPPs in the presence of HS as the order of their HS-binding affinities. Moreover, ITC experiment revealed that their heat capacities were $\Delta C_{P,TAT}^0 > \Delta C_{P,RA}^0 > 0 > \Delta C_{P,EF}^0 > \Delta C_{P,RF}^0$, showing that electrostatic

interactions were dominant in the HS-binding processes of Arg-containing peptides (RA) in comparison to Glu-containing peptides, whereas hydrophobic contributions were the primary mode of interactions of Phe-containing peptides (EF and RF) in comparison to Ala-containing peptides (RA). DLS analysis showed that the order of the HS-clustering abilities of these CPPs was $EF > RF > RA > EA$, which also correlated well with their CU activities in HeLa and A549 cells.

Furthermore, the molecular dynamics (MD) simulations in AMBER 12.0 were performed to reveal the detailed binding and clustering mechanisms of these four peptides with HS. The order of the HS-binding affinities and HS-clustering abilities of the four CPPs, which were obtained from the binding free energies calculated in MD simulations ($\Delta G_{RA} \approx \Delta G_{RF} < \Delta G_{EF} < 0 < \Delta G_{EA}$), almost agreed with the experimental data. Using free energy decomposition analyses, it was also indicated a similar result as given by ITC analysis. In comparison to Glu-containing peptides, the Arg-containing peptides were found to contribute more electrostatic energy to the binding enthalpy, whereas the Phe-containing peptides benefited more from non-electrostatic energy in comparison to the Ala-containing peptides. Moreover, although electrostatic interactions were revealed as the initial impetus of HS-CPP binding, non-electrostatic interactions might be more favorable during the HS-CPP clustering process because of their tolerance to local charge mismatch. Consequently, the CPPs which adopted a non-electrostatic dominant HS-binding mechanism, could be internalized by cells more efficiently.

These data, indicating the structure-HS binding affinity relationship of CPPs, will be helpful to identify and design new efficient CPPs for various functions, such as the delivery of macromolecular drugs to cells.

備考：論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note: Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1 copy of 800 Words (English).