## T2R2 東京科学大学 リサーチリポジトリ Science Tokyo Research Repository

## 論文 / 著書情報 Article / Book Information

題目(和文)	第10族遷移金属錯体の有機配位子移動反応を利用した環状分子および 高分子の合成		
Title(English)			
著者(和文)	吉越裕介		
Author(English)	Yusuke Yoshigoe		
出典(和文)	学位:博士(工学), 学位授与機関:東京工業大学, 報告番号:甲第10217号, 授与年月日:2016年3月26日, 学位の種別:課程博士, 審査員:小坂田 耕太郎,竹内 大介,穐田 宗隆,本倉 健,稲木 信介		
Citation(English)	Degree:, Conferring organization: Tokyo Institute of Technology, Report number:甲第10217号, Conferred date:2016/3/26, Degree Type:Course doctor, Examiner:,,,,		
学位種別(和文)	博士論文		
Category(English)	Doctoral Thesis		
 種別(和文)	論文要旨		
Type(English)	Summary		

## 論 文 要 旨

THESIS SUMMARY

学生氏名: 吉越 裕介 指導教員(主): 小坂田 耕太郎教授	専攻: Department of	13D32100	専攻	申請学位 (専攻分野) : 博士 工学 Academic Degree Requested Doctor of
Student's Name Academic Advisor(main)	学生氏名: Student's Name	吉越 裕介		指導教員(主): Academic Advisor(main) 小坂田 耕太郎教授
指導教員(副): Academic Advisor(sub) 竹内 大介准教授				指導教員(副): 竹内 大介准教授 Academic Advisor(sub)

要旨(英文 800 語程度) Thesis Summary (approx.800 English Words)

The author investigated ligand trasnsfer reactions of aryl platinum(II) complexes and macrocyclization of bridgied-Trögers base ligand in this thesis. Macrocyclization is one of the attractive research fields of synthetic chemistry. The author empolyed ligand transfer reaction of organometallic compounds to synthesize macrocycles because it was easy to control transmetallation by the condition such as temparature, solvent, and ancillary ligand. This doctral thesis is composed of 7 chapters as below.

In chapter 1, the author made an introduction of the thesis, which contained the previous research reports of organic ligand transfer reaction, especially about transmetallation. Also, typical strategies for synthesis of macrocyclic compound such as one-step cyclization, step-wise cyclization, template method, and metathesis method, were described. Among these strategies, the author emphasized that it was the most effective method to form macrocycles by metathesis of the equilibrated bond exchange (e.g. imine metathesis, alkyne metathesis, and alkene metathesis) under thermodynamic control. At the end of chapter 1, the author claimed the reversible transmetallation and similar organometallic reactions would be useful for synthetic application of the macrocyclization.

In chapter 2, the author developed new reversible intramolecular transmetallation of aryl platinum(II) complexes. Diarylplatinum complexs with 1,5-cyclooctadiene (cod) as ancillary ligand were emplyed. Reaction of  $Pt(C_6H_5)_2(cod)$  (1-H<sub>2</sub>) and  $Pt(C_6H_4-4-F)_2(cod)$  $(1-F_2)$  in C<sub>6</sub>D<sub>6</sub> at 50 °C gave a mixture of 1-H<sub>2</sub>, 1-F<sub>2</sub> and Pt(C<sub>6</sub>H<sub>4</sub>-4-F)(C<sub>6</sub>H<sub>5</sub>)(cod) (1-FH) in statical molar ratio (51:49:102). Dissolution of 1-FH in  $C_6D_6$  at 50 °C gave similar mixture of platinum complexes, indicating the reversibility of the ligand exchange reaction. The second-order rate constants ( $k_1$  and  $k_1$ ) of the comproportionation reaction of **1**-H<sub>2</sub> and **1**-F<sub>2</sub> were determined to be  $(6.4 \pm 0.6) \times 10^{-6}$  and  $(2.0 \pm 0.2) \times 10^{-6}$  M<sup>-1</sup>s<sup>-1</sup>, respectively. The kinetic parameters such as  $\Delta S^{\ddagger} = -46 \pm 84$  J mol K<sup>-1</sup>, were calculated by temparature dependance on the  $k_1$  and  $k_2$ , which indicated the associative transition state of two platinum complexes with four membered Pt-C-Pt-C ring. The existence of free cod in the reaction mixture did not affect rate constant significantly that indicated cod ligand coordinated tightly during the transmetallation. DFT studies revealed the effect of ancillary ligand that increased HOMO-LUMO interaction between the platinum complexes to form the above associative transition state.

In chapter 3, the author developed new method of synthesis for macrocycle via the dinuclear macrocyclic complexation by the combination of organometallic reactions. Tröger's base (TB), which has a pair of right-angled aromatic rings, was employed as

bridging ligands of dinuclear complexes. Reaction of Me<sub>3</sub>Sn-C<sub>6</sub>H<sub>3</sub>-C<sub>3</sub>H<sub>6</sub>N<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>-SnMe<sub>3</sub> (TB-(SnMe<sub>3</sub>)<sub>2</sub>) and PtCl<sub>2</sub>(cod) in THF under reflux condition yielded macrocyclic dinuclear Pt(II) complex composed of two Tröger's base ligands. Following reductive elimination was induced by addition of 10 equivalent of PPh<sub>3</sub> yielding macrocyclic dimer of Tröger's base (*cyclic*- $11_2$ ). Based on the NMR spectroscopy of *cyclic*- $11_2$  and DFT calculation, the author revealed cyclic-11<sub>2</sub> had most distorted TBs with angle of arene rings ( $\phi = 65^{\circ}$ ) reported so far. Reaction of HCC-C<sub>6</sub>H<sub>3</sub>-C<sub>3</sub>H<sub>6</sub>N<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>-CCH (TB-(CCH)<sub>2</sub>) and PtCl<sub>2</sub>(dppf) (dppf = 1,1'-bis(diphenylphosphino)) ferrocene) in the presence of CuI in THF/*i*Pr<sub>2</sub>NH at room induced transmetallation vield temparature similar to the macrocycle, *cyclic*-[(dppf)Pt-CC-TB-CC-]<sub>2</sub>. Oxidative addition of TB-I<sub>2</sub> to  $Pd_2(dba)_3$ ·CHCl<sub>3</sub> (dba = dibenzylideneacetone) in the presence of  $Me_2N-C_2H_4-NMe_2$ (TMEDA) gave 11-{Pd(tmeda)}<sub>2</sub>, followed by the transmetallation of bis(arylboronic acid-pinacol ester)-oligo ethylene  $(pin)B-C_6H_4O-(CH_2O)_4-C_6H_4-B(pin)$ glycol. or  $(pin)B-C_6H_4O-(CH_2O)_3-C_6H_4-(OCH_2)_4-OC_6H_4-B(pin)$  (pin = pinacolate), and reductive elimination to yield cyclic-11·14e or cyclic-11·15.

In chapter 4 and 5, the author investigated Lewis acid-promoted polycondensation of ferrocene and aldehyde as well as the characterization of the products which contained linearand cyclic-oligo(ferrocenylene alkylene)s. Reaction of ferocene and 1-alkylaldehyde, such as  $n-C_7H_{15}$ -CHO, Et-CHO and "Bu-CHO, in the presence of BF<sub>3</sub>·OEt<sub>2</sub> in (CH<sub>2</sub>Cl)<sub>2</sub> at 30 °C yielded a mixture of *linear*- and *cyclic*-oligo(ferrocenylene alkylene)s, while polycondensation of ferrocene and arylaldehyde, such as C<sub>6</sub>F<sub>5</sub>-CHO, CF<sub>3</sub>C<sub>6</sub>H<sub>4</sub>-4-CHO and MeC<sub>6</sub>H<sub>4</sub>-4-CHO, gave the linear oligomers exclusively. Their terminal groups of linear oligomers, -ferrocenyl or -CH<sub>2</sub>-aryl group, were charactarized by HRESI-TOMS. Condensations  $\alpha$ -hydroxyalkylferrocene, such 1-ferrocenyloctanol of as and ferrocenylmethanol, gave similar mixture of linear and cyclic oligomer, indicating the propagation of the above polycondensation contained terminal 1-hydroxyalkyl-ferrocenylene group,  $-Fe(C_5H_4)[C_5H_4\{CH(OH)R\}]$ . Polycondensation of diferrocenylmethane with paraformaldehyde dimethylacetal yielded a mixture of oligomers composed of an odd and an even number of ferrocenylene groups, indicating the above polycondensation contained depolymerization process. The trimer synthesized from ferrocene and paraformaldehydedimethylacetal contained 1,1'-, 1,2-, and 1,3-ferrocenylene units, suggesting that the oligomers obtained from alkyl and aryl aldehydes were also composed of these structural units. At the end of chapter 5, the author suggested the mechanism of condensation by consideration of the above result.

In chapter 6, the author described the summery of this thesis and outlooks of the method of the macrocyclization .

In chapter 7, experimental procedures and data of compounds were described in detail.

備考 : 論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note : Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1copy of 800 Words (English).

注意:論文要旨は、東工大リサーチリポジトリ(T2R2)にてインターネット公表されますので、公表可能な範囲の内容で作成してください。 Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).