

論文 / 著書情報
Article / Book Information

題目(和文)	
Title(English)	Study on Candida albicans adhesion in human oral cavity
著者(和文)	NGUYENHoa Thanh
Author(English)	Nguyenthanh Hoa
出典(和文)	学位:博士(学術), 学位授与機関:東京工業大学, 報告番号:甲第12020号, 授与年月日:2021年3月26日, 学位の種別:課程博士, 審査員:山本 直之,小畠 英理,蒲池 利章,小倉 俊一郎,柘植 丈治,梶原 将
Citation(English)	Degree:Doctor (Academic), Conferring organization: Tokyo Institute of Technology, Report number:甲第12020号, Conferred date:2021/3/26, Degree Type:Course doctor, Examiner:,,,,,
学位種別(和文)	博士論文
Type(English)	Doctoral Thesis

TOKYO INSTITUTE OF TECHNOLOGY



Tokyo Tech

Doctoral Thesis

**Study on *Candida albicans* adhesion in
human oral cavity**

Student: Nguyen Thanh Hoa

Supervisor: Professor Susumu Kajiwara

Associate Professor Kanami Orihara

DEPARTMENT OF LIFE SCIENCE

SCHOOL OF LIFE SCIENCES AND TECHNOLOGIES

TOKYO INSTITUTE OF TECHNOLOGY

ACKNOWLEDGEMENTS

First and foremost, I would like to express my deepest gratitude to Professor Kajiwara for his valuable guidance and much needed encouragement throughout this study. It has been a great honor and privilege to study under his supervision. He has provided me not only precious opportunities to work at the Kajiwara and Orihara Laboratory, but also to help me to be able to learn from many Professors especially from Imperial College London and University of Otago, which are very beneficial for my Ph.D's research and my future career.

I am immensely grateful to Associate Professor Orihara for her continuous supports and guidance. Without her help and careful feedback, my study would be incomplete.

My special thanks are dedicated to Dr. Massa Niimi, Dr. Kyoko Niimi and Professor Richard Cannon for providing scientific insights especially in the field of molecular biology and microbiology. Their passion on scientific findings and their critical thinking inspired me to work harder.

I would like to give my sincere thanks to all the past and present members of Kajiwara and Orihara laboratory especially Associate Prof. Chen, Associate Prof. Iwatani, and also Tejima, Ksusha, Agy, Tria for their true friendship, help, co-operation and encouragement. I am also grateful to all brilliant friends I have met in Japan, especially Elena, Ng. Tran, Dzung, Kim, Kana and Nana who believed in me and encouraged me to pursue my study.

To my dear parents, my husband Minh Hai, my daughters Thanh Ha and My An, from the bottom of my heart, I want to thank you all for your unconditional love and support to give me this opportunity to do my best in my Ph.D. study.

Finally, I also express my appreciation to the MEXT scholarship for providing financial supports.

ABSTRACT

Objectives: *Candida albicans* is an opportunistic human pathogen that causes oral candidiasis. The adhesion of this fungus to saliva-coated surfaces is an important early step in the infection process of human oral cavity. A previous study has shown that the *C. albicans* cell wall proteins Bgl2 and Ecm33 may mediate the interaction between the yeast and hydroxyapatite via saliva proteins. This interaction is an initial event in the colonization of the oral cavity, which leads to oral diseases such as candidiasis and denture stomatitis. The aim of this research was to investigate the roles of these two cell wall proteins in the adherence of *C. albicans* to saliva-coated hydroxyapatite as a model for the tooth surface.

Methods: *C. albicans* *BGL2* and *ECM33* null mutants were generated from wild-type strain SC5314 by using the *SATI*-flipper gene disruption method. A novel method for measuring fungal adherence, based on labeling the yeast with the fluorogenic compound Nile Red (NR), was used to investigate the adherence of *C. albicans*. Hydroxyapatite beads coated with various proteins were incubated with *C. albicans* cells to determine the adhesion of the yeast to the beads, and then the cells attached to beads were stained with NR. *C. albicans* gene expression was measured by qRT-PCR.

Results: The adhesion of the *bgl2* Δ and *ecm33* Δ null mutants to saliva-coated hydroxyapatite beads was 76.4 % and 64.8% of the wild-type strain, respectively. Interestingly, the adhesion of the *bgl2* Δ , *ecm33* Δ mutant (87.7 %) was higher than that of either single mutant. qRT PCR analysis indicated that the *ALS1* gene was over-expressed in the *bgl2* Δ ,*ecm33* Δ strain. This result suggested that the double mutant had developed a compensatory response to reduced cell wall integrity. Therefore, I disrupted all three genes (*bgl2* Δ , *ecm33* Δ , *als1* Δ) in *C. albicans* and the triple null mutant should a significantly reduced adherence to the beads, (37.6 %) compared to the wild-type.

Conclusion: This study has demonstrated that the fluorescent compound NR can be used to measure *C. albicans* adherence and the two cell wall proteins Bgl2 and Ecm33 contribute to interaction between *C. albicans* and saliva-coated surfaces. In addition, I propose that *ALS1* gene is associated with the compensatory responses in the fungal cells due to the deletion of both *BGL2* and *ECM33*.

Keywords: *C. albicans*; cell wall protein, cell adhesion, *BGL2*, *ECM33*, *ALS1*, saliva, tooth enamel

TABLE OF CONTENTS

CHAPTER 1. RESEARCH BACKGROUND	1
1.1 INTRODUCTION	1
1.2 Cell wall proteins research.....	2
1.3 Cell wall structure	3
1.3.1 Glucan	4
1.3.2 Chitin	5
1.3.3 Glycoproteins.....	6
1.4 Human saliva	7
1.4.1 Salivary proteins:	8
1.5 Conclusion	10
CHAPTER 2. Developing Nile Red staining assay for detection of <i>C. albicans</i> to adhere to hydroxyapatite (teeth model).....	13
2.1 Introduction.....	13
2.2 Materials and methods:	15
2.2.1 Autofluorescence of NR in different solvents	15
2.2.2 Excitation and Emission spectrum:.....	15
2.2.3 Emission spectrum of NR in the presence of <i>C. albicans</i> SC5314 at different cell numbers	16
2.2.4 The correlation between NR fluorescence and cell numbers	16
2.2.5 The correlation between different concentration of NR fluorescence	16
2.2.6 Time-dependent assay.....	16

2.2.7	Fluorescence of cells in different solvents and time measurement	17
2.2.8	Fluorescence intensity of <i>C. albicans</i> cells in different growth phases	17
2.3	Results and discussion:	18
2.3.1	Effect of different solvent to autofluorescence 1 µg/ml NR.....	18
2.3.2	Excitation and Emission spectrum.....	19
2.3.1	Emission spectrum of NR in the presence of <i>C. albicans</i> SC5314 at different cell numbers	20
2.3.2	The correlation between NR fluorescence and cell numbers	21
2.3.3	The correlation between different concentration of NR fluorescence	23
2.3.4	Time-dependent assay:	24
2.3.5	Fluorescence of cells at different NR concentrations	25
2.3.6	Fluorescence intensity of <i>C. albicans</i> cells in different growth phases	26
2.4	Conclusion	27
CHAPTER 3. The role of Bgl2p and ECM33p in the ability of <i>C. albicans</i> to adhere to saliva-coated oral surface.....		29
3.1	Introduction.....	29
3.2	Materials and methods:.....	33
3.2.1	Strains and growth conditions:	33
3.2.2	Gene disruption and re-insertion.....	33
3.2.3	Transformation of <i>C. albicans</i>	34
3.2.4	Preparation of saliva-coated hydroxyapatite (SHA) beads.....	35

3.2.5	Assay of <i>C. albicans</i> adhesion to SHA beads	35
3.2.6	The susceptibility of <i>C. albicans</i> strains to Calcofluor white (CFW). 36	
3.3	Results and discussion	42
3.3.1	Construction of revertant strain of <i>bgl2Δ/ecm33Δ</i> double mutant	42
3.3.2	Protein concentration	51
3.3.3	Adhesion assay with various protein coated HA beads	52
3.3.4	Cells susceptibility:	54
3.3.5	Adhesion assay:	55
3.3.6	Confocal images:	56
3.4	Conclusion	58
CHAPTER 4. The role of <i>ALS1</i> as a key protein of the complementary system in <i>BGL2</i> and <i>ECM33</i> double mutant of <i>C. albicans</i>		
60		
4.1	Introduction:	60
4.2	Material and Methods:	62
4.2.1	Strains and growth conditions:	62
4.2.2	Gene disruption and re-insertion	63
4.2.3	qRT-PCR	63
4.2.4	Assay of <i>C. albicans</i> adhesion to SHA beads	65
4.2.1	Fluorescence signals of different strains	65
4.2.2	Growth curve of different strains in GSB medium	65
4.2.3	The susceptibility of <i>C. albicans</i> strains to Calcofluor white (CFW). 65	
4.3	Results and discussion	66
4.3.1	<i>ALS1</i> gene was over-expressed in the <i>bgl2Δ, ecm33Δ</i> strain	66

4.3.2	The introduction of ALS1 gene disruptions in mutant strains and investigation of their effect on adherence to SHA beads.....	68
4.3.3	Fluorescence signals of different strains:.....	69
4.3.4	Growth curve of different strains in GSB medium:.....	70
4.3.5	Cells susceptibility.....	71
4.4	Conclusion	72
	CHAPTER 5. Conclusion.....	74
	Ethics approval and consent to participate	75
	REFERENCE.....	78
	Appendix.....	88

ABBREVIATION

Als	Agglutinin-Like-Sequence
AFR	Amyloid-Forming Region
CV	Crystal Violet
CWPs	Cell wall proteins
ECM	Extracellular Matrix
RT-qPCR	Reverse Transcriptase quantitative PCR
YPD	Yeast Extract–Peptone–Dextrose Medium
NT	N-terminal domain
NTC	Nourseothricin
TR	Tandem repeats
PBC	Peptide-Binding Cavity

FIGURES AND CHARTS

Figure 1-1: General steps in invasion process of <i>C. albicans</i>	1
Figure 1-2: Publications on <i>C. albicans</i> from 1964-2019.....	3
Figure 1-3: Transmission electron micrograph showed sections of the <i>C. albicans</i> cell wall and a schematic representation of the major components and the role of those components in immune activation and suppression. ¹⁷	4
Figure 1-4: Overview of salivary gland anatomy.	8
Figure 2-1: Number of publications from 1995 to 2019 concerning the use of NR. Source: Scopus, Keyword: NR, Last updated 9 th March 2020.	14
Figure 2-2: Effect of different solvent in NR staining. (A) Emission spectra of 1 µg/ml NR in KCl buffer containing 10% acetone, (B) Emission spectra of 1 µg/ml NR stock in acetone in KCl buffer containing 10% DMSO, (C) Emission spectra of 1 µg/ml NR in KCl buffer containing 10% DMSO.....	18
Figure 2-3: Excitation spectrum of NR stained <i>C. albicans</i> SC5314.....	19
Figure 2-4: Emission spectrum of NR in the presence of <i>C. albicans</i> SC5314 at different cell numbers.....	20
Figure 2-5: The correlation between NR fluorescence and optical density (600nm).	21
Figure 2-6: The correlation between NR fluorescence and optical density (600nm).	22
Figure 2-7: The correlation between different concentration of NR fluorescence at SC 5314 cells $OD_{600\text{ nm}}=1$ and $OD_{600\text{ nm}}=0.1$	23
Figure 2-8: Time dependent of fluorescence signals at excitation 530nm and emission 570nm.....	24
Figure 2-9: Fluorescence of cells in different solvents and time measurement. SC5314 cells was inoculated in GSB medium to $OD_{600\text{ nm}} = 4-5$, harvested and diluted cells to $OD_{600\text{ nm}} = 1$. Fluorescence intensity of stained cells in different solvents was measured at excitation 530nm. (A) After 5 minute staining, (B) After 30 minutes staining, (C) After 60 minute staining, (D) After 60 minute staining.....	25
Figure 2-10: Fluorescence intensity of NR-stained mid-log phase ($OD_{600\text{ nm}}=1.5$) and stationary phase ($OD_{600\text{ nm}}=4.5$) <i>C. albicans</i> SC5314 cells grown in GSB medium. Cells were harvested and diluted at different $OD_{600\text{ nm}}$ and then stained with 1 µg/ml NR in KCl buffer containing 10% DMSO. Fluorescence signals were measured after 60 minutes of exposure of cells to NR.	26
Figure 3-1: Chromosomal map of <i>C. albicans</i> -specific genes. Chromosomal coordinates for all 65 expressed <i>C. albicans</i> -specific genes were downloaded from the CGD and plotted as a function of	

chromosome location. Asterisks denote clusters of genes at the same loci; indicates the presence of a gene at the telomere.⁶⁴ 29

Figure 3-2: Plasmid construction. TG: Target gene (BGL2. ECM33. ALS1) 31

Figure 3-3: Construction of ECM33 reintegration cassette using PCR-based cloning strategy..... 31

Figure 3-4: (A) Schematic of ECM33 deletion cassette; (B) Lane 1: KpnI and SacI digested plasmid containing ECM33 deletion cassette..... 42

Figure 3-5: (A) Fragment 1 was amplified by primer 5 (ECM33 upstream forward) and primer 24 (ECM33 FRT-Frg1-reverse), (B) Lane 1: Fragment 1 was obtained and estimate the concentration of 100 ng/μl, Lane M: Marker 1kb..... 43

Figure 3-6: Fragment 2 was amplified by primer 25 (ECM33 FRT-Frg2-forward), and primer 20 (ECM33 downstream reverse), (B) PCR result Fragment 2 was purified and estimate the concentration of 100 ng/μl..... 44

Figure 3-7: ECM33 re-insertion cassette amplified by primer 5 and 20, templates are ECM33 fragment 1 and ECM33 fragment 2 (described as the above). B: 1μl loaded of PCR products, M: Marker 45

Figure 3-8: Transformation result of ECM33 re-insertion cassette into *bgl2Δ/ecm33Δ* double mutant ... 46

Figure 3-9: Testing the two clones after ECM33 re-insertion cassette by using primer 11 and primer 14. 47

Figure 3-10: : Testing the two clones after ECM33 re-insertion cassette by using primer 23 and primer 12. 48

Figure 3-11: Testing the clones after ECM33 re-insertion cassette 49

Figure 3-12: Removing SAT1 flipper in clone 1 and clone 2. 50

Figure 3-13: Protein concentration determination of different samples. Protein concentration of each sample was determined by Lowry method using (Bio-Rad DC Protein Assay; cat no. 500-0116). The measurements were carried out according to the manufacturer's instructions (Bio-Rad, Hercules, CA). The amount of 50% saliva diluted in KCl buffer is 1.3 mg/ml, therefore, 0.1% BSA coated HA beads, FBS 1/100 dilution, and skim milk dilution 1/10 in KCl buffer were used as controls for *C. albicans* adherence to hydroxyapatite beads. Results are mean of at three independent experiments. 51

Figure 3-14: Adherence of *C. albicans* SC5314 cells to HA beads coated with various proteins. *C. albicans* cells (1×10^7) grown to stationary phase in GSB were incubated with uncoated, BSA-, FBS-, saliva- or skim milk-coated HA beads (12 mg) for 1 h at 28°C The percentage of the input *C. albicans* cells attached to the HA beads was calculated. Results are the means of at least three independent experiments..... 52

Figure 3-15: *C. albicans* strains deleted in *BGL2* and *ECM33* genes are hypersensitive to CFW. *C. albicans* SC5314, BGL2M4, ECM33M4, and BEM4 cells were serially diluted and spotted onto YPD agar plates and YPD agar plates containing 12.5 µg/mL CFW and incubated at 30 °C for 24 h. 54

Figure 3-16: *C. albicans* strains SC5314 and revertant strain BGL2MK2, ECM33MK2 and BEMK2 restore sensitivity of CFW. Those strains were serially diluted and spotted onto YPD agar plates and YPD agar plates containing 12.5 µg/mL CFW and incubated at 30 °C for 24 h. 54

Figure 3-17: Adherence of wild type cells and homozygous *BGL2* and *ECM33* single and double mutants to SHA beads. The *C. albicans* cells (1×10^7) and SHA beads (12 mg) were incubated for 1 h at 28°C. Percent attached cells is calculated relative to SC5314 which is set to 100%. Results are means of at least three independent experiments. Statistical significance is shown (** $p > 0.01$ *** $p > 0.001$). 55

Figure 3-18: Confocal images of different cell numbers of *C. albicans* SC5314 cells adhered to saliva-coated beads. Z-stack images collected at 83.3 µm sections, total 246 slides were collected to construct the 3D image. Detection was made using excitation channel at 559nm, emission at 636 nm, the red fluorescence indicates yeast cells attached to saliva-coated beads. IMARIS (Bitplane, Inc.) software was used for confocal z-stack 4D and 3D reconstruction and counting the cells numbers. 56

Figure 3-19: Confocal microscopy Z-stack imaging to visualize *C. albicans* to saliva-coated beads. Z-stack images collected at 83.3 µm sections, total 246 slides were collected to construct the 3D image. Detection was made using excitation channel at 559nm, emission at 636 nm, the red fluorescence indicates yeast cells attached to saliva-coated beads. IMARIS (Bitplane, Inc.) software was used for confocal z-stack 4D and 3D reconstruction and counting the cells numbers. 57

Figure 4-1: Schematic of Als protein structure. Line drawing of a representative Als protein, using *C. albicans* Als3 as the example.⁷⁸ The various domains are NT: N-terminal domain, T: T domain, TR: Tandem repeats, and CT: C-terminal domain. 61

Figure 4-2: Primer for qRT PCR..... 64

Figure 4-3: The transcriptional expression of ALS family genes of *C. albicans* wild type and mutant strains. Four strains of *C. albicans* were cultured in YPD medium and then total RNA was isolated for qRT-PCR assay. *C. albicans* ACT1 gene was used as internal control in this experiment. These results are mean of at least three independent experiments. 67

Figure 4-4: The transcriptional expression of ALS family genes of *C. albicans* wild type and mutant strains. Four strains of *C. albicans* were cultured in YPD medium and then total RNA was isolated for qRT-PCR

assay. C. albicans ACT1 gene was used as internal control in this experiment. These results are mean of at least three independent experiments. 67

*Figure 4-5: Adherence of ALS1, BGL2, and ECM33 single, double, and triple null mutants compared to the wild type strain C. albicans SC5314. The C. albicans cells (1×10^7) and SHA beads (12 mg) were mixed for 1 h at 28°C. After the incubation, the beads were washed twice with KCl buffer and all input cells and washed cells were stained with NR in KCl buffer containing 10% DMSO at the final concentration of 1 μ g/ml. Results are means of at least three independent experiments. Statistical significance is shown (** $p > 0.01$ *** $p > 0.001$). 68*

Figure 4-6: Fluorescence intensity of different strains at $OD_{600nm} = 1$ 69

Figure 4-7: Growth curve of different strains in GSB medium 70

*Figure 4-8: The susceptibility of C. albicans strains to Calcofluor white (CFW). SC5314 (wild type), ALS1M4(*als1* Δ), ABM4 (*als1* Δ /*bgl2* Δ), AEM4 (*als1* Δ /*ecm33* Δ), and ABEM4 (*als1* Δ /*bgl2* Δ /*ecm33* Δ) cells were cultured in 5 mL YPD medium overnight at 30°C and diluted in YPD to an OD_{600} of 1. The cells were then serially diluted (10-fold) in YPD medium. Three microliters of each dilution were spotted onto YPD plates containing 12.5 μ g/mL CFW, and incubated at 30°C for 24h. Susceptibility of cells to CFW was indicated by their inability to grow on this medium.. 71*

TABLES

<i>Table 3-1: Strains list</i>	38
<i>Table 3-2: Plasmid list</i>	39
<i>Table 3-3: Primers for BGL2, ECM33, ALS1 deletion and reintegration</i>	40
<i>Table 4: Strain list in this chapter</i>	62

CHAPTER 1. RESEARCH BACKGROUND

1.1 INTRODUCTION

Candida spp. are opportunistic pathogens and some of the most common causes of fungal infections of humans.¹ Although they are considered harmless commensal organisms in healthy people, in immunocompromised individuals such as those with cancer, or who have undergone repeated surgery or the introduction of intravenous catheters, these fungi can become pathogenic. They can cause a variety of mucosal and systemic infections, including pseudomembranous candidiasis, erythematous candidiasis, hyperplastic candidiasis, and candidemia, which can be fatal.²⁻⁴ *Candida albicans* is the most frequent cause of oral candidiasis⁵ and possesses several factors and specific abilities that contribute to its high pathogenicity. These include polymorphism, the yeast-to-hypha morphological transition and biofilm formation. Figure 1-1 shows *C. albicans* biofilms formation in a sequential process including adhesion to host surfaces, induction of hyphal growth, production of extracellular matrix material, and finally dispersion of yeast cells from the biofilm complex.^{6,7} *Candida* cells in a biofilm matrix exhibit high resistance to antimicrobial drugs, and are hidden from the host immune system, and this contributes to the virulence of this fungus.




Figure 1-1: General steps in invasion process of *C. albicans*

C. albicans possesses many surface proteins involved in adhesion that bind to host receptors. These include the ALS protein family, the Sap protein family, Hwp1, Ywp1, Pra1, and Csh1 which have been well studied.⁸⁻¹³ In addition, surface mannoproteins, which constitute the outer layer of the cell wall, are thought to be major candidates for putative adhesins. Therefore, it is important to investigate the functions of cell wall mannoproteins in order to understand the adhesion mechanisms, biofilm formation and virulence of *C. albicans*.^{3,7,14,15}

1.2 Cell wall proteins research

The cell wall has two critical roles: (1) to maintain and protect the integrity of the cell and (2) to interact with the outer environment. The cell surface is also the contact point between *C. albicans* cell and the host surfaces including mucosal surfaces, or medical devices.⁸ Because of this important role, for *C. albicans* only, the cell wall has attracted a consistent focus of attention over the last several decades. Keyword searches for “*C. albicans* cell wall”, “*C. albicans* adhesion” in Pubmed (<https://www.ncbi.nlm.nih.gov/pubmed/>) showed that the number of publications on this topic increased by over 50% in the decade between 2000-2019 (Figure 1-2). One area that has consistent focus is research on *C. albicans* biofilm due to the availability of DNA sequences from Genomic Databases and the study of *C. albicans* cell wall proteins is increasing by utilizing a number of tools.⁸




Figure 1-2: Publications on *C. albicans* from 1964-2019.

The cell wall is a specific and complex structure comprised of glucans, chitin, chitosan, and glycosylated proteins. These components contribute to the cell wall rigidity, and its relationship with virulence, pathogenicity and the interaction with the host's systems.¹⁶ Therefore, in the following sections, the different components of the *C. albicans* cell wall will be discussed generally.

1.3 Cell wall structure

The fungal cell wall locates outside the plasma membrane and mediates all the relationships of the cell with the environment. Therefore, the cell wall allows interaction with the external environment since some of cell wall proteins are adhesins and receptors.¹⁶

The *C. albicans* cell wall contains 90% carbohydrate and 10% protein, Figure 1-3 showed electron microscopy of thin sections of the *C. albicans* cell wall which is comprised of differential abundances of cell wall constituents.¹⁷ There are two main layers in *C. albicans* structure which has different characteristics: (i) outer layer which is comprised of mannan and cell wall proteins; and (ii) the inner layer which is mostly enriched of chitin and polysaccharide matrix.

The *C. albicans* inner cell wall consist of a network including primarily branches of β -1,3- and β -1,6-glucans, chitin and mannoproteins.




Figure 1-3: Transmission electron micrograph showed sections of the *C. albicans* cell wall and a schematic representation of the major components and the role of those components in immune activation and suppression.¹⁷

1.3.1 *Glucan*

Glucan is the most important structural polysaccharide of the *C. albicans* cell wall and represents 50-60% of the dry weight of this matrix. The *C. albicans* inner cell wall constructs a network composed primarily branches of β -1,3- and β -1,6-glucans, to which other components of this structure are covalently linked. The β -1,3 – glucans is synthesized by a number of enzymes located in the plasma membrane named glucan synthases.

The β -1,6-glucans are side chains of variable lengths and distributions, they can form complex structures stabilized by interchain hydrogen bonds. They act as a linking components binding different cell wall proteins to the β -1,3-glucan chitin core through glycosylphosphatidyl inositol (GPI) proteins.¹⁸

1.3.1.1. Bgl2

In fungal cell wall, major polysaccharide components are glucans (including β -1,3 glucan and β -1,6-glucan). As can be seen on Figure 1-3, glucan is synthesized as

linear polymer, therefore, many enzymes are required to form the branching of polysaccharide from the linear polymer. Bgl2 is a manno protein from the surface of *C. albicans* and was identified as major β -1,3-glucosyltransferase.⁸ Bgl2 is categorized as non-covalent and secreted protein in *C. albicans* cell wall proteins matrix. Bgl2 has substrates within the cell wall and remain primarily cell associated.⁸

Since Bgl2 is a β -1,3-glucosyltransferase and a highly conserved cell wall protein, disruption of the *C. albicans* *BGL2* gene interferes cell wall integrity, and the mutant also became attenuated in virulence for mice when compared to its parent strain.¹⁹ Chen *et al.* investigated the morphological role of Bgl2p during the biofilm formation and found that after 24h of cultivation, both the wild-type SC5314 and the *bgl2 Δ /bgl2 Δ* mutant strains exhibited similar amount of biofilm. However, while wild-type strain composed of long filamentous cells and few yeast cells, the *bgl2 Δ /bgl2 Δ* mutant showed excess numbers of yeast cells in addition to some filamentous cells. This studied demonstrated a delay in the transition to filamentous form in the *bgl2 Δ /bgl2 Δ* mutant during initial step of biofilm formation. Further screening the expression level of some well-known transcription factors associated with hypha-associated genes showed that the deletion of *BGL2* gene stimulates a repression of the *CPH2/TEC1* pathway and leads to a delay in the transition of filamentous cells during biofilm formation.^{7,20}

1.3.2 Chitin

The chitin content of the cell wall varies according to the morphological phase of the yeast. It represents 1-2% of the dry weight of yeast cell wall while in filamentous form, it can reach up to 10-20%. The chitin content in the *C. albicans* hyphae wall is three times higher than that of yeast.²¹ Chitin is synthesized from n-acetylglucosamine by chitin synthase, which deposits chitin polymers in the extracellular space. Chitin chains can form tight antiparallel hydrogen-bonded structures associated with high insolubility, cell shape, viability and integrity.⁸

Chitin plays an important role in the interaction of *C. albicans* with the host. Chitin-deficient mutant strains showed attenuated virulence in the immunocompetent and immunosuppressed hosts^{22,23}.

Chitin is important for caspofungin resistance in some *Candida* species, especially *C. albicans*. For example, strains with increased levels of chitin in the *C. albicans* cell wall had higher resistance to echinocandin and showed a systematic *in vivo* infection model of candidiasis.^{16,24}

1.3.3 *Glycoproteins*

As shown in Figure 1-3, the outer layer of *C. albicans* cell wall is packed with mannoproteins. One major class of this layer is glycosylphosphatidyl inositol (GPI)-anchored proteins.¹⁶ N-linked mannans are composed of α -1,6-mannose backbone with α -1,2-oligomannose sidechains. O-linked mannans are found associated with cell wall glycoproteins. Some protein mannosyltransferases are responsible for the first steps in the O-linked mannans biosynthesis, adding a mannose residue to a serine or threonine residue. Some other additional mannoses are synthesized by α -1,2-mannosyltransferases with produces a short α -1,2-mannose chain. The last step results in the addition of an α -1,3-mannoses by α -1,3-mannosyltransferases.²²

Mannan layers are less rigid compared to β -glucans and chitin, therefore, they do not influence the cell shape. However, they have low permeability and porosity which refer to the resistance of yeast cell to antifungal drugs and host defense mechanisms.¹⁷ In addition, since the outer mannan layer covers the inner layers of the cell wall, it is considered to play an important role in the invasion process. Mannans has been described as pathogen associated molecular pattern (PAMP) ligands and many host receptors are proven to participate in their recognition.²⁵⁻²⁷

1.3.3.2. ECM33

Ecm33 is a glycosylphosphatidyl inositol-linked cell wall protein which is required for cell wall integrity and the yeast-to-hyphal transition. It is also required for normal virulence in the mouse model which is hematogenously disseminated candidiasis.²⁸ A mutation of *ECM33* in *Saccharomyces cerevisiae* increases the sensitivity to Calcofluor White²⁸. In *C. albicans*, the filamentation process of *ecm33Δ/ecm33Δ* was delayed in YEPD with serum, and fewer hyphal form were produced.²⁹ In addition, the *ecm33Δ/ecm33Δ* mutant had delayed hypha formation in liquid and solid media. Consistent with these *in vitro* defects, this mutant had severely attenuated virulence in the mouse model of hematogenously disseminated candidiasis²⁸.

Examination of *ecm33Δ/ecm33Δ* mutants by transmission electron microscopy revealed that the cell wall of these strains had an abnormally electron-dense outer mannoprotein layer, which may represent a compensatory response to reduced cell wall integrity.³⁰ Therefore, these results suggested that Ecm33 is required for normal cell wall architecture as well as normal function and expression of cell surface proteins in *C. albicans*³⁰.

Wachtler *et al.* constructed 26 mutants in order to define the genes of *C. albicans* that mediate adhesion, invasion and damage during interaction with oral epithelial cells. One of the most interesting results was the *ecm33Δ* mutant showed significant reduction in adhesion (45%), and also had a higher decreased ability (more than 60%) to invade into epithelial cells, this finding suggested that there is a distinct adhesion and invasion function of *ECM33* gene.³¹

1.4 Human saliva

Within the human body, oral cavity is one of the most heavily colonized by microorganisms. Although there are several antimicrobial systems in human saliva, apparently the saliva also contains substrates for growth of different bacteria and yeast

cells. In the oral cavity, saliva plays an important role in supporting the maintenance or even excessive of microbial colonization³². Studies have showed that salivary constituents can be used by *C. albicans* for growth and survival and adherence to teeth surfaces. In addition, it was found that *C. albicans* can grow in human saliva without addition of glucose in more than 400 hours³².




Figure 1-4: Overview of salivary gland anatomy.

Saliva is secreted from different location of saliva glands, which are the parotid, submandibular and sublingual glands.³³ Under stimulated conditions, the major glands produce about 90% of saliva production, whereas minor glands in the mouth and lips produce the rest of needed saliva.³⁴ The main composition of saliva is a number of electrolytes (sodium, potassium, calcium, magnesium, bicarbonate, and phosphate), proteins (immunoglobulins, enzymes, mucins, bRPs) and nitrogenous products (urea, and ammonia).^{14,34,35} Because of these variety of components, saliva takes part in various interactions including hydration, buffering, tissue coating, mineralization, antimicrobial activities as well as facilitating taste, swallowing and speech.³⁵ The normal pH value of saliva is ranging from 6 to 7, which means that it is slightly acidic.³⁴

1.4.1 *Salivary proteins:*

It is challenging to study the saliva proteome due to the relative composition of salivary proteins varies from person to person as well as physiological conditions. In

general, saliva is a relatively dilute complex mixture of protein including α -amylase, lysozyme, esterase, peroxidase, immunoglobulins, mucins, solid food and trapping of microorganisms.^{33,34,36,37}

- Mucins: Mucins are large O-linked glycosylated proteins which have gel-forming properties.³⁸ There are two carbohydrate rich mucin populations in human saliva, which are MG1 encoded by *MUC5B* gene and MG2 encoded by *MUC7* gene.³⁸ These mucins are synthesized by different cells of the salivary glands, salivary gel-forming mucin *MUC5B* is a large polymeric disulphide bonded protein which can facilitate the dental plaque formation, by contrast, non-gel forming mucin *MUC7* is reported to possess inhibitory activity against a number of bacterial, fungal and viral strains.³⁸⁻⁴⁰

- Alpha-Amylase: this is the most abundant enzyme in saliva, which is mainly formed in the parotid glands and plays an important role in the adhesion of bacteria in the tooth surface by facilitating carbohydrate metabolism.⁴¹

- Proline-rich proteins (PRPs): the proline-rich proteins are described by high number of proline residue (25-42%), and they are categorized into three classes: basic (bPRPs), glycosylated (gPRPs), and acidic (aPRPs).⁴² Acidic PRPs bind to hydroxyapatite, and calcium ions in the cavity and therefore they present in the acquired pellicle.⁴³ When absorbed onto hydroxyapatite, the acidic PRPs are capable of exposing receptors for bacterial attachment.⁴³ The basic PRPs, which are expressed only in parotid saliva, form insoluble complexes with tannin and tannic acid and therefore might play a role in protection against harmful dietary tannins.⁴⁴ Glycosylated PRPs also binds to microorganisms and might facilitate the adherence of bacteria to oral surfaces or clearance from the mouth based on their presence in the tooth pellicle or saliva.⁴⁴

- Statherin: Statherin and its variants exhibit various functions such as binding to hydroxyapatite, and inhibiting crystal growth of calcium phosphate salts. These interactions suggest that it is a potential precursor of the acquired enamel pellicle.⁴³

- Cystatins: Cystatins are widely distributed in human tissues and fluids including saliva, they are natural physiological inhibitors of cysteine proteinase and could protect the oral tissues against degradation of these enzymes.⁴⁵
- Salivary IgA: Immunoglobulin A is the main class of antibodies present in the body secreted fluids including saliva, tears, and mucus of intestines. It is considered the first line defense against harmful environmental factors. The amount of secretory IgA in saliva is stimulated by various factors such as stress and physical activity. IgA is responsible for the agglutination of bacteria as well as inhibition of bacterial adhesion, and neutralization of viruses, toxins, and enzymes produced by microorganism.^{45,46}
- Histatin: histatins are several low molecular-weight histidine rich peptides which are secreted from parotid and submandibular glands. Like PRPs, histatin 1 inhibits crystal growth of calcium phosphate salts but does not inhibit its spontaneous precipitations.⁴³

1.5 Conclusion

Biofilm characteristics of *Candida* species depends on the ability to produce extracellular polymeric substances (EPS), display dimorphic growth, carbon source availability and some other factors.⁴⁷ More importantly, the expression of some adhesins which may proceed to biofilm formation, filamentation displays great complexity and diversity of this pathogenic yeast. Those factors not only have implication in the persistence of colonization and infections but also on antifungal resistance typically found in *C. albicans* cells. Despite many challenges, one of the most promising strategies is to reduce biofilm formation and virulence of *C. albicans* by preventing the adhesion to teeth, medical devices or cell surfaces.

While *C. albicans* is generally considered as a commensal, however, under certain circumstances, it can cause diseases. *C. albicans* is one of the principal causes of opportunistic mycoses worldwide. Adhesion is important for establishing the *C. albicans*-host interaction and may lead to biofilm and antibiotic resistance.

In previous work other researchers discovered that two cell surface mannoproteins, Bgl2 (35 kDa) and another highly glycosylated mannoprotein (97.4 kDa) inhibited the attachment of *C. albicans* cells to bPRP-coated hydroxyapatite beads. The importance of cell wall proteins in biofilm formation is supported by studies showing that mutations of these proteins in *C. albicans* affect cell morphology, cell integrity, adherence to different surfaces, and compromise biofilm development.^{7,48,49} In particular, Bgl2 on the surface of *C. albicans* is required for cell wall maintenance, and disruption of the *C. albicans* *BGL2* gene interferes cell wall integrity⁵⁰ and the dimorphic transition⁵¹ and the null-mutant is attenuated in virulence for mice when compared to its parent strain⁵⁰. The N-terminal sequencing of the 97.4 kDa putative *C. albicans* adhesin gave the amino acid sequence ANXXXLXXAXP which is consistent with the ANNSTLTTATP sequence in Ecm33, (residues 19-29) a glycosylphosphatidylinositol (GPI) – anchored cell wall protein. *C. albicans* Ecm33 is required for normal cell wall integrity and the yeast-to-hyphal transition.²⁸ Deletion of *ECM33* in *C. albicans* leads to delayed hypha formation and severely attenuated virulence in mouse model.²⁸ Therefore, both Bgl2 and Ecm33 are required for normal cell wall architecture as well as normal function and expression of cell surface proteins in *C. albicans*.³⁰ Thus, I hypothesized that the two mannoproteins Bgl2 and Ecm33 mediate specific interactions of *C. albicans* with adhesion receptors in saliva-coated surfaces.

In this research, I investigated the roles of Bgl2 and Ecm33 in the adhesion of *C. albicans* to saliva-coated hydroxyapatite by generating null mutants of these genes and using a novel NR staining assay. I found that the two cell wall proteins Bgl2 and Ecm33 contribute to interaction between *C. albicans* and saliva-coated surfaces. Interestingly, I observed that the deletion of both genes triggered compensatory responses in *C. albicans* cells. In particular, the *ALS1* gene, encoding a major GPI-anchored adhesin, was overexpressed in the *bgl2Δ/ecm33Δ* double disrupted strain, and the deletion of all three

genes caused a significant decrease in the adhesion of *C. albicans* to saliva-coated hydroxyapatite (SHA) beads.

CHAPTER 2. DEVELOPING NILE RED STAINING ASSAY FOR DETECTION OF *C. ALBICANS* TO ADHERE TO HYDROXYAPATITE (TEETH MODEL).

2.1 Introduction

In order to detect and quantify the microbial adhesion to teeth surfaces, several assays have been developed including radiolabeling of the yeast⁵, and/or microscopy using crystal violet (CV) staining¹⁴. In general, radiolabeling is considered high sensitivity and reproducibility compared to quantifying with microscopy or viable cell counts¹⁴. However, all of the above-mentioned assays based on radiolabeling require specialized handling and equipment, laborious work and time consuming. Therefore, a novel alternative method that is fast, cheap and sensitive to detect *C. albicans* adhesion to HA is much needed.

Recently, fluorescence dyes have been extensively used for many cell fluorescence imaging purposes. Although encoded fluorescence proteins have been used widely, many chemical fluorescence dyes are still useful when conjugated to proteins or ligands.⁵² Several fluorescence dyes used for visualizing and quantifying different cells are available. In particular, CFDA (5-Carboxyfluorescein diacetate, acetoxymethyl ester), one of the esterase dependent green fluorogenic probes used for flow cytometric method was developed.⁵³ CFDA provides rapid quantitative information on the vitality and vigor of yeast cell cultures and it was validated by cell sorting and analysis of live, heat killed, and UV-treated yeast.⁵³ However, as CFDA fluorescence signals depend on the amount of esterase, I found that if I use CFDA for quantifying yeast cells, the esterase in human saliva presents in saliva-coated beads will greatly interfere with the adhesion assay results. In 2019, Anna Ligasova *et al.* published a new method using DAPI and Hoechst stains, which based on DNA staining, for the purpose of quantification of fixed adherent cells. In this research, DAPI fluorescence showed excellent linear correlations of cell

quantification based on the measurement of DNA content with the enhancement of SDS.⁵⁴ However, using DAPI staining method for fixed cells requires several washing steps, which will be a major drawback for quantification of yeast cells in the suspension.

In general, fluorescence dyes for *C. albicans* assay about adhesion to saliva-coated HA beads must meet some criteria, including: (i) chemical stability under experiment conditions, (ii) labelling of yeasts to high fluorescence intensity and with uniformity, (iii) relatively short labelling time, (iv) retention of the fluorescence dyes within the labelled yeasts, (v) minimum effects on yeast adherence to HA beads, (vi) controllable nonspecific fluorescence labelling of saliva in the adherence reaction.

Nile Red (NR, 9-diethylamino-5Hbenzo- α -phenoxazine5-5-one), a lipophilic, is one of the most used fluorescent dyes which can bind to intracellular neutral⁵⁵. NR is almost nonfluorescent in water and other polar solvents but shows increased fluorescence signals in nonpolar environments⁵⁶. Due to this characteristic, NR has proven for detection and quantification of intracellular lipid droplets in various organisms including yeasts, algae, filamentous fungi, bacteria and mammalian cells such as macrophages⁵⁵.




Figure 2-1: Number of publications from 1995 to 2019 concerning the use of NR. Source: Scopus, Keyword: NR, Last updated 9th March 2020.

In general, NR is dissolved in an organic solvent that serves as a stain carrier to enhance the permeability of NR into cells. Once NR has diffused into yeast cells, it binds

to the intracellular neutral lipid droplets, resulting in a yellow-gold fluorescence. The magnitude of the fluorescence signals depends on various parameters including stain carriers, cells numbers, cell growth, incubation time and NR concentrations⁵⁷. Therefore, it is crucial to develop a fluorescent staining method using NR for *C. albicans* strains. This method needs to ensure that interactions between NR and intracellular lipids droplets give fluorescence values that are relevant to the calibration from fluorescence to neutral lipid contents and thus, cells numbers. This chapter focuses on the use of NR as a fluorescence dye to quantify *C. albicans* which can be used for further investigation of yeast cells that adhered to saliva-coated HA beads.

2.2 Materials and methods:

2.2.1 *Autofluorescence of NR in different solvents*

NR in acetone stock and DMSO stock were diluted into 3 different solvents to the final concentration of 1 µg/ml, in particular, sample (A) 1 µg/ml NR in KCl buffer containing 10% Acetone, (B) 1 µg/ml NR stock in Acetone in KCl buffer containing 10% DMSO, (C) 1 µg/ml NR in KCl buffer containing 10% DMSO. Two hundred µl of above mentioned solutions were put into different wells in 98 wells plate. Emission spectrum of stained cells in different solvents was measured at fixed excitation 530nm at different time frame.

2.2.2 *Excitation and Emission spectrum:*

SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 5$, yeast cells were harvested by centrifuging at 5000 rpm in 5 minutes and diluted in KCl buffer to $OD_{600nm} = 1$. Cells were then stained with NR in 1 µg/ml in KCl buffer containing 10% DMSO. After 1 hour staining, 200 µl of stained cells and 200 µl of NR 1 µg/ml in KCl containing 10% DMSO were put in to wells in 98 wells plate. Excitation spectrum of stained cells in different solvents was measured at fixed emission 570nm. Emission spectrum of stained cells in different solvents was measured at fixed excitation 530nm.

2.2.3 Emission spectrum of NR in the presence of *C. albicans* SC5314 at different cell numbers

SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 4-5$, yeast cells were harvested by centrifuging at 5000 rpm in 5 minutes and diluted in KCl buffer to OD_{600nm} ranging from 0.1 to 1. Fluorescence spectra of stained cells were measured at excitation 530 nm.

2.2.4 The correlation between NR fluorescence and cell numbers

SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 5$, yeast cells were harvested by centrifuging at 5000 rpm in 5 minutes and diluted in KCl buffer to different OD_{600nm} numbers using ASV11D spectrophotometer. Fluorescence intensity of NR-stained cells at 530 nm excitation and 570 nm emission wavelengths.

2.2.5 The correlation between different concentration of NR fluorescence

SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 5$, yeast cells were harvested by centrifuging at 5000 rpm in 5 minutes and diluted in KCl buffer to $OD_{600nm} = 1$. Cells were then stained with different concentration of NR ranging from 0.25 $\mu\text{g/ml}$ to 4 $\mu\text{g/ml}$ in KCl buffer containing 10% DMSO. After 1 hour staining, 200 μl of stained cells and 200 μl of NR 1 $\mu\text{g/ml}$ in KCl containing 10% DMSO were put in to wells in 98 wells plate. Fluorescence intensity of NR-stained cells at 530 nm excitation and 570 nm emission wavelengths.

2.2.6 Time-dependent assay

SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 4-5$, yeast cells were harvested by centrifuging at 5000 rpm in 5 minutes and diluted in KCl buffer to $OD_{600nm} = 1$. Cells were then stained with different concentration of NR 1 $\mu\text{g/ml}$ in KCl buffer containing 10% DMSO. Fluorescence intensity were measured of NR-stained cells at 530 nm excitation and 570 nm emission wavelengths at different time.

2.2.7 *Fluorescence of cells in different solvents and time measurement*

SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 4-5$, yeast cells were harvested by centrifuging at 5000 rpm in 5 minutes and diluted in different solvents to $OD_{600nm} = 1$. Fluorescence intensity of stained cells in different solvents was measured at excitation 530 nm after 5 minute, 30 minutes, 60 minutes, and 80 minutes staining.

2.2.8 *Fluorescence intensity of C. albicans cells in different growth phases*

SC5314 cells was inoculated in GSB medium to mid-log phase ($OD_{600nm}=1.5$) and stationary phase ($OD_{600nm}=4.5$). Yeast cells were then harvested by centrifuging at 5000 rpm in 5 minutes and diluted in KCl buffer to $OD_{600nm}=1$ Fluorescence signals were measured at excitation wavelength 530 nm and emission wavelength at 570 nm after 60 minutes of staining of cells with 1 μ g/ml NR in KCl buffer containing 10% DMSO.

2.3 Results and discussion:

2.3.1 Effect of different solvent to autofluorescence 1 $\mu\text{g/ml}$ NR




Figure 2-2: Effect of different solvent in NR staining. (A) Emission spectra of 1 $\mu\text{g/ml}$ NR in KCl buffer containing 10% acetone, (B) Emission spectra of 1 $\mu\text{g/ml}$ NR stock in acetone in KCl buffer containing 10% DMSO, (C) Emission spectra of 1 $\mu\text{g/ml}$ NR in KCl buffer containing 10% DMSO.

In general NR is dissolved in organic solvents and serves as a stain carrier to enhance the diffusion of NR into cells. Previous studies suggested that the addition of solvents such as acetone or DMSO at different concentration can be used for enhancing fluorescence signal.¹⁹ Therefore, I investigate the possibility of using DMSO, acetone at different concentrations, and combination of these two solvents at different time of fluorescence acquisition. Fluorescence spectra was measured at excitation 530nm. This result showed that NR gives the best fluorescence intensity when in KCl buffer containing 10% DMSO.

2.3.2 Excitation and Emission spectrum

As NR assay is sensitive, different excitation/emission wavelengths have been used for measurements of various biological samples⁵⁶, therefore, in this experiment, I re-examined the excitation and emission spectrum using Varioskan LUX fluorescence spectrophotometer using 96 well plate.




Figure 2-3: Excitation spectrum of NR stained *C. albicans* SC5314

As shown in Figure 2-3, although 650 nm was the best emission wavelengths for measuring NR fluorescence, but the autofluorescence of this solution is also very high. In contrast, the autofluorescence of emission wavelength at 570 nm was relatively low. Therefore, it would be better to choose 570 nm as emission wavelengths rather than 650nm.

2.3.1 Emission spectrum of NR in the presence of *C. albicans* SC5314 at different cell numbers




Figure 2-4: Emission spectrum of NR in the presence of *C. albicans* SC5314 at different cell numbers.

This figure showed that fluorescence intensities of NR at 570 nm were proportional to the cell number (dashed line) without autofluorescence. Autofluorescence signal (blank) increased above 600 nm, and reached the highest (8 arbitrary unit) at 660 nm. Results are mean of three independent experiments.

2.3.2 The correlation between NR fluorescence and cell numbers




Figure 2-5: The correlation between NR fluorescence and optical density (600nm).

Figure 2-5 showed that the more NR stained cells number tested, the higher the fluorescence values. The fluorescence signals become stable or saturated when the number of cells reached $OD_{600nm}=2$. This phenomenon can be explained that due to the very high density of cells sink into the bottom of the well at OD_{600nm} above 2, the fluorescence signals that can be detected in this assay get saturated. This result indicates that the detection limit for this system would be best with the OD_{600nm} ranging from 0.1-2. Therefore, more tested was done with lower OD_{600nm} value in order to confirm the linear relationship of NR stained *C. albicans* and their fluorescence signals.




Figure 2-6: The correlation between NR fluorescence and optical density (600nm).

More detailed of NR stained at smaller OD_{600nm} was tested. *C. albicans* SC5314 cells were adjusted in different OD_{600nm} ranging from 0.125 to 1. Fluorescence signal was obtained after 60 minutes staining. Figure 2-6 clearly shows that there is a strong correlation and linear relationship between the NR fluorescence and the *C. albicans* cell number. This indicated that the NR fluorescence could be used to quantify the number of *C. albicans* cells adhered to hydroxyapatite beads.

2.3.3 The correlation between different concentration of NR fluorescence




Figure 2-7: The correlation between different concentration of NR fluorescence at SC 5314 cells $OD_{600\text{ nm}}=1$ and $OD_{600\text{ nm}}=0.1$.

Figure 2-7 showed that the more NR added ($>1\ \mu\text{g/ml}$) the fluorescence values becomes stable or saturated. This phenomenon can be explained that quenching effects might occurred because too little neutral lipid being available for staining, in addition, high NR/cells ratios may cause stacking of NR molecules which leads NR more exposed to quenchers in the samples.

2.3.4 Time-dependent assay:




Figure 2-8: Time dependent of fluorescence signals at excitation 530nm and emission 570nm.

To test the stability of fluorescence signals of cells at different time of measurement, OD₆₀₀ was fixed at 1 and stained with NR 1 μ g/ml. Emission fluorescence unit was measured at 570 nm, as showed in Figure 2-8, the fluorescence values becomes stable after 1-5h after staining.

2.3.5 Fluorescence of cells at different NR concentrations




Figure 2-9: Fluorescence of cells in different solvents and time measurement. SC5314 cells was inoculated in GSB medium to $OD_{600nm} = 4-5$, harvested and diluted cells to $OD_{600nm} = 1$. Fluorescence intensity of stained cells in different solvents was measured at excitation 530nm. (A) After 5 minute staining, (B) After 30 minutes staining, (C) After 60 minute staining, (D) After 60 minute staining

In general organic solvent served as a stain carrier to enhance the diffusion of NR into yeast cells. Some studies suggested that addition of solvents such as acetone or DMSO can be used for this purpose.² Therefore, in addition to autofluorescence investigation that was mentioned above, I also investigated the effect of 10% DMSO, 10% acetone and combination of those two solvents at different exposure time on fluorescence intensity. The results also clearly showed that DMSO10% is the best solvent that enhances the fluorescence signals. In addition, this data also strengthen the stability of fluorescence signals after even 1-1.5h of staining.

2.3.6 Fluorescence intensity of *C. albicans* cells in different growth phases




Figure 2-10: Fluorescence intensity of NR-stained mid-log phase ($OD_{600nm}=1.5$) and stationary phase ($OD_{600nm}=4.5$) *C. albicans* SC5314 cells grown in GSB medium. Cells were harvested and diluted at different OD_{600nm} and then stained with $1\mu g/ml$ NR in KCl buffer containing 10% DMSO. Fluorescence signals were measured after 60 minutes of exposure of cells to NR.

The main importance for using fluorescence staining using different fluorescence dyes is that the fluorescence signals depends on the metabolic state of cell populations. As metabolic state can be related to the cell density of the cell population, this may lead to non-linear dependence of the signals on the cell numbers.⁵⁴ In case of using NR staining yeast cells, as can be seen on Figure 2-10, the lipid amount in the stationary cells is different from that in the mid-log cells, however, most of stationary cells have similar lipid amount each other. Therefore, the fluorescence signals always showed excellent linear correlations with cell numbers even in different growth states. Cells at mid-log phase contain less lipid constituents than cells grow at stationary phase, therefore, the fluorescence intensity of NR stained cells at mid-log phase was less than in stationary phase. However, the fluorescence signals were always in a linear dependence with cell populations. This data was in agreement with all other above mentioned results where cells at stationary phase ($OD_{600nm}=4-5$) were used.

2.4 Conclusion

Cell quantification is a common procedure for many assays. In particular, assays used to detect and quantify the microbial adhesion to teeth surfaces have been developed. Some of the most common assays used radiolabeling of the yeast⁵, and/or microscopy using crystal violet (CV) staining¹⁴. Although radiolabeling is considered high sensitivity and reproducibility compared to quantifying using microscopy or viable cell counts¹⁴, all of the above-mentioned assays based on radiolabeling require specialized handling and equipment, laborious work and time consuming.

NR (NR) is a lipophilic fluorescent dye that has been used to stain mammalian cells⁵⁸, bacteria⁵⁹, microalgae,^{56,57,60} and yeast cells⁶¹. I developed a novel adhesion assay based on NR staining of *C. albicans* cells to overcome the limitations of radiolabelling and microscopy techniques. First I optimized and validated the assay. I determined the optimum excitation (530 nm) and emission wavelengths (570 nm) to use to measure NR fluorescence, and ensured there was no autofluorescence of yeast cells at these wavelengths. The best solvent for the NR also investigated, in particular, NR gave the highest fluorescence signal in KCl buffer containing 10% DMSO as staining carrier and the signal remained stable 1 hour after staining. Then, I found that there was a linear relationship between the NR fluorescence and the *C. albicans* cell number (OD₆₀₀ ranging from 0.125 to 1). This indicated that the NR fluorescence could be used to quantify the number of *C. albicans* cells adhered to hydroxyapatite beads.

In this chapter, I have optimized different parameters to used NR as a fluorescence dye to develop a novel method for quantifying *C. albicans* cells adhesions to hydroxyapatite beads. There are many advantages of using NR compared with conventional methods including: (i) robust and reliable staining, (ii) fluorescence signals are stable up to 5 hours after staining, (iii) no washing steps are needed, this is especially important for *C. albicans* adhesion assays as I need to wash cells that do not attached to

HA beads twice, the cell numbers can be small and if staining with fluorescence dyes need washing, human errors might easily occurred, (iv) NR offers a much cheaper and easier solution, and (v) NR has been widely used and it is safe for human.

The method of using NR as a novel fluorescence dye is suitable for cells quantifications which provides high linearity.

CHAPTER 3. THE ROLE OF BGL2P AND ECM33P IN THE ABILITY OF *C. ALBICANS* TO ADHERE TO SALIVA-COATED ORAL SURFACE.

3.1 Introduction

C. albicans is one of the most frequently isolated fungal pathogens of human, it affects immunocompromised patients ranging from premature infants to AIDS patients. Systemic infections due to *C. albicans* attributed mortality rate of 30-50%.⁶² *C. albicans* is always a diploid organism and has 8 sets of chromosome pairs, its genome size is about 16 Mb (haploid). The complete sequence of *C. albicans* are available at <http://www.candidagenome.org/>.^{63,64}




Figure 3-1: Chromosomal map of *C. albicans*-specific genes. Chromosomal coordinates for all 65 expressed *C. albicans*-specific genes were downloaded from the CGD and plotted as a function of chromosome location. Asterisks denote clusters of genes at the same loci; indicates the presence of a gene at the telomere.⁶⁴

In recent years, the free *Candida* Genome Database is established to facilitate the genome sequences research about *C. albicans*.⁶³ Several gene manipulation techniques are available to allow useful genetic approaches to study gene functions in *C. albicans*

including gene disruption, controlled gene expression, protein tagging, gene re-intergration, and overexpression.⁶³ All of the mentioned techniques have the same principles, which are based on homologous recombination between exogenous complementary sequences and genomic sequences in *C. albicans*.⁶³ In general, genes that are not essential for survival in *C. albicans* can be disrupted or replaced by transformation with DNA cassettes containing selectable markers such as auxotrophic or drug-resistant marker.^{63,65}

The *SAT1*-flipper cassette consists of a nourseothricin resistance marker *C. albicans SAT1* gene and the FLP-mediated gene which encodes a site-specific recombinase FLP, and two FLP recombination sites. Target genes disruption plasmids are usually created by subcloning upstream and downstream regions of the target gene onto both sides of the *SAT1*-flipper cassette.⁶³ After that, the linearized of target gene deletion cassette is introduced into *C. albicans* and successful transformant will be selected in YPD medium containing nourseothricin. The transformants which contain *SAT1*-flipper cassette will be inserted into the *C. albicans* genome and the target gene in one allele is deleted accordingly. The *SAT1*-flipper region is the liberated by transferring the transformants to YPD medium, under the regulation of maltose promoter of FLP gene, FLP recombinase expression is activated. To delete the second allele of the target gene, the process is repeated one more time. This method requires many cloning steps and is therefore somewhat laborious, however, it has the advantage of deleting certain genes in most of *C. albicans* strains, including clinical isolates that lack auxotrophic markers.⁶⁶ Based on the same techniques, the re-insertion of target gene cassette can be constructed and used to generate revertant of null mutants.




Figure 3-2: Plasmid construction. TG: Target gene (BGL2, ECM33, ALS1)

Comparing with classical cloning *SAT1*-flipper cassette method, the deletion cassette construction procedures are considered laborious and time consuming due to several cloning steps. In our experiment, in addition to constructing the deletion cassette using classical cloning strategy, I used an alternative method which relies on PCR-based strategies, using long primers (i.e. 45-100 nt) to flank both sides of the re-insertion cassette which contain both upstream region, target gene sequence, *SAT1*-flipper cassette, and downstream region. Although this technique relies heavily on PCR-based cloning strategy, but it is much faster and can eliminate the hurdle of choosing restriction enzymes that are appropriate for cloning.




Figure 3-3: Construction of ECM33 reintegration cassette using PCR-based cloning strategy.

In previous work the other researchers discovered that two cell surface mannoproteins, Bgl2 (35 kDa) and another highly glycosylated mannoprotein (97.4 kDa) inhibited the attachment of *C. albicans* cells to bPRP-coated hydroxyapatite beads. The importance of cell wall proteins in biofilm formation is supported by studies showing that mutations of these proteins in *C. albicans* affect cell morphology, cell integrity, adherence to different surfaces, and compromise biofilm development.^{7,48,49} In particular, Bgl2 on the surface of *C. albicans* is required for cell wall maintenance, and disruption of the *C. albicans* *BGL2* gene interferes cell wall integrity⁵⁰ and the dimorphic transition⁵¹ and the null-mutant is attenuated in virulence for mice when compared to its parent strain⁵⁰. The N-terminal amino acid sequencing of the 97.4 kDa putative *C. albicans* adhesin gave the sequence ANXXXLXXAXP which is consistent with the ANNSTLTTATP sequence in Ecm33, (residues 19-29) a glycosylphosphatidylinositol (GPI) – anchored cell wall protein. *C. albicans* Ecm33 is required for normal cell wall integrity and the yeast-to-hyphal transition.²⁸ Deletion of *ECM33* in *C. albicans* leads to delayed hypha formation and severely attenuated virulence in mouse model.²⁸ Therefore, both Bgl2 and Ecm33 are required for normal cell wall architecture as well as normal function and expression of cell surface proteins in *C. albicans*.³⁰ In this research, I hypothesized that two mannoproteins Bgl2 and Ecm33 mediate specific interactions of *C. albicans* with adhesion receptors in saliva-coated surfaces. This interaction is an initial event in the colonization of human oral cavity, which leads to oral diseases such as candidiasis and denture stomatitis. Therefore, I propose that the deletion of these two genes triggered compensatory responses in the yeast cells.

3.2 Materials and methods:

3.2.1 Strains and growth conditions:

C. albicans strains used in this study were listed in Table 3-1. Yeast cells were maintained in YPD medium (1% [w/v] yeast extract, 2% [w/v] peptone, 2% [w/v] glucose) containing 50 % [v/v] glycerol at -80°C. They were routinely grown in glucose/salts/biotin medium (GSB) (10 [g/l] glucose, 1.0 [g/l] (NH₄)₂SO₄, 2.0 [g/l] KH₂PO₄, 0.05 [g/l] MgSO₄·7H₂O, 0.05 [g/l] CaCl₂·2H₂O; and 0.05 [mg/l] biotin)⁶⁷.

3.2.2 Gene disruption and re-insertion

C. albicans gene sequences were obtained from the Candida Genome Database web site (<http://www.candidagenome.org/cgi-bin/seqTools>). The *bgl2* null mutant (BGL2M4) and the *BGL2*-restored strain (BGL2MK2 from BGL2M4) had been constructed previously by our group using the *SAT1* flipper method.⁷ In the present study, the same method was used to construct the *ecm33* null mutant (ECM33M4). In brief, the *SAT1*-flipper method was used to construct the mutants by using *SAT1*, a nourseothricin resistance marker. An *ECM33* deletion cassette was constructed using the plasmid pSFS2. One *KpnI*-*XhoI* fragment of the upstream region (564 bp) of the *ECM33* gene was amplified from SC5314 genomic DNA with *ECM33* upstream forward and reverse primers. Another *NotI*-*SacI* fragment of the downstream region (555 bp) of the *ECM33* gene was amplified with primers *ECM33* downstream forward and reverse primers. These two fragments were inserted into pSFS2 on both sides of the *SAT1*-flipper. The resulting plasmid was digested with *KpnI* and *SacI* to excise *ECM33* deletion cassette.

A recombinant PCR method was used to construct the revertant of the *ecm33*Δ strain (revertant ECM33MK2 containing a functional *ECM33* gene), and revertant of *bgl2*Δ/*ecm33*Δ strain (revertant BEMK2). Firstly, using SC5314 genomic DNA as a template, a fragment containing *ECM33* and its upstream region was PCR amplified using the *ECM33* upstream forward primer and the *ECM33*-FRT-Frg1-reverse primer (Table 3,

Fig. S7(b)), which contained an overlapping region (15 nucleotides) with the ECM33-FRT-Frg2-forward primer. An ECM33 deletion cassette, which contained the nourseothricin resistance marker *SATI*, the *FLP* recombinase, and the *ECM33* downstream region, was PCR amplified using ECM33 deletion cassette and the ECM33-FRT-Frg2-forward primer and ECM33 downstream reverse primer. These two DNA fragments were gel extracted and purified and used as templates for generating an *ECM33* reinsertion cassette using the ECM33 upstream forward primer and ECM33 downstream reverse primer (Supplementary Figure S7). The resultant *ECM33* re-insertion cassette contained the *ECM33* upstream region, *ECM33* coding region and *SATI*-flipper followed by the *ECM33* downstream region.

3.2.3 Transformation of *C. albicans*

C. albicans SC5314 cells were transformed with DNA fragments using the lithium acetate method⁶⁸ with slight modifications. The overnight-cultured of *C. albicans* SC5314 cells were diluted to 10^7 cells/ml in 50 mL fresh YPD medium and grown at 30°C for two generations (~ 4 h). The cells were collected by centrifuging at 5000 rpm for 5 minutes and washed with distilled water. Washed cells were resuspended in 600 μ l distilled water, and 100 μ l of cell suspension was mixed with transformation mixture (72 μ L 1 M lithium acetate, 100 μ L of 2.0 mg/mL single-stranded carrier DNA, 68 μ L *ECM33* deletion cassette suspension). The transformation mixture was incubated at 30°C with rotatory mixing at 20 rpm for 2 h before heat shocking in a 42°C water bath for 45 min. Cells were harvested by centrifuging at 5000 rpm for 5 minutes and resuspended in 1 mL fresh YPD medium before being incubated for 4 h at 30°C with shaking at 200rpm for recovery. Finally, the cells were collected by centrifuging and resuspended in 200 μ l YPD and, then the cell suspension were spread on YPD plates containing 200 μ g/mL nourseothricin and incubated at 30°C for 1-2 days.

The same procedure was repeated using *ECM33* re-insertion cassette or *BGL2* re-insertion cassette into BEM4 (*bgl2* Δ , *ecm33* Δ) double mutants.

3.2.4 Preparation of saliva-coated hydroxyapatite (SHA) beads

Saliva-coated hydroxyapatite beads were prepared as previously described⁵ with modifications. Unstimulated whole saliva collected from 8 donors was stored on ice, and an equal volume of saliva from each donor was pooled. The pooled saliva was clarified by centrifugation at 3000g for 1 h and the supernatant was diluted with an equal volume of KCl buffer (2 mM KH₂PO₄, 2 mM K₂HPO₄·3H₂O, 5 mM KCl, 1 mM CaCl₂, pH 6.5). Proteinase inhibitor (Protease Inhibitor Cocktail, EDTA free, Nacalai tesque, Japan) was added to the diluted saliva at a final concentration of 1 x protease inhibitor solution to prevent proteolysis of saliva proteins. Prepared saliva solution was aliquoted in glass tubes and stored at -80°C.

Hydroxyapatite beads (Micro-Prep Ceramic Hydroxyapatite Type I 80 µm diameter, Bio-rad) were suspended in KCl buffer, washed with dH₂O 10 times and dried in an oven at 60°C. For each adhesion assay 12 mg beads were placed in a 1.5ml eppendorf and hydrated by incubation in 1 mL KCl buffer at 4°C for 16 h prior to use in an adhesion assay. The KCl buffer was aspirated from the hydrated beads and 1 ml saliva solution (50% [v/v] in KCl buffer) was added. The tube was mixed with end-over-end rotation at 15 rpm at room temperature for 1 h to allow adsorption of saliva proteins to the hydroxyapatite beads. The supernatant was aspirated, and the beads were washed twice with 1 mL KCl buffer. As a negative control, hydroxyapatite beads were incubated in KCl buffer (non saliva-coated beads [NSHA]).

3.2.5 Assay of *C. albicans* adhesion to SHA beads

One colony of yeast cells of SC5314 (wild type), BGL2M4 (*bgl2*Δ), ECM33M4 (*ecm33*Δ), and BEM4 (*bgl2*Δ, *ecm33*Δ) were pre-cultured in 5mL GSB medium overnight at 30°C. Cells were harvested by centrifuging 5000 rpm for 5 minutes and then diluted in KCl buffer to OD₆₀₀ of 1.

SHA beads were incubated with stationary phase cells to determine the adhesion ability of yeast cells to the saliva-coated beads. Yeast cells (1 ml) of OD_{600nm} ranging from 0.05 to 1 using ASV11D spectrophotometer in KCl buffer were incubated with SHA or NSHA beads at room temperature with end-over-end mixing for 1.5 h. After incubation, loosely adhered *C. albicans* cells were removed by washing the beads twice with 1 ml KCl buffer. Cells on the beads were stained with NR (final concentration 1 µg/ml in 10% DMSO) for 1 h, and fluorescence intensity associated with the cells was measured using a Varioskan Lux LL154104 (Thermo Fisher) at 530 nm and 570 nm, excitation and emission wavelength, respectively. The fluorescence intensity was determined using a fluorescence-intensity standard curve generated by mixing NR with various number of yeast cells. (Supplement Figure S6).

*Adhesion ability was calculated by following formula. Adhesion ability of wild type strain SC5314 was considered as 100%.

Adhesion ability (%)

$$= \frac{(\text{Fluorescence intensity of input cells} - \text{Fluorescence intensity of unattached cells})}{\text{Fluorescence Intensity of input cells}} \times 100\%$$

As other reference experiment, 1mg of Skim milk (Nacalai Tesque), 1mg BSA (Nacalai Tesque), and FBS (Gibco) were diluted in KCl buffer and protein concentration of each sample was determined by Lowry method using (Bio-Rad DC Protein Assay; cat no. 500-0116). As the protein concentration of 50% whole saliva in KCl buffer contained 1.3 mg/ml protein, the SM, BSA and FBS and were diluted accordingly to the same amount of protein presented in 50% saliva in KCl buffer, those solution were used to coat HA beads in the same way with saliva-coated hydroxyapatite beads mention above.

3.2.6 The susceptibility of *C. albicans* strains to Calcofluor white (CFW)

SC5314 (wild type), BGL2M4 (*bgl2*Δ), ECM33M4 (*ecm33*Δ), and BEM4 (*bgl2*Δ, *ecm33*Δ) cells were cultured in 5 mL YPD medium overnight at 30°C and diluted in YPD to an OD₆₀₀ of 1. The cells were then serially diluted (10-fold) in YPD medium. Three

microliters of each dilution were spotted onto YPD plates containing 12.5 $\mu\text{g}/\text{mL}$ CFW, and incubated at 30°C for 24h. Susceptibility of cells to CFW was indicated by their inability to grow on this medium.

Table 3-1: Strains list

Strain	Genotype	Source/Reference
SC5314	Wild type strain	69
BGL2M4	SC5314, <i>bgl2Δ::FRT/bgl2Δ::FRT</i>	70
BGL2MK2	SC5314, <i>bgl2Δ::FRT/BGL2Δ::FRT</i>	70
ECM33M4	SC5314, <i>ecm33Δ::FRT/ecm33Δ::FRT</i>	70
ECM33MK2	SC5314, <i>ecm33Δ::FRT/ECM33Δ::FRT</i>	this study
BEM4	SC5314, <i>bgl2Δ::FRT/bgl2Δ::FRT, ecm33Δ::FRT/ecm33Δ::FRT</i>	70
BEMK2	SC5314, <i>bgl2Δ::FRT/BGL2Δ::FRT, ecm33Δ::FRT/ECM33Δ::FRT</i>	this study

Table 3-2: Plasmid list

Name	Description	Source/Reference
pSFS2	Contains SAT1 flipper for constructing the revertant strains	⁶⁵
pSFS2.BGL2	Deletes <i>BGL2</i> gene	this study
pSFS2.ECM33	Deletes <i>ECM33</i> gene	this study
pSFS2.ALS1	Deletes <i>ALS1</i> gene	this study

Table 3-3: Primers for *BGL2*, *ECM33*, *ALS1* deletion and reintegration

Primer name	Sequence 5'-3'
<i>BGL2</i> upstream forward	AAGGTACCGATGTTGCTATAAAATAGTACTCACATGA
<i>BGL2</i> upstream reverse	AACTCGAGATGGGAAGAAGTGGGTATAAGAATG
<i>BGL2</i> downstream forward	AAGCGGCCGCTTTAACACTCAATTCATTTTTTATCGTAACTTGG
<i>BGL2</i> downstream reverse	AAGAGCTCCATGACAATATATTGTGCCAGTTC
<i>ECM33</i> upstream forward	AAGGTACCTCTCCCTCTGAACATTACATCTAG
<i>ECM33</i> upstream reverse	AACTCGAGGAAGGGAAGAAACAAATGGAAAGG
<i>ECM33</i> downstream forward	AAGCGGCCGCGGAACCAACACAAAGAAGTATCCAC
<i>ECM33</i> downstream reverse	AAGAGCTCGGATTGAAAAACAAACGTGCAACTTCG
<i>BGL2</i> check forward	CTGGTGGTAGTGTAATCATTACACA
<i>BGL2</i> check reverse	TCCCATGTTGCCAAATTAACATTGC
<i>ECM33</i> check forward	ACAGAGTTCAAACTTCTACACCATCTG
<i>ECM33</i> check reverse	CATGGAATCACCGAATATGGAATTGC
<i>SAT1</i> check forward	GTGAAGTGTGAAGGGGGAGAT
<i>SAT1</i> check reverse	CTGACGAACAAGCACCTTAGG
on <i>BGL2</i> check reverse	AACACCAAGGTTGAAAGCCAAATCACC
on <i>ECM33</i> check forward	TCATCTTGAAGACTGAGGGTA
<i>ECM33</i> -Check1-R	TCACTGCACACAATCCATCCA
<i>ECM33</i> -Check2-R	GTTGGAAGCAGTAGCGGTGA
<i>SAT1</i> -Check-F	AGTCTCGAACGAAACAGCG
<i>ECM33-FRT-Frg1</i> - reverse	CTAGAAAGTATAGGAACTTCTTAGAATAAAGCAACACCAA
<i>ECM33-FRT-Frg2</i> - forward	TTGGTGTGCTTTATTCTAAGAAGTTCCTATACTTTCTAG
<i>BGL2-FRT-Frg1</i> - Reverse	CTAGAAAGTATAGGAACTTCTTAGTTGAATTTACAGTCAA
<i>BGL2-FRT-Frg2</i> - forward	TTGACTGTAAATTCAACTAAGAAGTTCCTATACTTTCTAG
<i>ALS1</i> upstream forward	AAGGTACCCTCGAATCTGCAATGAAAACG
<i>ALS1</i> upstream reverse	AACTCGAGATTTCAATTGAGAGGAGGAAAGAGC
<i>ALS1</i> downstream forward	AAGCGGCCGCGATCACCTTTTTGGCTTGATCT
<i>ALS1</i> downstream reverse	AAGAGCTCGAAGCGAATGCTAGAAAATGAACTG
<i>ALS1</i> check forward	CTGAAGGGAGATGGGAGAGAAT
<i>ALS1</i> check reverse	GGTCAGGTGTGATCGAGGATA

on *ALS1* check reverse

ACCCAAAACAGCATTCCAAG

3.3 Results and discussion

3.3.1 Construction of revertant strain of *bgl2Δ/ecm33Δ* double mutant

3.3.1.1. Positive control using SAT1 cassette




Figure 3-4: (A) Schematic of *ECM33* deletion cassette; (B) Lane 1: *KpnI* and *SacI* digested plasmid containing *ECM33* deletion cassette

The aim of this experiment is to use the *ECM33* deletion cassette which was constructed by previous student as a positive control for introduction of *ECM33* reinsertion cassette into *bgl2Δ/ecm33Δ* double mutant. Figure 3-4 showed that the *ECM33* deletion cassette which contains *ECM33* upstream region, *SAT1*-flipper, and the *ECM33* downstream region was generated by cutting the deletion cassette using *KpnI* and *SacI*.

3.3.1.2. Amplification of ECM33 fragment 1:




Figure 3-5: (A) Fragment 1 was amplified by primer 5 (*ECM33* upstream forward) and primer 24 (*ECM33* FRT-Frg1-reverse), (B) Lane 1: Fragment 1 was obtained and estimate the concentration of 100 ng/ μ l, Lane M: Marker 1kb

The aim of this experiment was to amplify fragment 1 which contains the *ECM33* coding sequence and the upstream region for re-insertion of *ECM33* into the double mutant *bgl2 Δ /ecm33 Δ* by PCR using genomic *C. albicans* as the template. Figure 3-5 showed that the *ECM33* fragment 1 (2820 bp) was successfully amplified with the concentration around 100 ng/ μ l.

3.3.1.3. Amplification of ECM33 fragment 2:




Figure 3-6: Fragment 2 was amplified by primer 25 (*ECM33* FRT-Frg2-forward), and primer 20 (*ECM33* downstream reverse), (B) PCR result Fragment 2 was purified and estimate the concentration of 100 ng/ μ l

The aim of this experiment was to amplify fragment 2 which contains the *SAT1*-flipper cassette and *ECM33* downstream region for re-insertion of *ECM33* into the double mutant *bgl2 Δ /ecm33 Δ* . The *ECM33* deletion cassette was used as the template for this step. Primer 24 (*ECM33* FRT-Frg1-reverse) and primer 25 (*ECM33* FRT-Frg2-forward) have an overlapping region so that *ECM33* fragment 2 has an 15 nucleotide overlapping region with and *ECM33* fragment 1. Figure 3-6 showed that the *ECM33* fragment 2 (4.7 kb) was successfully amplified with the concentration around 100 ng/ μ l.

3.3.1.4. ECM33 Fragment 1 and Fragment 2 for PCR recombination PCR




Figure 3-7: *ECM33* re-insertion cassette amplified by primer 5 and 20, templates are *ECM33* fragment 1 and *ECM33* fragment 2 (described as the above). B: 1 μ l loaded of PCR products, M: Marker .

The aim of this experiment was to amplify *ECM33* re-insertion fragment which contains (i) *ECM33* upstream region, (ii) *ECM33* coding region, (iii) *SAT1*-flipper cassette, and (iv) *ECM33* downstream region. In this recombination PCR protocol, *ECM33* fragment 1 and *ECM33* fragment 2 were used as two templates for this reaction, and primer 5 and primer 20 was used to amplify the whole *ECM33* re-insertion cassette of 7.5 kb. Figure 3-7 showed that the *ECM33* reinsertion cassette (7.5 kb) was successfully amplified, however, due to the overlapping region of FRT sequence, there was another small band occurred at 3.3kb. The 7.5 kb fragment was then purified, and 0.18 μ g of this cassette was used for the transformation step to re-insert *ECM33* coding region back into *bgl2A/ecm33A*.

3.3.1.5. Transformation result:




Figure 3-8: Transformation result of *ECM33* re-insertion cassette into *bgl2Δ/ecm33Δ* double mutant

Based on the dilution methods, about 9×10^8 cells was plated on each plate of YPD containing $400 \mu\text{g}$ NTC/ml. Using the positive control which was the *ECM33* deletion cassette generated by enzyme digestion method, as the amount of input DNA was around $3.2 \mu\text{g}$, 1048 clones on positive control were obtained. Since recombination PCR method produce much little DNA amount, only $0.18 \mu\text{g}$ DNA of *ECM33* re-insertion cassette was used, 2 clones in *ECM33* re-insertion transformation was obtained. There was no other colonies grew in negative control. As can be seen on Figure 3-8, the transformation efficiency of using high amount of DNA was much higher than the recombination PCR method, however, generation of *ECM33* re-insertion cassette by using conventional cloning methods is often considered time-consuming and laborious. Although the recombination PCR provided a much smaller amount of DNA, the *ECM33* re-insertion cassette could be constructed quickly using only PCR amplification protocols. The next steps were to test the 2 obtained clones in order to confirm the re-insertion of *ECM33* coding region into *bgl2Δ/ecm33Δ* mutant.

3.3.1.6. Confirmation of ECM33 reinsertion (1):




Figure 3-9: Testing the two clones after *ECM33* re-insertion cassette by using primer 11 and primer 14.

Genomic DNA of these two clones mentioned above were extracted, and the region which contained both *ECM33* upstream region and *ECM33* coding region attached to *SATI*-flipper cassette was amplified with primer 11 and 14. As can be seen on Figure 3-9, no band occurred WT and double mutant which showed that only the two obtained clones contained *SATI*-flipper cassette, and the expected band was at exactly 3.4kb. This result showed that *ECM33* reinsertion cassette has been successfully integrated into *bgl2Δ/ecm33Δ* mutant.

3.3.1.7. Confirmation of ECM33 reinsertion (2):




Figure 3-10: : Testing the two clones after ECM33 re-insertion cassette by using primer 23 and primer 12.

The aim of this experiment was to confirm that *SAT1*, the FRT, and *ECM33* downstream region were successfully inserted into *bgl2Δ/ecm33Δ* mutant. Using primer 23 and primer 12, and gDNA of two clones as templates, the results in Figure 3-10 showed that there were no band at WT and *bgl2Δ/ecm33Δ* double mutant which mean *SAT1*-flipper region, and there was an expected band at exactly 0.87 kb from clone 1 and clone 2. This result strongly confirmed that *ECM33* reinsertion cassette has been successfully transformed into *bgl2Δ/ecm33Δ* mutant. *SAT1* gene helped the two clones survive in strong NTC selection pressure.

3.3.1.8. Confirmation of ECM33 reinsertion (3):




Figure 3-11: Testing the clones after *ECM33* re-insertion cassette

The aim of this experiment was to confirm that the whole *ECM33* re-insertion cassette was successfully transformed into *bgl2Δ/ecm33Δ* mutant. Using primer 5 and primer 20, and gDNA of two clones were used as primers and templates by PCR, the results in Figure 3-11 showed that there was a band at 3.3 kb of WT and the two clones which mean *ECM33* upstream region, *ECM33* coding region, and *ECM33* downstream region presented. This result strongly confirmed that *ECM33* reinsertion cassette has been successfully transformed into *bgl2Δ/ecm33Δ* mutant, and *SAT1* gene helped the two clones survive in strong NTC selection pressure. The other bright band at 1.4kb showed that this two transformants still had one allele that lack *ECM33* coding region. The band at 3.3 kb of the two clones showed that some cells had flipped out *SAT1* during the liquid incubation overnight in order to obtain the gDNA. There was one band at 7.5 kb which showed that *ECM33* and *SAT1* cassette has been inserted into *bgl2Δ/ecm33Δ* double mutant. Once again, this result showed that *ECM33* reinsertion cassette has been successfully intergrated into *bgl2Δ/ecm33Δ* mutant, and the *SAT1* region helped the 2 clones survive in strong NTC selection pressure. The next important step was to remove the *SAT1* region by cultivating these strains into YPD medium without NTC.

3.3.1.9. Removing SAT1 flipper in 2 clones:




Figure 3-12: Removing SAT1 flipper in clone 1 and clone 2.

As stated before, the clone 1 and clone 2 were inoculated into YPD plates containing 25 $\mu\text{g/ml}$ Nourseothricin for their smaller colony size compared to Nour^R strains, and then the new clones were inoculated in YPD liquid medium without nourseothricin to allow FLP-mediated excision of *SAT1* flippers. In summary, 5 colonies were selected and these gDNA were extracted and used as templates for different PCR amplification confirmation procedure. As a result, 4 of them showed no *SAT1* region (check by primer 23-12), and those 4 colonies contained *ECM33* coding region (check by primer 11-12). Finally, the sequence of *ECM33* in this revertant strain was confirmed by sequencing and comparing with *ECM33* sequence of WT SC5314.

In summary, *ECM33* was successfully re-inserted into *bgl2 Δ /ecm33 Δ* , the genome of this strain was *bgl2 Δ ::FRT/bgl2 Δ ::FRT/ecm33 Δ ::FRT/ECM33::FRT*. Therefore, the next step is to reinsert *BGL2* coding region into this revertant strain. The same procedure of using recombination PCR method was applied to re-insert *BGL2* coding region into *bgl2 Δ ::FRT/bgl2 Δ ::FRT/ecm33 Δ ::FRT/ECM33::FRT* strain.

3.3.2 Protein concentration




Figure 3-13: Protein concentration determination of different samples. Protein concentration of each sample was determined by Lowry method using (Bio-Rad DC Protein Assay; cat no. 500-0116). The measurements were carried out according to the manufacturer's instructions (Bio-Rad, Hercules, CA). The amount of 50% saliva diluted in KCl buffer is 1.3 mg/ml, therefore, 0.1% BSA coated HA beads, FBS 1/100 dilution, and skim milk dilution 1/10 in KCl buffer were used as controls for *C. albicans* adherence to hydroxyapatite beads. Results are mean of at three independent experiments.

Saliva plays an important role in oral surfaces and promotes the adhesions of *C. albicans* to the teeth. Therefore, it is important to identify the protein concentration in saliva in oral adhesion studies. In addition, to determine whether this promotion of adherence was specific to salivary proteins, other proteins including bovine serum albumin (BSA), fetal bovine serum (FBS), and skim milk (SM) were also recruited. In order to maintain the same concentration of proteins used in each experiment, the protein concentration of 50% saliva in KCl buffer were defined as 1.3 mg/ml, and therefore, the

same amount of protein in other solutions were determined and used for the study. In particular, 0.1% BSA in KCl buffer, 1% FBS dilution, and 10% skim milk with dilution in KCl buffer were used for next experiments.

3.3.3 Adhesion assay with various protein coated HA beads




Figure 3-14: Adherence of *C. albicans* SC5314 cells to HA beads coated with various proteins. *C. albicans* cells (1×10^7) grown to stationary phase in GSB were incubated with uncoated, BSA-, FBS-, saliva- or skim milk-coated HA beads (12 mg) for 1 h at 28°C. The percentage of the input *C. albicans* cells attached to the HA beads was calculated. Results are the means of at least three independent experiments.

All oral surfaces are coated with a salivary protein pellicle. Therefore, it is important to include saliva in oral adhesion studies. In this study, I used hydroxyapatite (HA) beads as a model for the tooth surface and so investigated the influence of human saliva on adherence of *C. albicans* cells to HA beads. HA beads were incubated in human saliva for 1 h (to make SHA beads), washed and then *C. albicans* cells (SC5314) were added. It was found that 93.8 ± 4.9 % of the input yeast cells bound to the SHA beads, whereas only 17.0 ± 9.0 % of SC5314 cells bound to uncoated HA beads (Figure 3-14). This result demonstrated that saliva promoted *C. albicans* adherence HA beads significantly (increased adherence 4.5-fold), and was consistent with our previous results^{5,14}. To determine whether this promotion of adherence was specific to salivary proteins, HA beads were coated with other protein mixture: bovine serum albumin (BSA), fetal bovine serum (FBS), and skim milk (SM). It was found that FBS and BSA, unlike saliva,

did not promote adhesion greatly (by 0.28-fold and 1.74-fold, respectively). Interestingly, SM promoted the adhesion of *C. albicans* to HA beads to almost the same extent as saliva (4.3-fold). From these results, I inferred that saliva, and skim milk, act as receptors for *C. albicans* adherence to HA beads.

3.3.4 Cells susceptibility:




Figure 3-15: *C. albicans* strains deleted in BGL2 and ECM33 genes are hypersensitive to CFW. *C. albicans* SC5314, BGL2M4, ECM33M4, and BEM4 cells were serially diluted and spotted onto YPD agar plates and YPD agar plates containing 12.5 µg/mL CFW and incubated at 30 °C for 24 h.




Figure 3-16: *C. albicans* strains SC5314 and revertant strain BGL2MK2, ECM33MK2 and BEMK2 restore sensitivity of CFW. Those strains were serially diluted and spotted onto YPD agar plates and YPD agar plates containing 12.5 µg/mL CFW and incubated at 30 °C for 24 h.

To investigate the impact of gene disruptions on *C. albicans* cell wall integrity, I tested the sensitivity of null mutants to calcofluor white (CFW), a chemical which interferes with cell wall construction.⁷¹ *C. albicans* SC5314 (wild type), BGL2M4

(*bgl2*Δ), ECM33M4 (*ecm33*Δ), and BEM4 (*bgl2*Δ, *ecm33*Δ) cells were serially diluted and inoculated on YPD agar and on YPD agar containing CFW (12.5 μg/mL), and incubated for 24 h at 30°C. All of the null mutants were more susceptible to CFW than the wild type parental strain (Figure 3-15). BEM4 (*bgl2*Δ, *ecm33*Δ) showed the highest CFW sensitivity. This result indicated that when these cell surface mannoproteins are not expressed there is a decrease in the integrity of the *C. albicans* cell wall.

3.3.5 Adhesion assay:




Figure 3-17: Adherence of wild type cells and homozygous *BGL2* and *ECM33* single and double mutants to SHA beads. The *C. albicans* cells (1×10^7) and SHA beads (12 mg) were incubated for 1 h at 28°C. Percent attached cells is calculated relative to SC5314 which is set to 100%. Results are means of at least three independent experiments. Statistical significance is shown (** $p > 0.01$ *** $p > 0.001$).

To determine whether Bgl2 or Ecm33 might be adhesins involved in adhesion to saliva-coated HA beads, the adherence of *C. albicans* null mutant strains to SHA beads was compared to that of the parental strain SC5314. The null mutants with the genes restored, BGL2MK2, ECM33MK2, and BEMK2 were used in adherence assays as controls. Deletion of both alleles on either the *BGL2* or the *ECM33* gene reduced *C. albicans* adherence by more than 20% compared to the adherence of the wild type strain. Returning the *BGL2* or the *ECM33* gene to the null mutants (BGL2MK2, ECM33MK2) restored their ability to bind to SHA beads (Figure 3-17). These results suggested that both Bgl2 and Ecm33 confer the ability to adhere to SHA beads. Surprisingly, I found

that disruption of the *ECM33* gene in the *BGL2* null mutant did not reduce adhesion further but in fact increased adhesion to a level that was not significantly different to that of the wild type (Figure 3-17; BEM4). Returning the *BGL2* and *ECM33* genes to the double disruptant BEM4 did not alter adhesion to SHA beads significantly (Figure 3-17; BEMK2). This increase in adherence when two potential adhesins had been deleted led me to hypothesize that there was a compensatory induction of expression of another adhesin in the *bgl2Δ, ecm33Δ* double mutant.

3.3.6 Confocal images:

In order to visualize the adhesion properties of *C. albicans* to saliva-coated beads, confocal images of different cell numbers of *C. albicans* SC5314 cells adhered to saliva-coated beads were obtained. As can be seen on Figure 3-18, the higher the OD_{600nm} number of cells added to the adhesion assay, the more cells adhered to the beads surfaces.




Figure 3-18: **Confocal microscopy Z-stack imaging to visualize *C. albicans* adhering to saliva-coated beads.** The *C. albicans* SC5314 cells that attached to hydroxyapatite beads were stained with NR. Confocal images were captured with a Zeiss LSCM780 laser scanning microscope. Scale-bar: 20 μm . (A) Bead that had been incubated with *C. albicans* SC5314 cells at OD_{600nm} = 1, (B) Bead that had been incubated with *C. albicans* SC5314 cells at OD_{600nm} = 0.1.

Theoretically, IMARIS (Bitplane, Inc.) software could be used for directly counting the numbers of cells that attached to each saliva-coated beads. This method

could be the one of the most accurate and direct way to quantitatively identify how many *C. albicans* cells interact with saliva-coated HA beads or other protein-coated HA beads. However, this methods is time consuming and requires a great deal of memory on disks. Therefore, quantitatively defining the number of cells that adhered to saliva coated beads surfaces by measuring the fluorecence signals of the unattached cells and total input cells gives more robust results. More detailed steps for constructing the z-stack 3D images was described in Figure 3-18.




Figure 3-19: Confocal microscopy Z-stack imaging to visualize *C. albicans* to saliva-coated beads. Z-stack images collected at 83.3 μm sections, total 246 slides were collected to construct the 3D image. Detection was made using excitation channel at 559nm, emission at 636 nm, the red fluorecence indicates yeast cells attached to saliva-coated beads. IMARIS (Bitplane, Inc.) software was used for confocal z-stack 4D and 3D reconstruction and counting the cells numbers.

3.4 Conclusion

C. albicans is an opportunistic pathogen responsible for 50-90% of oral candidiasis although it is considered commensal microorganism in healthy individuals³⁷. In addition, it was found that *C. albicans* can grow in human saliva without addition of glucose in more than 400 hours⁷². The surface mannoproteins, which constitute the outer layer of cell wall in *C. albicans*, are believed to be the most likely candidates for putative adhesins. Previous studies have shown that two *C. albicans* cell wall proteins, Bgl2 and Ecm33, may mediate the interaction between the yeast and human teeth hydroxyapatite via saliva in human oral cavity⁷. This interaction is an initial event in the colonization of the oral cavity, which leads to oral diseases such as candidiasis and denture stomatitis yeast. The aim of this chapter was to investigate the roles of these two cell wall proteins Bgl2, Ecm33 in the adherence of *C. albicans* to saliva-coated hydroxyapatite as a model for the tooth surface. Since, *C. albicans* *BGL2* and *ECM33* null mutants have been generated by previous students, these revertant strains of BEMK2 and ECM33MK2 were generated from *bgl2Δ/ecm33Δ* by using the *SAT1*-flipper gene disruption method. The novel method for measuring *C. albicans* adherence, based on labeling the yeast with the NR, was used to investigate the adherence of *C. albicans*. Labelling cells using NR is an effective way to count the cell and either by undirect method by quantifying the unattached cells or by direct method using 3D confocal images studies.

Hydroxyapatite beads coated with various proteins were incubated with *C. albicans* cells to determine the adhesion of the yeast to the beads, and then the cells attached to beads were stained with NR. The adhesion of the *bgl2Δ* and the *ecm33Δ* null mutants to saliva-coated hydroxyapatite beads was 76.4 % and 64.8% of the wild-type strain, respectively. Interestingly, the adhesion of the *bgl2Δ, ecm33Δ* mutant (87.7 %) was higher than that of both single mutants. This result suggested that the double mutant had developed a compensatory response to reduced cell wall integrity. In depth investigation

about the mechanism and possible compensatory phenomenon was continued in the next chapter.

CHAPTER 4. THE ROLE OF *ALS1* AS A KEY PROTEIN OF THE COMPLEMENTARY SYSTEM IN *BGL2* AND *ECM33* DOUBLE MUTANT OF *C. ALBICANS*

4.1 Introduction:

In the oral cavity, saliva performs a number of functions. It contains antimicrobial factors such as lysozyme, histatins, and amylase (to begin the breakdown of starch), but it also contains several nutrients that microorganisms can metabolise. Thus, saliva enables microbial colonization of the mouth and it has been reported that *C. albicans* could utilize salivary constituents for the growth and the colonization of tooth surfaces.^{14,32} Salivary proteins adsorb to oral surfaces to form the acquired pellicle. The main component of human tooth enamel is hydroxyapatite and the adsorption of salivary proteins to hydroxyapatite can promote microbial adhesion. Salivary basic proline-rich proteins (bPRP) within the pellicle are thought to act as main receptors for the adherence of *C. albicans*.^{73,74,19}

C. albicans possesses many surface proteins involved in adhesion that bind to host receptors. These include the ALS protein family, the Sap protein family, Hwp1, Ywp1, Pra1, and Csh1 which have been well studied.⁸⁻¹³ In addition, surface mannoproteins, which constitute the outer layer of the cell wall, are thought to be major candidates for putative adhesins. Therefore, it is important to investigate the functions of cell wall mannoproteins in order to understand the adhesion mechanisms, biofilm formation and virulence of *C. albicans*.^{3,7,14,15}

ALS family and adhesion

The *ALS* (agglutinin-like-sequence) gene family of *C. albicans* encodes a large cell surface glycoprotein containing three domains structure that are critical in the process of adhesion to host surfaces. The *ALS* family includes eight genes, from *ALS1* to *ALS7* and *ALS9*, whereas *ALS3* and *ALS8* are the same gene but reside at separate loci. In *C. albicans*,

ALS genes are found on three different chromosomes: chromosomes 3 (*ALS6* and *ALS7*), R (*ALS3*), and 6 (*ALS1*, *ALS2*, *ALS4*, *ALS5*).^{14,27,75-77} Each Als protein contains a relatively conserved N-terminal domain, a central domain consisting of tandem copies of a highly conserved 108-bp unit of a repeated motif, and a serine-threonine-rich C-terminal domain that is relatively variable across the family (Figure 4-1).⁸ The tandem repeats is a potential site for allelic variation, which occur within the same cells with differences in the two alleles borne by the diploid cell or between strains. For example, in the wild type *C. albicans* SC5314, the larger alleles of *ALS1*, *ALS2*, *ALS4*, *ALS5*, and *ALS9* are on one chromosome, while smaller alleles are on the other. The last two domains have multiple glycosylation sites that are considered to lead N-terminal binding domain from the attachment site.



Figure 4-1: Schematic of Als protein structure. Line drawing of a representative Als protein, using *C. albicans* Als3 as the example.⁷⁸ The various domains are NT: N-terminal domain, T: T domain, TR: Tandem repeats, and CT: C-terminal domain.

In the above chapter, I investigated the roles of Bgl2 and Ecm33 in the adhesion of *C. albicans* to saliva-coated hydroxyapatite by generating null mutants of these genes and using a novel NR staining assay. I found that the two cell wall proteins Bgl2 and Ecm33 contribute to interaction between *C. albicans* and saliva-coated surfaces. Interestingly, I observed that the deletion of both genes triggered compensatory responses in *C. albicans* cells. In this chapter, the *ALS* protein family, encoding a major GPI-anchored adhesin, was investigated whether there was a connection between the roles of those proteins in the adhesion ability of *C. albicans* to saliva-coated HA beads.

4.2 Material and Methods:

4.2.1 Strains and growth conditions:

C. albicans strains used in this study were listed in Table 3-1. Yeast cells were maintained in YPD medium (1% [w/v] yeast extract, 2% [w/v] peptone, 2% [w/v] glucose) containing 50 % [v/v] glycerol at -80°C. They were routinely grown in glucose/salts/biotin medium (GSB) (10 [g/l] glucose, 1.0 [g/l] (NH₄)₂SO₄, 2.0 [g/l] KH₂PO₄, 0.05 [g/l] MgSO₄·7H₂O, 0.05 [g/l] CaCl₂·2H₂O; and 0.05 [mg/l] biotin)⁶⁷.

Table 4: Strain list in this chapter

Strain	Genotype	Source/Reference
SC5314	Wild type strain	69
ALS1M4	SC5314, <i>als1Δ::FRT/als1Δ::FRT</i>	79
ABM4	SC5314, <i>bgl2Δ::FRT/bgl2Δ::FRT, als1Δ::FRT/als1Δ::FRT</i>	79
AEM4	SC5314, <i>als1Δ::FRT/als1Δ::FRT, ecm33Δ::FRT/ecm33Δ::FRT</i>	79
ABEM4	SC5314, <i>als1Δ::FRT/als1Δ::FRT, bgl2Δ::FRT/bgl2Δ::FRT, ecm33Δ::FRT/ecm33Δ::FRT</i>	79

4.2.2 Gene disruption and re-insertion

C. albicans gene sequences were obtained from the Candida Genome Database web site (<http://www.candidagenome.org/cgi-bin/seqTools>). The *als1* null mutant (ALS1M4); *als1, bgl2* null mutant (ABM4); *als1, ecm33* null mutant (AEM4); and *als1, bgl2, ecm33* triple mutant (ABEM4) had been constructed previously by our group using the *SAT1* flipper method.⁷

4.2.3 qRT-PCR

C. albicans cells were cultured in YPD medium at 30°C overnight. The culture was diluted to an OD₆₀₀ of 0.2 in 50 ml fresh YPD medium, and incubated at 30°C until the OD₆₀₀ reached 1–2. Total RNA was extracted using the glass beads lysis method⁶⁸. The RNA samples were used as templates to synthesize first strand cDNA using the ReverTra Ace qPCR RT Master Mix with gDNA Remover kit (Toyobo, Osaka, Japan). Real time PCR was performed using the THUNDERBIRD SYBR qPCR Mix (Toyobo) under the following conditions: 95°C for 10 min, followed by 40 cycles of 15 sec at 95°C, 30 seconds at 56°C, 1 min at 72°C. *ACT1* mRNA was used for normalization, and gene expression levels were determined using Applied Biosystems software and the $\Delta\Delta C_t$ method.

Figure 4-2: Primer for qRT PCR

Primer name	Sequence 5'-3'
<i>ACT1</i> forward	TGTGTAAAGCCGGTTTTGCC
<i>ACT1</i> reverse	TTGGATTGGGCTTCATCACC
<i>ALS1</i> forward	AGCGGTTCTCATGAATCAGC
<i>ALS1R</i> reverse	CAGAAGAAACAGCAGGTGATGG
<i>ALS2</i> forward	AGCGGTTCTCATGAATCAGC
<i>ALS2</i> reverse	CAGAAGAAACAGCAGGTGATGG
<i>ALS3</i> forward	TCGTTGACATGGACAAGAGC
<i>ALS3</i> reverse	ACCAGCCCAAACAGCATTTC
<i>ALS4</i> forward	TGCCGTTATCGTCCATTTG
<i>ALS4</i> reverse	ATGTGAAAGGTGCACGTTGC
<i>ALS5</i> forward	ACCCTGTGTTGACAACAAGC
<i>ALS5</i> reverse	ATAACCACTGTCGCAGTTGC
<i>ALS6</i> forward	TCCAATGCTGGCAATTACGC
<i>ALS6</i> reverse	AATGACCAACCCAACACAGC
<i>ALS7</i> forward	GCTGTTTTAGGTTGGAGCTTGG
<i>ALS7</i> reverse	TTGAAAACGCAGGGCATGAC
<i>ALS9</i> forward	ATCGACCTTTTGTGGATGCG

4.2.4 Assay of *C. albicans* adhesion to SHA beads

One colony of yeast cells of SC5314 (wild type), ALS1M4(*als1*Δ), ABM4 (*als1*Δ/*bgl2*Δ), AEM4 (*als1*Δ/*ecm33*Δ), and ABEM4 (*als1*Δ/*bgl2*Δ/*ecm33*Δ) were pre-cultured in 5mL GSB medium overnight at 30°C. Cells were harvested by centrifuging 5000 rpm for 5 minutes and then diluted in KCl buffer to OD₆₀₀ of 1. Adhesion assays were done in the same method described in chapter 3.

4.2.1 Fluorescence signals of different strains

One yeast colony of each 8 strains were inoculated in GSB medium overnight to OD₆₀₀=4-5. Cell pellet were harvested by centrifuging at 5000 rpm in 5 minutes and then were suspend in KCl buffer to OD_{600 nm} =1. Cells were then stained with NR in 10% DMSO with final concentration of 1 μg/ml. After 1 hour, two hundred μl of cell suspension were put in to 96 well plate and fluorescence signals were measured with excitation wavelength of 530 nm and 570 nm using Varioskan Lux system.

4.2.2 Growth curve of different strains in GSB medium

Overnight culture of 8 strains in YPD medium were harvested by centrifuging at 5000 rpm in 5 minutes and then were diluted to OD_{600 nm} =0.01 and inoculated in GSB medium with shaking 200 rpm. The OD_{600nm} values were measured after 2-4 hours and the growth curve were constructed.

4.2.3 The susceptibility of *C. albicans* strains to Calcofluor white (CFW)

One colony of yeast cells of SC5314 (wild type), ALS1M4(*als1*Δ), ABM4 (*als1*Δ/*bgl2*Δ), AEM4 (*als1*Δ/*ecm33*Δ), and ABEM4 (*als1*Δ/*bgl2*Δ/*ecm33*Δ) cells were cultured in 5 mL YPD medium overnight at 30°C and diluted in YPD to an OD₆₀₀ of 1. The cells were then serially diluted (10-fold) in YPD medium. Three microliters of each dilution were spotted onto YPD plates containing 12.5 μg/mL CFW, and incubated at

30°C for 24h. Susceptibility of cells to CFW was indicated by their inability to grow on this medium.

4.3 Results and discussion

4.3.1 *ALS1 gene was over-expressed in the bgl2Δ, ecm33Δ strain*

Since the BEM4 (*bgl2Δ, ecm33Δ*) strain showed higher adhesion to SHA beads than BGL2M4 (*bgl2Δ*) or ECM33M4 (*ecm33Δ*), I examined the expression of a family of important *C. albicans* adhesin genes in these strains. The *ALS* (Agglutinin-Like Sequence) genes encode a family of GPI-anchored cell surface glycoproteins, eight of which (Als1-Als7 and Als9) have been reported to be involved in *C. albicans* adhesion.^{27,75,80} Interestingly, qRT PCR indicated that *ALS1* mRNA was more than 6 fold higher in the *bgl2Δ, ecm33Δ* double mutant BEM4 (Figure 4-3) than that of *C. albicans* SC5314. In contrast, there was no significant increase in expression of any of the other *ALS* gene in BGL2M4, ECM33M4 and BEM4 (Figure 4-3). This indicates that the deletion of the *BGL2* and *ECM33* genes induces expression of *ALS1* which may be responsible for the increase in adherence of *C. albicans* to SHA beads.




Figure 4-3: The transcriptional expression of ALS family genes of *C. albicans* wild type and mutant strains. Four strains of *C. albicans* were cultured in YPD medium and then total RNA was isolated for qRT-PCR assay. *C. albicans* ACT1 gene was used as internal control in this experiment. These results are mean of at least three independent experiments.




Figure 4-4: The transcriptional expression of ALS family genes of *C. albicans* wild type and mutant strains. Four strains of *C. albicans* were cultured in YPD medium and then total RNA was isolated for qRT-PCR assay. *C. albicans* ACT1 gene was used as internal control in this experiment. These results are mean of at least three independent experiments.

4.3.2 The introduction of ALS1 gene disruptions in mutant strains and investigation of their effect on adherence to SHA beads




Figure 4-5: Adherence of ALS1, BGL2, and ECM33 single, double, and triple null mutants compared to the wild type strain *C. albicans* SC5314. The *C. albicans* cells (1×10^7) and SHA beads (12 mg) were mixed for 1 h at 28°C. After the incubation, the beads were washed twice with KCl buffer and all input cells and washed cells were stained with NR in KCl buffer containing 10% DMSO at the final concentration of 1 µg/ml. Results are means of at least three independent experiments. Statistical significance is shown (**p > 0.01 ***p > 0.001).

From these results, previously, *als1*Δ (ALS1M4), *als1*Δ and *bgl2*Δ (ABM4), *als1*Δ and *ecm33*Δ (AEM4), and *als1*Δ, *bgl2*Δ, and *ecm33*Δ (ABEM4) mutants were constructed from SC5314, BGL2M4, ECM33M4 and BEM4 strains.⁷⁹ After confirming the disruption of the *ALS1* gene in each strain, the adherence of these strains to SHA beads was measured. Deletion of *ALS1* in SC5314 (to form ALS1M4) reduced adherence ability and this percentage was as similar as (72.6±2.5 % of SC5314) as deletion of *BGL2* (76.4±13.0 %) or *ECM33* (64.8±9.1 %) (Figure 4-5). Deletion of *ALS1* in either a *bgl2*Δ or a *ecm33*Δ background did not reduce adhesion to SHA beads further (Figure 4-5). As predicted, however, deletion of *ALS1* in the *bgl2*Δ, *ecm33*Δ double disruptant (to form ABEM4) resulted in a significant reduction in adherence to SHA beads (37.6±4.3 % of

SC5414 adherence; Figure 4-5). These results indicated that reliance on Als1 for adherence to SHA beads is only triggered when both Bgl2 and Ecm33 are inactivated, and highlight the importance of all three proteins for the attachment of *C. albicans* cells to tooth surfaces via saliva in the oral cavity.

4.3.3 Fluorescence signals of different strains:




Figure 4-6: Fluorescence intensity of different strains at $OD_{600nm} = 1$

Although the adhesion ability of each strain was measured by the ratio of the fluorescence intensity of attached cells vs. total input cell, therefore, the errors were already minimized. However, there might be a difference of lipid concentrations between *C. albicans* WT SC5314 and all other mutants. Therefore, fluorescence intensity of all strains constructed in this study were compared. Yeast cells of all strains were inoculated in GSB medium to $OD_{600nm} = 4-5$, these cells were harvested and diluted cells to $OD_{600nm} = 1$. Fluorescence intensity of stained cells was measured at excitation 530 nm and emission 570 nm. This result in Figure 4-6 showed that most cells have similar fluorescence signals with each other which might answer the questions of the level of lipid contents in the cells. By deleting either one, two, or three different genes that play

roles in cell membrane, the concentration of lipid did not change drastically, therefore, the fluorescence signals of NR stained are similar between those strains.

4.3.4 Growth curve of different strains in GSB medium:




Figure 4-7: Growth curve of different strains in GSB medium

The aim of this experiment is to evaluate possible effect of deletion *BGL2*, *ECM33*, *ALS1*, double mutants and triple mutants of those genes to the cell growth. Overnight culture of these 8 strains were diluted to $OD_{600\text{ nm}} = 0.01$ and inoculated in GSB medium with shaking 200 rpm. The $OD_{600\text{ nm}}$ values were measured after 2-4 hours and the growth curve. As shown in Figure 4-7, all strains had similar growth curve, which indicated that there were no strong effect on the cell growth by deleting those three genes in *C. albicans* SC5314 cells.

4.3.5 Cells susceptibility




Figure 4-8: The susceptibility of *C. albicans* strains to Calcofluor white (CFW). SC5314 (wild type), ALS1M4(*als1*Δ), ABM4 (*als1*Δ/*bgl2*Δ), AEM4 (*als1*Δ/*ecm33*Δ), and ABEM4 (*als1*Δ/*bgl2*Δ/*ecm33*Δ) cells were cultured in 5 mL YPD medium overnight at 30°C and diluted in YPD to an OD₆₀₀ of 1. The cells were then serially diluted (10-fold) in YPD medium. Three microliters of each dilution were spotted onto YPD plates containing 12.5 µg/mL CFW, and incubated at 30°C for 24h. Susceptibility of cells to CFW was indicated by their inability to grow on this medium..

To investigate the impact of gene disruptions on *C. albicans* cell wall integrity, I tested the sensitivity of null mutants to calcofluor white (CFW), a chemical which interferes with cell wall construction.⁷¹ SC5314 (wild type), ALS1M4(*als1*Δ), ABM4 (*als1*Δ/*bgl2*Δ), AEM4 (*als1*Δ/*ecm33*Δ), and ABEM4 (*als1*Δ/*bgl2*Δ/*ecm33*Δ) cells were serially diluted and inoculated on YPD agar and on YPD agar containing CFW (12.5 µg/mL), and incubated for 24 h at 30°C. All of the null mutants were more susceptible to CFW than the wild type parental strain (Figure 4-8). ABEM4 (*als1*Δ/*bgl2*Δ/*ecm33*Δ) showed the highest CFW sensitivity. This result indicated that lack of cell surface mannoproteins leads to a significant decrease in the integrity of the *C. albicans* cell wall.

4.4 Conclusion

As an opportunistic microorganism, *C. albicans* adheres to ligands that retain the yeast cells in the host through different surfaces.¹⁴ These interactions prevent human fluids, such as saliva, from removing *C. albicans* from the host. Many methods were used to identify these interactions and the ability of the organism to bind to a variety of surfaces (cell, protein, or other substrate). Therefore, a number of *C. albicans* adhesins were detected including Als family of GPI-CWPs, Hwp1p, ECM33p, and extracellular hydrolytic enzymes.⁸¹ Bgl2p (β -1,3-glucosyltransferase) and Ecm33p (a typical feature of glycosylphosphatidylinositol (GPI) – anchored protein) are known to be cell wall proteins and related to the cell wall integrity^{6, 18}. In this study, it was confirmed that BGL2M4 (*bgl2Δ*) and ECM33M4 (*ecm33Δ*) showed higher sensitivity to CFW than SC5314 (wild type). Whereas the CFW susceptibility of BEM4 (*bgl2Δ, ecm33Δ*) was highest in all four strains. Then, it was suggested that the disruption of these double genes severely influenced the cell wall construction of *C. albicans*. Surprisingly, although the adhesion of BGL2M4 and ECM33M4 to SHA beads were 76.4% and 64.8% of SC5314, respectively, BEM4 strains showed 87.7 % of adhesion activity of the wild type and this activity was higher than BGL2M4 and ECM33M4. From these results, I assumed that a certain compensatory system to maintain the adherence activity was activated in BEM4 because lack of two cell wall proteins leads to drastic change in the cell wall integrity of the fungal cells. To investigate this assumption, *ALS* gene family, which encodes large cell surface glycoproteins and are critical in the process of adhesion to host surfaces, was focused. The *ALS* family includes eight genes, from *ALS1* to *ALS7* and *ALS9* and the transcriptional expression of these eight genes were analyzed. Interestingly, the expression level of *ALS1* in BEM4 is higher than those of wild-type and single gene disruptants. *ALS1* is reported to take part in *C. albicans* adhesion to human epithelial, endothelial cells and even abiotic surfaces such as silicon⁴². This protein is considered

key component needed for biofilm, hyphal development of *C. albicans*^{27,75} This result suggested that the double mutant had developed a compensatory response to overexpression of *ALSI* gene then recovered decrease in adhesion activity to SHA beads. In previous research, Chen *et al.* showed the *bgl2Δ* mutant (BGL2M4) took a delay in transition to filamentous cell morphology due to repression of transcription factor *CPH2/TECI*. *TECI* was identified as a transcriptional factor that is required for hypha formation and virulence.⁸² In addition, *TECI* is also reported to be an important transcription factor of *ECM33* and *ALSI*^{83,84}. Therefore, I hypothesized that in case of reduction of *BGL2* and *ECM33*, the expression of *ALSI* may be induced by *TECI*. To clarify this transcriptional mechanism, the further experiments on the protein network and functions of *C. albicans* outer cell wall constituents will be needed.

Collectively, in this chapter, I investigated the roles of Bgl2 and ECM33 in the ability of *C. albicans* to adhere to saliva-coated oral surface by generating several null mutants of these genes and a novel NR staining assay. I found that the two cell wall proteins Bgl2p, Ecm33p contribute to interaction between *C. albicans* and saliva-coated surfaces. However, I observed that the deletion of both two genes triggered compensatory responses in this yeast cells. In particular, the *ALSI* gene, encoding a major GPI-modified protein, was overexpressed in the *bgl2Δ/ecm33Δ* double disrupted strain. Then deletion of all of these three genes caused a significant decrease in the adhesion of *C. albicans* to saliva-coated hydroxyapatite (SHA) beads.

CHAPTER 5. CONCLUSION

Amongst *Candida* species, *C. albicans* is most commonly isolated from the oral cavity and is responsible for most superficial and systemic fungal infections¹⁵. The oral cavity provides many saliva-coated surfaces and niches that are easily colonized by *C. albicans*. Therefore, it is important to investigate how this opportunistic yeast interacts with tooth surfaces when covered by saliva components. The yeast cell wall is the vital point of contact between *C. albicans* cell and the saliva-coated oral surfaces.⁸ Because of this important role, the *C. albicans* cell wall has attracted a lot of attention over the last few decades.

A method used previously for measuring *C. albicans* adherence to saliva-coated surfaces involved labelling the yeast with [³⁵S]-methionine⁵. This method is reliable, reproducible and sensitive; however, it is time-consuming, expensive and poses a risk to human health. Recently, several alternative fluorescent staining strategies have been developed to provide fast detection and quantification of yeast cells. In my research, important criteria for choosing the optimum fluorescent dye to measure yeast adhesion were chemical stability under the experiment conditions, low-cost, safety of use, and no interference with other substances in saliva. I investigated using, carboxyfluorescein diacetate succinimidyl ester (CFDA) for staining of *C. albicans* cells as this compound is cleaved by intracellular esterase enzymes to form a fluorescent amine-reactive product. I found that CFDA stained *C. albicans* cells and gave a strong fluorescence signal but there were also many esterases in saliva resulting in a high background signal in our SHA bead adhesion assay. I then focused on NR, a commonly used fluorescent lipophilic dye that binds to intracellular neutral lipids. NR has low fluorescence in water and other polar solvents, but gives strong fluorescence signals in nonpolar environments.⁵⁵ I found, by using spectroscopy and confocal microscopy, that once NR was absorbed by *C. albicans* cells, the signal was stable, even after 2-5 h of staining. This characteristic is a major

advantage of using NR because once yeast cells were stained with this dye, no washing steps were needed. Therefore, I used NR to stain *C. albicans* cells for adhesion assays.

While uncoated HA beads showed low non-specific binding of 17% of SC5314 cells, I found that 89% yeast cells adhered to SM-coated HA beads, which is relatively equal to the amount of SHA beads. This is consistent with previous reports which have shown that bPRP and statherin promote attachment of *C. albicans* cells to hydroxyapatite beads^{5,14}. Control proteins FBS and BSA did not promote much adherence of *C. albicans* cells to hydroxyapatite beads but, surprisingly, skim milk did. It would be interesting to determine which components of skim milk promote this adhesion.

The proteins Bgl2 (β -1,3-glucosyltransferase) and Ecm33 (a GPI-anchored protein) are cell wall proteins contributing to cell wall integrity^{6,18}. In this study, it was confirmed that BGL2M4 (*bgl2* Δ) and ECM33M4 (*ecm33* Δ) showed higher sensitivity to CFW than SC5314 (wild type) and the double disruptant (*bgl2* Δ , *ecm33* Δ ; BEM4) showed even higher sensitivity. As well as having compromised cell wall integrity, BGL2M4 and ECM33M4 showed reduced adherence to SHA beads than SC5314. Surprisingly, the double disruptant BEM4 demonstrated increased adhesion to SHA beads. Deletion of ALS1 in either the *bgl2* Δ or the *ecm33* Δ background did not result in a strain with increased adherence to SHA beads (as was seen with the *bgl2* Δ , *ecm33* Δ strain). These results demonstrated that *ALS1* expression is only induced when both *BGL2* and *ECM33* are deleted, which was substantiated by the significant reduction in adherence of the *bgl2* Δ , *ecm33* Δ strain when *ALS1* was also disrupted. Als1p along with other Als proteins are well recognized adhesins, but their expression has been predominantly studied under hypha inducing conditions.^{80,85} Much is known about *ALS1* expression during morphogenesis and biofilm formation where adhesin gene expression is under the control of transcriptional regulators such as Efg1p, Bcr1p, Tup1p and Nrg1p.^{86,87} However, there have been very few studies on the expression of adhesins in yeast cells. Iare interested in

adhesin expression by yeast cells as they are responsible for the colonization of saliva-coated surfaces. *ALS1* is expressed by yeast cells when released from stationary phase, but expression decreases during yeast growth. In contrast, expression of *ALS1* is induced greatly during hyphal formation.⁸⁸ The present study shows that the expression of *ALS1* in yeast cells is increased in cells deleted in *BGL2* and *ECM33*. Chen *et al.* have previously reported that disruption of *BGL2* resulted in decreased expression of transcriptional regulator *CPH2* during yeast growth⁵¹ and this may be responsible, in part, for changes in *ALS1* expression. Further experiments are needed to elucidate the transcriptional network controlling the expression of Bgl2p, Ecm33p and Als1p under yeast and mycelia growth conditions.

It must be acknowledged that I only looked for compensatory expression of *ALS* genes in the disrupted strains. While there is strong evidence that there is induction of *ALS1* when *BGL2* and *ECM33* are deleted, I cannot exclude the possibility that expression of other adhesins is also induced. Further research, including complete RNA sequencing of mutant strains, may provide a more complete mechanistic insight into the role of the cell wall protein network in the adhesion ability of *C. albicans*. This study does, however, provide strong evidence that Bgl2p, Ecm33p and Als1p contribute to *C. albicans* adherence to saliva-coated surfaces.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Approval to conduct the study was obtained from the Research Ethics Committee of Tokyo Institute of Technology in accordance with the Ethical Guidelines for Clinical Research and Ethical Guidelines for Epidemiological Research (issued by the Ministry of Health, Labour and Welfare and Ministry of Education, Culture, Sports, Science and Technology - Japan).

REFERENCE

1. ten Cate, J. M., Klis, F. M., Pereira-Cenci, T., Crielaard, W. & de Groot, P. W. J. Molecular and cellular mechanisms that lead to Candida biofilm formation. *J. Dent. Res.* **88**, 105–15 (2009).
2. Pages, J. M. & Amaral, L. Mechanisms of drug efflux and strategies to combat them: challenging the efflux pump of Gram-negative bacteria. *Biochim Biophys Acta* **1794**, 826–833 (2009).
3. Patil, S., Rao, R. S., Majumdar, B. & Anil, S. Clinical appearance of oral Candida infection and therapeutic strategies. *Frontiers in Microbiology* (2015) doi:10.3389/fmicb.2015.01391.
4. Dühning, S. *et al.* Host-pathogen interactions between the human innate immune system and *Candida albicans*-understanding and modeling defense and evasion strategies. *Frontiers in Microbiology* (2015) doi:10.3389/fmicb.2015.00625.
5. Cannon, R. D., Nand, A. K. & Jenkinson, H. F. Adherence of *Candida albicans* to human salivary components adsorbed to hydroxylapatite. *Microbiology* (1995) doi:10.1099/00221287-141-1-213.
6. Mayer, F. L., Wilson, D. & Hube, B. *Candida albicans* pathogenicity mechanisms. *Virulence* (2013) doi:10.4161/viru.22913.
7. Chen, X. *et al.* The role of Bgl2p in the transition to filamentous cells during biofilm formation by *Candida albicans*. *Mycoses* **60**, 96–103 (2017).
8. Chaffin, W. L. *Candida albicans* Cell Wall Proteins. *Microbiol. Mol. Biol. Rev.* (2008) doi:10.1128/mubr.00032-07.
9. Naglik, J. R., Challacombe, S. J. & Hube, B. *Candida albicans* Secreted Aspartyl Proteinases in Virulence and Pathogenesis. *Microbiol. Mol. Biol. Rev.* (2003)

doi:10.1128/mmbr.67.3.400-428.2003.

10. Staab, J. F., Datta, K. & Rhee, P. Niche-specific requirement for hyphal wall protein 1 in virulence of *Candida albicans*. *PLoS One* (2013) doi:10.1371/journal.pone.0080842.
11. Citiulo, F. *et al.* *Candida albicans* scavenges host zinc via Pra1 during endothelial invasion. *PLoS Pathog.* (2012) doi:10.1371/journal.ppat.1002777.
12. Alves, R. *et al.* Adapting to survive: How *Candida* overcomes host-imposed constraints during human colonization. *PLoS pathogens* (2020) doi:10.1371/journal.ppat.1008478.
13. Singleton, D. R., Fidel, P. L., Wozniak, K. L. & Hazen, K. C. Contribution of cell surface hydrophobicity protein 1 (Csh1p) to virulence of hydrophobic *Candida albicans* serotype A cells. *FEMS Microbiol. Lett.* (2005) doi:10.1016/j.femsle.2005.02.010.
14. Cannon, R. D. *et al.* Adhesion of yeast and bacteria to oral surfaces. in *Methods in Molecular Biology* (2017). doi:10.1007/978-1-4939-6685-1_10.
15. Pereira, D., Seneviratne, C. J. & Koga-Ito, C. Y. Is the oral fungal pathogen *Candida albicans* a cariogen? *Oral Dis* (2017) doi:10.1111/odi.12691.
16. Garcia-Rubio, R., de Oliveira, H. C., Rivera, J. & Trevijano-Contador, N. The Fungal Cell Wall: *Candida*, *Cryptococcus*, and *Aspergillus* Species. *Frontiers in Microbiology* (2020) doi:10.3389/fmicb.2019.02993.
17. Gow, N. A. R. & Hube, B. Importance of the *Candida albicans* cell wall during commensalism and infection. *Current Opinion in Microbiology* (2012) doi:10.1016/j.mib.2012.04.005.
18. Klis, F. M., De Groot, P. & Hellingwerf, K. Molecular organization of the cell wall

- of *Candida albicans*. *Medical Mycology, Supplement* (2001)
doi:10.1080/mmy.39.1.1.8-0.
19. Sarthy, A. V *et al.* Phenotype in *Candida albicans* of a disruption of the BGL2 gene encoding a 1,3-beta-glucosyltransferase. *Microbiology* **143** (Pt 2, 367–376 (1997).
 20. Kalebina, T. S. *et al.* Deletion of BGL2 results in an increased chitin level in the cell wall of *Saccharomyces cerevisiae*. *Antonie Van Leeuwenhoek* **84**, 179–184 (2003).
 21. Chattaway, F. W., Holmes, M. R. & Barlow, A. J. Cell wall composition of the mycelial and blastospore forms of *Candida albicans*. *J. Gen. Microbiol.* (1968)
doi:10.1099/00221287-51-3-367.
 22. Free, S. J. Fungal Cell Wall Organization and Biosynthesis. in *Advances in Genetics* (2013). doi:10.1016/B978-0-12-407677-8.00002-6.
 23. Bulawa, C. E., Miller, D. W., Henry, L. K. & Becker, J. M. Attenuated virulence of chitin-deficient mutants of *Candida albicans*. *Proc. Natl. Acad. Sci. U. S. A.* (1995) doi:10.1073/pnas.92.23.10570.
 24. Lee, K. K. *et al.* Elevated cell wall chitin in *Candida albicans* confers echinocandin resistance in vivo. *Antimicrob. Agents Chemother.* (2012)
doi:10.1128/AAC.00683-11.
 25. Brown, G. D. *et al.* Dectin-1 is a major β -glucan receptor on macrophages. *J. Exp. Med.* (2002) doi:10.1084/jem.20020470.
 26. Brown, G. D. & Gordon, S. Immune recognition. A new receptor for beta-glucans. *Nature* (2001) doi:10.1038/35092620.
 27. Sundstrom, P. Adhesion in *Candida* spp. *Cellular Microbiology* (2002)
doi:10.1046/j.1462-5822.2002.00206.x.

28. Martinez-Lopez, R., Park, H., Myers, C. L., Gil, C. & Filler, S. G. Candida albicans Ecm33p Is Important for Normal Cell Wall Architecture and Interactions with Host Cells. *Eukaryot Cell* **5**, 140–147 (2006).
29. Martinez-Lopez, R., Monteoliva, L., Diez-Orejas, R., Nombela, C. & Gil, C. The GPI-anchored protein CaEcm33p is required for cell wall integrity, morphogenesis and virulence in Candida albicans. *Microbiology* (2004) doi:10.1099/mic.0.27320-0.
30. Hoyer, L. L., Payne, T. L., Bell, M., Myers, A. M. & Scherer, S. Candida albicans ALS3 and insights into the nature of the ALS gene family. *Curr Genet* **33**, 451–459 (1998).
31. Wächtler, B., Wilson, D., Haedicke, K., Dalle, F. & Hube, B. From attachment to damage: Defined genes of Candida albicans mediate adhesion, invasion and damage during interaction with oral epithelial cells. *PLoS One* (2011) doi:10.1371/journal.pone.0017046.
32. Valentijn-Benz, M., Nazmi, K., Brand, H. S., van't Hof, W. & Veerman, E. C. Growth of Candida albicans in human saliva is supported by low-molecular-mass compounds. *FEMS Yeast Res* **15**, (2015).
33. Porcheri, C. & Mitsiadis, T. Physiology, Pathology and Regeneration of Salivary Glands. *Cells* (2019) doi:10.3390/cells8090976.
34. Humphrey, S. P. & Williamson, R. T. A review of saliva: Normal composition, flow, and function. *J. Prosthet. Dent.* (2001) doi:10.1067/mpr.2001.113778.
35. Ghannam, M. G. & Singh, P. Anatomy, Head and Neck, Salivary Glands. in (2020).
36. Crosara, K. T. B. *et al.* Revealing the Amylase Interactome in Whole Saliva Using Proteomic Approaches. *Biomed Res. Int.* **2018**, 15 (2018).

37. Norris, H. L. *et al.* Salivary metals, age, and gender correlate with cultivable oral *Candida* carriage levels. *J Oral Microbiol* **10**, 1447216 (2018).
38. Mall, A. S., Habte, H., Mthembu, Y., Peacocke, J. & De Beer, C. Mucus and Mucins: Do they have a role in the inhibition of the human immunodeficiency virus? *Virology Journal* (2017) doi:10.1186/s12985-017-0855-9.
39. Gabryel-Porowska, H. *et al.* Mucin levels in saliva of adolescents with dental caries. *Med. Sci. Monit.* (2014) doi:10.12659/MSM.889718.
40. Wickström, C. & Svensäter, G. Salivary gel-forming mucin MUC5B - A nutrient for dental plaque bacteria. *Oral Microbiol. Immunol.* (2008) doi:10.1111/j.1399-302X.2007.00407.x.
41. Boehlke, C., Zierau, O. & Hannig, C. Salivary amylase - The enzyme of unspecialized euryphagous animals. *Archives of Oral Biology* (2015) doi:10.1016/j.archoralbio.2015.05.008.
42. Bennick, A. Salivary proline-rich proteins. *Mol. Cell. Biochem.* (1982) doi:10.1007/BF00223503.
43. Lamkin, M. S. & Oppenheim, F. G. Structural features of salivary function. in *Critical Reviews in Oral Biology and Medicine* (1993). doi:10.1177/10454411930040030101.
44. Hajishengallis, G. & Russell, M. W. Innate Humoral Defense Factors. in *Mucosal Immunology: Fourth Edition* (2015). doi:10.1016/B978-0-12-415847-4.00015-X.
45. Henskens, Y. M. C. *et al.* Protein composition of whole and parotid saliva in healthy and periodontitis subjects: Determination of cystatins, albumin, amylase and IgA. *J. Periodontal Res.* (1996) doi:10.1111/j.1600-0765.1996.tb00464.x.
46. Trochimiak, T. & Hübner-Woźniak, E. Effect of exercise on the level of

- immunoglobulin a in saliva. *Biol. Sport* (2012) doi:10.5604/20831862.1019662.
47. Hoyer, L. L. & Cota, E. Candida albicans agglutinin-like sequence (Als) family vignettes: A review of als protein structure and function. *Frontiers in Microbiology* (2016) doi:10.3389/fmicb.2016.00280.
 48. Desai, J. V. & Mitchell, A. P. Candida albicans Biofilm Development and Its Genetic Control. *Microbiol. Spectr.* (2015) doi:10.1128/microbiolspec.mb-0005-2014.
 49. Lin, C. H. *et al.* Genetic Control of Conventional and Pheromone-Stimulated Biofilm Formation in Candida albicans. *PLoS Pathog.* (2013) doi:10.1371/journal.ppat.1003305.
 50. Sarthy, A. V. *et al.* Phenotype in Candida albicans of a disruption of the BGL2 gene encoding a 1,3- β -glucosyltransferase. *Microbiology* (1997) doi:10.1099/00221287-143-2-367.
 51. Chen, X. *et al.* The role of Bgl2p in the transition to filamentous cells during biofilm formation by Candida albicans. *Mycoses* **60**, 96–103 (2017).
 52. Hayashi-Takanaka, Y., Stasevich, T. J., Kurumizaka, H., Nozaki, N. & Kimura, H. Evaluation of chemical fluorescent dyes as a protein conjugation partner for live cell imaging. *PLoS One* (2014) doi:10.1371/journal.pone.0106271.
 53. Hernlem, B. & Hua, S. S. Dual fluorochrome flow cytometric assessment of yeast viability. *Curr Microbiol* **61**, 57–63 (2010).
 54. Ligasová, A. & Koberna, K. Quantification of fixed adherent cells using a strong enhancer of the fluorescence of DNA dyes. *Sci. Rep.* (2019) doi:10.1038/s41598-019-45217-9.
 55. Rostron, K. A. & Lawrence, C. L. Nile Red Staining of Neutral Lipids in Yeast.

- Methods Mol Biol* **1560**, 219–229 (2017).
56. Chen, W., Zhang, C., Song, L., Sommerfeld, M. & Hu, Q. A high throughput Nile red method for quantitative measurement of neutral lipids in microalgae. *J Microbiol Methods* **77**, 41–47 (2009).
 57. Aleman-Nava, G. S. *et al.* How to use Nile Red, a selective fluorescent stain for microalgal neutral lipids. *J Microbiol Methods* **128**, 74–79 (2016).
 58. Genicot, G., Leroy, J. L. M. R., Van Soom, A. & Donnay, I. The use of a fluorescent dye, Nile red, to evaluate the lipid content of single mammalian oocytes. *Theriogenology* (2005) doi:10.1016/j.theriogenology.2004.06.006.
 59. Izard, J. & Limberger, R. J. Rapid screening method for quantitation of bacterial cell lipids from whole cells. *J. Microbiol. Methods* (2003) doi:10.1016/S0167-7012(03)00193-3.
 60. Rostron, K. A., Rolph, C. E. & Lawrence, C. L. Nile red fluorescence screening facilitating neutral lipid phenotype determination in budding yeast, *Saccharomyces cerevisiae*, and the fission yeast *Schizosaccharomyces pombe*. *Antonie Van Leeuwenhoek* **108**, 97–106 (2015).
 61. Ivnitski-Steele, I. *et al.* Identification of Nile Red as a fluorescent substrate of the *Candida albicans* ABC transporters Cdr1p and Cdr2p and the MFS transporter Mdr1p. *Anal. Biochem.* **394**, 87–91 (2009).
 62. Seneviratne, C. J., Jin, L. & Samaranayake, L. P. Biofilm lifestyle of *Candida*: a mini review. *Oral Dis* **14**, 582–590 (2008).
 63. Xu, Q. R. *et al.* Molecular genetic techniques for gene manipulation in *Candida albicans*. *Virulence* (2014) doi:10.4161/viru.28893.
 64. Wilson, D. *et al.* Distinct roles of *Candida albicans*-specific genes in host-pathogen

- interactions. *Eukaryot. Cell* (2014) doi:10.1128/EC.00051-14.
65. Reuß, O., Vik, Å., Kolter, R. & Morschhäuser, J. The SAT1 flipper, an optimized tool for gene disruption in *Candida albicans*. *Gene* (2004) doi:10.1016/j.gene.2004.06.021.
 66. Noble, S. M. & Johnson, A. D. Genetics of *Candida albicans*, a Diploid Human Fungal Pathogen. *Annu. Rev. Genet.* (2007) doi:10.1146/annurev.genet.41.042007.170146.
 67. Holmes, A. R., Cannon, R. D. & Shepherd, M. G. Effect of calcium ion uptake on *Candida albicans* morphology. *FEMS Microbiol. Lett.* (1991) doi:10.1111/j.1574-6968.1991.tb04345.x.
 68. Burke, D. J., Dawson, D. & Stearns, T. Methods in Yeast Genetics: A Cold Spring Harbor Laboratory Course Manual. *A Cold Spring Harb. Lab. Course Man.* (2000).
 69. Gillum, A. M., Tsay, E. Y. H. & Kirsch, D. R. Isolation of the *Candida albicans* gene for orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae* *ura3* and *E. coli* *pyrF* mutations. *MGG Mol. Gen. Genet.* (1984) doi:10.1007/BF00328721.
 70. RUOYU, Z. The Function Analysis of Bgl2p and Ecm33p in *Candida albicans* Adhesion to Human Oral Cavity. (2012).
 71. Ram, A. F. J. & Klis, F. M. Identification of fungal cell wall mutants using susceptibility assays based on Calcofluor white and Congo red. *Nat. Protoc.* (2006) doi:10.1038/nprot.2006.397.
 72. Valentijn-Benz, M., Nazmi, K., Brand, H. S., Hof, W. V. & Veerman, E.C.I. Growth of *Candida albicans* in human saliva is supported by low-molecular-mass compounds. *FEMS Yeast Res.* (2015) doi:10.1093/femsyr/fov088.

73. Jeng, H. W., Holmes, A. R. & Cannon, R. D. Characterization of two *Candida albicans* surface mannoprotein adhesins that bind immobilized saliva components. *Med. Mycol.* (2005) doi:10.1080/13693780410001731637.
74. O'Sullivan, J. M., Cannon, R. D., Sullivan, P. A. & Jenkinson, H. F. Identification of salivary basic proline-rich proteins as receptors for *Candida albicans* adhesion. *Microbiology* **143** (Pt 2, 341–348 (1997).
75. Ho, V. *et al.* An amyloid core sequence in the major *Candida albicans* adhesin Als1p mediates cell-cell adhesion. *MBio* (2019) doi:10.1128/mBio.01766-19.
76. Jung, S. I. *et al.* Yeast casein kinase 2 governs morphology, biofilm formation, cell wall integrity, and host cell damage of *Candida albicans*. **12**, e0187721 (2017).
77. Hoyer, L. L. The ALS gene family of *Candida albicans*. *Trends in Microbiology* (2001) doi:10.1016/S0966-842X(01)01984-9.
78. Hoyer, L. L., Green, C. B., Oh, S. H. & Zhao, X. Discovering the secrets of the *Candida albicans* agglutinin-like sequence (ALS) gene family - A sticky pursuit. *Med. Mycol.* (2008) doi:10.1080/13693780701435317.
79. Inokawa. *Candida albicans*の細胞接着に関わる 細胞表層タンパク質生産とその役割. (2014).
80. Fu, Y. *et al.* *Candida albicans* Als1p: An adhesin that is a downstream effector of the EFG1 filamentation pathway. *Mol. Microbiol.* (2002) doi:10.1046/j.1365-2958.2002.02873.x.
81. Biswas, S., Van Dijck, P. & Datta, A. Environmental Sensing and Signal Transduction Pathways Regulating Morphopathogenic Determinants of *Candida albicans*. *Microbiol. Mol. Biol. Rev.* (2007) doi:10.1128/mubr.00009-06.
82. Liu, H. Transcriptional control of dimorphism in *Candida albicans*. *Current*

- Opinion in Microbiology* (2001) doi:10.1016/S1369-5274(01)00275-2.
83. Modrzewska, B. & Kurnatowski, P. Adherence of *Candida* sp. to host tissues and cells as one of its pathogenicity features. *Annals of parasitology* (2015).
 84. Roudbarmohammadi, S. *et al.* ALS1 and ALS3 gene expression and biofilm formation in *Candida albicans* isolated from vulvovaginal candidiasis . *Adv. Biomed. Res.* (2016) doi:10.4103/2277-9175.183666.
 85. Zhao, X. *et al.* ALS3 and ALS8 represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p. *Microbiology* (2004) doi:10.1099/mic.0.26943-0.
 86. Argimón, S. *et al.* Developmental regulation of an adhesin gene during cellular morphogenesis in the fungal pathogen *Candida albicans*. *Eukaryot. Cell* (2007) doi:10.1128/EC.00340-06.
 87. Chen, H. F. & Lan, C. Y. Role of SFP1 in the regulation of *Candida albicans* biofilm formation. *PLoS One* (2015) doi:10.1371/journal.pone.0129903.
 88. Green, C. B., Zhao, X., Yeater, K. M. & Hoyer, L. L. Construction and real-time RT-PCR validation of *Candida albicans* PALS-GFP reporter strains and their use in flow cytometry analysis of ALS gene expression in budding and filamenting cells. *Microbiology* (2005) doi:10.1099/mic.0.27696-0.

APPENDIX

A1. Adhesion procedure

