T2R2 東京科学大学 リサーチリポジトリ Science Tokyo Research Repository

論文 / 著書情報 Article / Book Information

Title	A definition of the asymptotic phase for quantum nonlinear oscillators from the Koopman operator viewpoint
Authors	Yuzuru Kato, Hiroya Nakao
Citation	Chaos: An Interdisciplinary Journal of Nonlinear Science, Volume 32, Issue 6,
Pub. date	2022, 6
URL	http://dx.doi.org/10.1063/5.0088559
note	This article may be downloaded for personal use only. Any other use

ARTICLE

View Onlin

A definition of the asymptotic phase for quantum nonlinear oscillators from the Koopman operator viewpoint **o**

Cite as: Chaos **32**, 063133 (2022); doi: 10.1063/5.0088559 Submitted: 18 February 2022 · Accepted: 24 May 2022 · Published Online: 24 June 2022

Yuzuru Kato^{1,a)} 🕩 and Hiroya Nakao^{2,b)} 🕩

AFFILIATIONS

¹Department of Complex and Intelligent Systems, Future University Hakodate, Hokkaido 041-8655, Japan ²Department of Systems and Control Engineering, Tokyo Institute of Technology, Tokyo 152-8552, Japan

a)Author to whom correspondence should be addressed: katoyuzu@fun.ac.jp

^{b)}E-mail: nakao@sc.e.titech.ac.jp

ABSTRACT

We propose a definition of the asymptotic phase for quantum nonlinear oscillators from the viewpoint of the Koopman operator theory. The asymptotic phase is a fundamental quantity for the analysis of classical limit-cycle oscillators, but it has not been defined explicitly for quantum nonlinear oscillators. In this study, we define the asymptotic phase for quantum oscillatory systems by using the eigenoperator of the backward Liouville operator associated with the fundamental oscillation frequency. By using the quantum van der Pol oscillator with a Kerr effect as an example, we illustrate that the proposed asymptotic phase appropriately yields isochronous phase values in both semiclassical and strong quantum regimes.

Published under an exclusive license by AIP Publishing. https://doi.org/10.1063/5.0088559

Spontaneous rhythmic oscillations and synchronization are observed in a wide variety of classical rhythmic systems, and recent progress in nanotechnology is facilitating the analysis of quantum rhythmic systems. The asymptotic phase plays a fundamental role in the analysis of classical limit-cycle oscillators, but a fully quantum-mechanical definition for quantum limit-cycle oscillators has been lacking. In this study, we propose a definition of the asymptotic phase for quantum nonlinear oscillators, which naturally extends the definition of the asymptotic phase for classical stochastic oscillatory systems¹ from the Koopman-operator viewpoint² and provides us with appropriate phase values for characterizing quantum synchronization.

I. INTRODUCTION

Synchronization of spontaneous rhythmic oscillations is widely observed in nature.^{3–8} It has been extensively studied in a variety of classical systems in physics, chemistry, and biology. Recently, much progress has been made in the experimental realization of synchronization in micro- and nanoscale systems, such as nanoelectromechanical oscillators,⁹ microlasers,¹⁰ spin torque oscillators,¹¹ and optomechanical oscillators.¹² Stimulated by the experimental developments, quantum synchronization has attracted much attention recently,^{13–41} and theoretical investigations of quantum signatures in synchronization, such as quantum fluctuations,^{13–16,31} quantum entanglement,^{18–20} discrete nature of the energy spectrum,^{21–23} and effects of quantum measurement,^{24–28} have been carried out. Experimental realizations of quantum phase synchronization in spin-1 atoms²⁹ and on the IBM Q system³⁰ have also been reported recently.

In classical deterministic systems, spontaneous rhythmic oscillations are typically modeled as stable limit cycles of nonlinear dynamical systems. The *asymptotic phase*,^{3–7} defined by the oscillator's vector field and increases with a constant frequency in the basin of the limit cycle, plays a central role in analyzing synchronization properties of limit-cycle oscillators. It is the basis for *phase reduction*,^{3–7} which gives low-dimensional phase equations approximately describing the oscillators under weak perturbations. Recently, it has been clarified that the asymptotic phase, which was originally introduced from a geometrical viewpoint,³ has a natural relationship with the Koopman eigenfunction associated with the fundamental frequency of the oscillator.^{2,42–45}

For classical stochastic oscillatory systems, Thomas and Lindner¹ proposed a definition of the asymptotic phase in terms of

the slowest decaying eigenfunction of the backward Fokker–Planck (Kolmogorov) operator describing the mean first passage time, which appropriately yields isochronous phase values that increase with a constant frequency on average even for strongly stochastic oscillations, in a similar way to the ordinary asymptotic phase for deterministic oscillators. We recently pointed out that their definition can be considered a natural extension of the deterministic definition from the Koopman operator viewpoint² (see Refs. 46–48 for the details of the stochastic Koopman operator).

The classical definitions of the asymptotic phase are applicable to quantum nonlinear oscillators in the semiclassical regime, where the system is described by a stochastic differential equation for the phase-space state along a deterministic classical trajectory under the effect of small quantum noise.^{2,16,31} However, in the stronger quantum regime, we cannot rely on the semiclassical approximation and how to define the asymptotic phase in a fully quantum-mechanical manner is an open question. In this study, we propose a definition of the asymptotic phase for quantum nonlinear oscillatory systems by using the eigenoperator of the adjoint Liouville operator, which is a counterpart of the backward Fokker–Planck operator in classical stochastic systems. We illustrate the validity of our definition by using a quantum van der Pol oscillator with the quantum Kerr effect in both semiclassical and strong quantum regimes.

II. ASYMPTOTIC PHASE FOR CLASSICAL OSCILLATORY SYSTEMS

In this section, we briefly review the definitions of the asymptotic phase for deterministic^{3–7,43–45} and stochastic^{1,2} classical oscillatory systems.

A. Deterministic oscillatory systems

Consider a deterministic dynamical system

$$\dot{X}(t) = A(X(t)), \tag{1}$$

where $X(t) \in \mathbb{R}^N$ is the system state at time t, $A(X) \in \mathbb{R}^N$ is a vector field representing the system dynamics, and () represents time derivative. We assume that this system has an exponentially stable limit-cycle solution $X_0(t)$ with a natural period T and frequency $\Omega_c = 2\pi/T$, satisfying $X_0(t+T) = X_0(t)$, and denote its basin of attraction as $B \subseteq \mathbb{R}^N$. The asymptotic phase $\Phi_c(X) : B \to [0, 2\pi)$ is defined such that $A(X) \cdot \nabla \Phi_c(X) = \Omega_c$ is satisfied for $\forall X \in B$, where $\nabla = \partial/\partial X$ represents the gradient with respect to X^{3-7} . The asymptotic phase $\phi(t) = \Phi_c(X(t))$ of the system state X(t) then obeys

$$\dot{\phi}(t) = \dot{\Phi}_c(\boldsymbol{X}(t)) = \boldsymbol{A}(\boldsymbol{X}(t)) \cdot \nabla \Phi_c(\boldsymbol{X}(t)) = \Omega_c, \quad (2)$$

i.e., ϕ always increases with a constant frequency Ω_c as X evolves in B. Thus, the asymptotic phase Φ_c gives a nonlinear transformation of the system state X to a phase value ϕ such that the dynamics of ϕ takes a simple linear form, $\phi(t) = \Omega_c t + \text{constant}$. The simplicity of the phase equation (2) has facilitated detailed studies of synchronization and collective dynamics in coupled-oscillator systems.³⁻⁷ The level sets of $\Phi_c(X)$ are called isochrons.

The linear operator $A = A(X) \cdot \nabla$ in the definition of the asymptotic phase $\Phi_c(X)$ is the infinitesimal generator of the *Koopman operator* describing the evolution of observables for the system

described in Eq. (1) (see Refs. 42–45 for details). The complex exponential $\Psi_c(X) = e^{i\Phi_c(X)}$ of $\Phi_c(X)$ is an eigenfunction of A with the eigenvalue $i\Omega_c$, namely, it satisfies the eigenvalue equation $A\Psi_c(X) = i\Omega_c\Psi_c(X)$. Therefore, the asymptotic phase $\Phi_c(X)$ has a natural operator-theoretic interpretation as the argument of the Koopman eigenfunction $\Psi_c(X)$ associated with the eigenvalue $i\Omega_c$, characterized by the fundamental frequency Ω_c of the oscillator.^{42–45} We can, thus, define the asymptotic phase by using the Koopman eigenfunction $\Psi_c(X)$ of A as

$$\Phi_c(\mathbf{X}) = \arg \Psi_c(\mathbf{X}). \tag{3}$$

B. Stochastic oscillatory systems

For stochastic oscillatory systems, we cannot use the deterministic limit-cycle solution in defining the asymptotic phase unless the noise is sufficiently weak. Thomas and Lindner¹ defined the asymptotic phase for classical stochastic oscillators without relying on the deterministic limit cycle by using the eigenfunction with the slowest decay rate of the backward Fokker–Planck operator.

Consider a stochastic oscillator described by an Ito stochastic differential equation (SDE)

$$d\mathbf{X}(t) = \mathbf{A}(\mathbf{X}(t)) \, dt + \mathbf{B}(\mathbf{X}(t)) \, d\mathbf{W}(t), \tag{4}$$

where $X(t) \in \mathbb{R}^N$ is the system state at time t, $A(X) \in \mathbb{R}^N$ is a drift vector representing the deterministic dynamics, $B(X) \in \mathbb{R}^{N \times N}$ is a matrix characterizing the effect of the noise, and $W(t) \in \mathbb{R}^N$ is a *N*-dimensional Wiener process. This system is assumed to be *oscillatory* in the sense explained below. The transition probability density p(X, t|Y, s) ($t \ge s$) of Eq. (4) obeys the forward and backward Fokker–Planck equations,⁴⁹

$$\frac{\partial}{\partial t}p(\mathbf{X},t|\mathbf{Y},s) = L_{\mathbf{X}}p(\mathbf{X},t|\mathbf{Y},s)$$
(5)

and

$$\frac{\partial}{\partial s}p(\mathbf{X},t|\mathbf{Y},s) = -L_{\mathbf{Y}}^{*}p(\mathbf{X},t|\mathbf{Y},s), \tag{6}$$

respectively, where the (forward) Fokker–Planck operator is given by

$$L_{\mathbf{X}} = -\frac{\partial}{\partial \mathbf{X}} \mathbf{A}(\mathbf{X}) + \frac{1}{2} \frac{\partial^2}{\partial \mathbf{X}^2} \mathbf{D}(\mathbf{X})$$
(7)

and the backward Fokker–Planck operator is given by (in terms of X)

$$L_X^* = A(X)\frac{\partial}{\partial X} + \frac{1}{2}D(X)\frac{\partial^2}{\partial X^2}.$$
 (8)

Here, $D(X) = B(X)B(X)^{\mathsf{T}} \in \mathbb{R}^{N \times N}$ is a matrix of diffusion coefficients (T indicates the matrix transposition). The forward and backward operators L_X and L_X^* are mutually adjoint, i.e., $\langle L_X G(X), H(X) \rangle_X = \langle G(X), L_X^* H(X) \rangle_X$, where the inner product is defined as $\langle G(X), H(X) \rangle_X = \int \overline{G(X)} H(X) \, dX$ for two functions $G(X), H(X) : \mathbb{R}^N \to \mathbb{C}$ (the overline indicates complex conjugate and the integration is taken over the whole range of X).

As explained in Appendix A, the backward Fokker–Planck operator L_X^* in Eq. (8) is the infinitesimal generator of the Koopman operator for Eq. (4). We denote the probability density function

of *X* at time *t* as $p_t(X) \in \mathbb{R}$, which also obeys the Fokker–Planck equation $\partial p_t(X)/\partial t = L_X p_t(X)$, and an observable of the system at time *t* as $g_t : \mathbb{R}^N \to \mathbb{C}$, which maps the system state *X* to a complex value. The evolution of the expectation $\langle g \rangle_t = \int p_t(X)g_t(X) dX = \langle p_t(X), g_t(X) \rangle_X$ of the observable *g* at $t = t_0$ can be expressed as

$$\frac{d}{dt} \langle g \rangle_t \Big|_{t=t_0} = \left\langle \frac{\partial}{\partial t} p_t(\mathbf{X}) \Big|_{t=t_0}, g_{t_0}(\mathbf{X}) \right\rangle_X$$

$$= \langle L_X p_{t_0}(\mathbf{X}), g_{t_0}(\mathbf{X}) \rangle_X = \langle p_{t_0}(\mathbf{X}), L_X^* g_{t_0}(\mathbf{X}) \rangle_X$$

$$= \left\langle p_{t_0}(\mathbf{X}), \frac{\partial}{\partial t} g_t(\mathbf{X}) \Big|_{t=t_0} \right\rangle_X, \tag{9}$$

where $g_t(X)$ remains constant and $p_t(X)$ evolves in the second expression, while $p_t(X)$ remains constant and $g_t(X)$ evolves in the last expression.

The linear differential operators L_X and L_X^* have a biorthogonal eigensystem $\{\lambda_k, P_k, Q_k\}_{k=0,1,2,...}$ of the eigenvalue λ_k and eigenfunctions $P_k(X)$ and $Q_k(X)$ satisfying

$$L_{\mathbf{X}}P_{k}(\mathbf{X}) = \lambda_{k}P_{k}(\mathbf{X}), \quad L_{\mathbf{X}}^{*}Q_{k}(\mathbf{X}) = \lambda_{k}Q_{k}(\mathbf{X}),$$

$$\langle P_{k}(\mathbf{X}), Q_{l}(\mathbf{X})\rangle_{\mathbf{X}} = \delta_{kl},$$

(10)

where k, l = 0, 1, 2, ... and δ_{kl} represents the Kronecker delta.⁴⁹ We assume that, among the eigenvalues, one eigenvalue λ_0 is zero, which is associated with the stationary state $p^S(\mathbf{X})$ of the system satisfying $L_{\mathbf{X}}p^S(\mathbf{X}) = 0$, and all other eigenvalues have negative real parts. It is assumed that the eigenvalues with the largest non-negative real part (or the smallest absolute real part, i.e., the slowest decay rate) are given by a complex–conjugate pair. We denote these eigenvalues as

$$\lambda_1 = \mu_s - i\Omega_s, \quad \overline{\lambda_1} = \mu_s + i\Omega_s, \tag{11}$$

where $|\mu_s|(\mu_s < 0)$ is the decay rate and $\Omega_s = \text{Im }\overline{\lambda_1}$ represents the fundamental oscillation frequency of the system. The oscillatory property of the system is embodied in this assumption. We also assume that this pair of principal eigenvalues are well separated from other branches of eigenvalues and this oscillatory mode is dominant in the system.

Thomas and Lindner¹ proposed a definition of the *stochastic* asymptotic phase of the system described by Eq. (4) by using the argument of the complex conjugate of the eigenfunction $Q_1(\mathbf{X})$ of $L_{\mathbf{X}}^*$ associated with $\overline{\lambda_1}$ as (in the notation used here)

$$\Phi_s(\mathbf{X}) = \arg Q_1(\mathbf{X}). \tag{12}$$

This definition is natural from the Koopman operator viewpoint,² as L_X^* is the infinitesimal generator of the Koopman operator for Eq. (4) that formally goes back to the linear operator $A = A(X) \cdot \nabla$, i.e., to the infinitesimal generator of the Koopman operator for the deterministic system Eq. (1) in the noiseless limit $D(X) \rightarrow 0$. The exponential average of Φ_s satisfies²

$$\frac{d}{dt}\arg \mathbb{E}^{X_0}[e^{i\Phi_s(X(t))}] = \frac{d}{dt}\arg \mathbb{E}^{X_0}[Q_1(X(t))] = \operatorname{Im}\overline{\lambda_1} = \Omega_s, \quad (13)$$

where \mathbb{E}^{X_0} represents the average over the stochastic trajectories of Eq. (4) starting from an initial point $X_0 \in \mathbb{R}^N$. Thus, the asymptotic phase defined by Eq. (12) increases with a constant frequency Ω_s

on average with the stochastic evolution of the system and can be considered a natural generalization of the deterministic asymptotic phase in Eq. (3). It is noted that we may also choose the eigenvalue λ_1 and eigenfunction $\overline{Q_1(X)}$ to define the stochastic asymptotic phase, which reverses its direction of increase.

III. ASYMPTOTIC PHASE FOR QUANTUM OSCILLATORY SYSTEMS

Our aim in this study is to propose a quantum-mechanical definition of the asymptotic phase that does not rely on classical trajectories. In Ref. 16, we developed a semiclassical phase reduction theory for quantum limit-cycle oscillators, but the definition of the asymptotic phase was based on the deterministic limit cycle in the classical limit and could not be applied in stronger quantum regimes. In Ref. 2, we considered the asymptotic phase of quantum limit-cycle oscillators using the definition for the classical stochastic oscillators explained in Sec. II, but it was valid only in the semiclassical regime. Here, we consider a quantum master equation describing the evolution of the density operator of the system and define the asymptotic phase by using the eigenoperator of the adjoint Liouville operator. We use standard notations for open quantum systems without a detailed explanation; see, e.g., Refs. 51–53 for details.

A. Quantum master equation

We consider quantum oscillatory systems with a single degree of freedom coupled to reservoirs. The system's quantum state is represented by a Hermitian density operator ρ ($=\rho^{\dagger}$) and the observable is described by an operator *F*, where \dagger represents Hermitian conjugate. Introducing an inner product $\langle X, Y \rangle_{tr} = \text{Tr}(X^{\dagger}Y)$ of two operators *X* and *Y*, the expectation of the observable *F* is expressed as

$$\langle F \rangle = \operatorname{Tr} (\rho F) = \langle \rho, F \rangle_{tr}.$$
 (14)

In the Schrödinger picture, the quantum state ρ evolves with time while the observable *F* remains constant. Assuming that the interactions of the system with the reservoirs are instantaneous and Markovian approximation can be employed, the evolution of the density operator ρ_t at time *t* obeys the quantum master equation⁵¹⁻⁵³

$$\dot{\rho}_t = \mathcal{L}\rho_t,\tag{15}$$

where the Liouville operator \mathcal{L} (sometimes called a superoperator because it acts on an operator) is given by

$$\mathcal{L}X = -i[H, X] + \sum_{j=1}^{n} \mathcal{D}[C_j]X.$$
(16)

Here, $H(=H^{\dagger})$ is a system Hamiltonian, C_j is a coupling operator between the system and *j*th reservoir (j = 1, ..., n), [A, B] = AB - BA is the commutator, $\mathcal{D}[C]X = CXC^{\dagger} - (XC^{\dagger}C + C^{\dagger}CX)/2$ is the Lindblad form, and the reduced Planck's constant is set as $\hbar = 1$.

In the Heisenberg picture, the quantum state ρ remains constant while the observable *F* evolves with time as

$$\dot{F}_t = \mathcal{L}^* F_t, \tag{17}$$

where the time dependence of *F* is explicitly shown. Here, \mathcal{L}^* is the adjoint operator of \mathcal{L} satisfying

$$\langle \mathcal{L}X, Y \rangle_{tr} = \langle X, \mathcal{L}^*Y \rangle_{tr}$$
 (18)

and can be explicitly calculated as

$$\mathcal{L}^* X = i[H, X] + \sum_{j=1}^n \mathcal{D}^+[C_j]X,$$
 (19)

where $\mathcal{D}^+[C]X = C^{\dagger}XC - (XC^{\dagger}C + C^{\dagger}CX)/2$ is the adjoint Lindblad form.⁵³ The evolution of the expectation $\langle F \rangle_t = \langle \rho_t, F_t \rangle_{tr}$ of *F* with respect to ρ at $t = t_0$ can be expressed as

$$\frac{d}{dt} \langle F \rangle_t \Big|_{t=t_0} = \langle \dot{\rho}_t |_{t=t_0}, F_{t_0} \rangle_{tr} = \langle \mathcal{L} \rho_{t_0}, F_{t_0} \rangle_{tr}$$
$$= \langle \rho_{t_0}, \mathcal{L}^* F_{t_0} \rangle_{tr} = \langle \rho_{t_0}, \dot{F}_t |_{t=t_0} \rangle_{tr}, \quad (20)$$

where *F* remains constant and ρ evolves in the second expression (Schrödinger picture), while ρ remains constant and *F* evolves in the last expression (Heisenberg picture). Equation (20) corresponds to Eq. (9) for the expectation of the observable for classical stochastic systems. Thus, the adjoint operator \mathcal{L}^* is a counterpart of the backward Fokker–Planck operator L_X^* in Sec. II, namely, \mathcal{L}^* corresponds to the infinitesimal generator of the Koopman operator.

We assume that the operators \mathcal{L} and \mathcal{L}^* have a biorthogonal eigensystem $\{\Lambda_k, U_k, V_k\}_{k=0,1,2,...}$ consisting of the eigenvalue Λ_k and right and left eigenoperators U_k and V_k , satisfying

$$\mathcal{L}U_k = \Lambda_k U_k, \quad \mathcal{L}^* V_k = \overline{\Lambda_k} V_k, \quad \langle U_k, V_l \rangle_{tr} = \delta_{kl},$$
(21)

for $k, l = 0, 1, 2, ...^{54}$ Among the eigenvalues, one eigenvalue Λ_0 is always 0, which is associated with the stationary state ρ^S of the system satisfying $\mathcal{L}\rho^S = 0$, and all other eigenvalues have negative real parts; this also indicates that the system has no decoherece free subspace.⁵⁰ We assume that, reflecting the system's oscillatory dynamics, the eigenvalues with the largest non-vanishing real part (i.e., with the slowest decay rate) are given by a complex-conjugate pair. We denote these eigenvalues as

$$\Lambda_1 = \mu_q - i\Omega_q, \quad \overline{\Lambda_1} = \mu_q + i\Omega_q, \tag{22}$$

where $|\mu_q|$ ($\mu_q < 0$) is the decay rate and $\Omega_q = \text{Im }\overline{\Lambda_1}$ gives the fundamental frequency of the oscillation. As in the case of the stochastic oscillatory systems in Sec. II, the oscillatory property of the system is embodied in this assumption.

B. Phase space representation

The density operator ρ can also be represented by using quasiprobability distributions in the phase space such as the *P*, *Q*, and Wigner distributions.^{51,52,55} We use the *P* representation and express ρ as

$$\rho = \int p(\boldsymbol{\alpha}) |\alpha\rangle \langle \alpha| \, d\boldsymbol{\alpha}, \tag{23}$$

where $|\alpha\rangle$ is a coherent state specified by a complex value $\alpha \in \mathbb{C}$, or equivalently by a complex vector $\boldsymbol{\alpha} = (\alpha, \overline{\alpha})^T \in \mathbb{C}^2, p(\boldsymbol{\alpha}) : \mathbb{C}^2 \to \mathbb{R}$

is a quasiprobability distribution of α , $d\alpha = d\alpha \, d\overline{\alpha}$, and the integral is taken over \mathbb{C} . Defining the *P* representation of an observable *F* as

$$f(\boldsymbol{\alpha}) = \langle \boldsymbol{\alpha} | F | \boldsymbol{\alpha} \rangle, \tag{24}$$

where *F* is expressed in the normal order, 51, 52, 55 the expectation of *F* is expressed as

$$\langle F \rangle = \operatorname{Tr}(\rho F) = \int p(\boldsymbol{\alpha}) f(\boldsymbol{\alpha}) \, d\boldsymbol{\alpha} = \langle p(\boldsymbol{\alpha}), f(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}}.$$
 (25)

Here, we defined the L^2 inner product $\langle g(\boldsymbol{\alpha}), h(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}} = \int g(\boldsymbol{\alpha}) h(\boldsymbol{\alpha}) d\boldsymbol{\alpha}$ of two functions $g(\boldsymbol{\alpha}), h(\boldsymbol{\alpha}) : \mathbb{C}^2 \to \mathbb{C}$, where the integral is taken over the complex plane.

In the Schrödinger picture, the time evolution of $p_t(\boldsymbol{\alpha})$ (dependence on *t* is explicitly denoted) corresponding to the master equation (15) is described by a partial differential equation

$$\frac{\partial}{\partial t}p_t(\boldsymbol{\alpha}) = L_{\boldsymbol{\alpha}}p_t(\boldsymbol{\alpha}), \qquad (26)$$

where the differential operator L_{α} is related to the Liouville operator \mathcal{L} in Eq. (15) via

$$\mathcal{L}\rho_t = \int L_{\boldsymbol{\alpha}} p_t(\boldsymbol{\alpha}) |\alpha\rangle \langle \alpha | \, d\boldsymbol{\alpha}$$

and can be explicitly calculated from Eq. (15) by using the standard calculus for the phase-space representation.^{51,52,55}

The corresponding evolution of the *P* representation $f_t(\alpha)$ of the observable F_t in the Heisenberg picture is given by

$$\frac{\partial}{\partial t} f_t(\boldsymbol{\alpha}) = L_{\boldsymbol{\alpha}}^* f_t(\boldsymbol{\alpha}), \qquad (27)$$

where the differential operator L^*_{α} is the adjoint of L_{α} , i.e.,

$$\langle L_{\alpha}g(\boldsymbol{\alpha}), h(\boldsymbol{\alpha})\rangle_{\boldsymbol{\alpha}} = \langle g(\boldsymbol{\alpha}), L_{\boldsymbol{\alpha}}^*h(\boldsymbol{\alpha})\rangle_{\boldsymbol{\alpha}},$$
 (28)

and satisfies

$$L^*_{\boldsymbol{\alpha}} f_t(\boldsymbol{\alpha}) = \langle \boldsymbol{\alpha} | \mathcal{L}^* F_t | \boldsymbol{\alpha} \rangle.$$
⁽²⁹⁾

Thus, L^*_{α} is the generator of the Koopman operator in the *P* representation describing the evolution of $f(\alpha)$, which corresponds to the adjoint Liouville operator \mathcal{L}^* in Eq. (19).

Corresponding to the Liouville operators \mathcal{L} and \mathcal{L}^* , the differential operators L_{α} and L^*_{α} also possess a biorthogonal eigensystem $\{\Lambda_k, u_k(\alpha), v_k(\alpha)\}_{k=0,1,2,...}$ of eigenvalue Λ_k and eigenfunctions u_k and v_k , satisfying

$$L_{\alpha}u_k = \Lambda_k u_k, \quad L_{\alpha}^* v_k = \overline{\Lambda_k} v_k, \quad \langle u_k, v_l \rangle_{\alpha} = \delta_{kl},$$
 (30)

which has one-to-one correspondence with Eq. (21). The eigenvalues $\{\Lambda_k\}_{k=0,1,2,...}$ are the same as those of \mathcal{L} , and the eigenfunctions u_k and v_k of L_{α} are related to the eigenoperators U_k and V_k of \mathcal{L} via

$$U_k = \int u_k(\boldsymbol{\alpha}) |\boldsymbol{\alpha}\rangle \langle \boldsymbol{\alpha} | \, d\boldsymbol{\alpha}, \quad v_k(\boldsymbol{\alpha}) = \langle \boldsymbol{\alpha} | V_k | \boldsymbol{\alpha}\rangle, \qquad (31)$$

which follow from

$$\mathcal{L}U_{k} = \int u_{k}(\boldsymbol{\alpha}) \left\{ \mathcal{L}|\alpha\rangle\langle\alpha| \right\} d\boldsymbol{\alpha} = \int \left\{ L_{\boldsymbol{\alpha}} u_{k}(\boldsymbol{\alpha}) \right\} |\alpha\rangle\langle\alpha| d\boldsymbol{\alpha}$$
$$= \int \Lambda_{k} u_{k}(\boldsymbol{\alpha}) |\alpha\rangle\langle\alpha| d\boldsymbol{\alpha} = \Lambda_{k} U_{k}$$
(32)

ARTICLE

and

$$L^*_{\alpha}v_k = L^*_{\alpha}\langle \alpha | V_k | \alpha \rangle = \langle \alpha | \mathcal{L}^* V_k | \alpha \rangle = \overline{\Lambda_k} \langle \alpha | V_k | \alpha \rangle = \overline{\Lambda_k} v_k.$$
(33)

C. Quantum asymptotic phase

Generalizing the definition for classical stochastic oscillatory systems in Sec. II, we here propose a definition of the quantum asymptotic phase. We note that, in quantum systems, the system state is given by the density operator ρ and individual trajectories as in the classical stochastic systems cannot be considered.

First, we define the quantum asymptotic phase $\Phi_q(\alpha)$ of the coherent state α in the *P* representation. Considering the definition Eq. (12) of the asymptotic phase in terms of the eigenfunction $Q_1(X)$ of the backward Fokker–Planck operator L_X^* in the classical stochastic case, we define the quantum asymptotic phase $\Phi_q(\alpha)$ as the argument of the complex conjugate of the eigenfunction $v_1(\alpha)$ in the *P* representation associated with the principal eigenvalue $\overline{\Lambda}_1$ as

$$\Phi_a(\boldsymbol{\alpha}) = \arg v_1(\boldsymbol{\alpha}) = \arg \langle \boldsymbol{\alpha} | V_1 | \boldsymbol{\alpha} \rangle. \tag{34}$$

Next, considering that the general quantum state ρ is represented as a superposition of coherent states with the weight $p(\alpha)$, Eq. (23), we define the asymptotic phase of ρ as

$$\Phi_{a}(\rho) = \arg \langle p(\boldsymbol{\alpha}), v_{1}(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}} = \arg \langle \rho, V_{1} \rangle_{tr}.$$
 (35)

It can be shown that the asymptotic phase $\Phi_q(\rho)$ evolves with a constant frequency Ω_q as the quantum state ρ evolves according to the master equation (15). Defining

$$\Psi_{q}(\rho_{t}) = \langle p_{t}(\boldsymbol{\alpha}), v_{1}(\boldsymbol{\alpha}) \rangle_{\boldsymbol{\alpha}} = \langle \rho_{t}, V_{1} \rangle_{tr}, \qquad (36)$$

where the dependence on time *t* is explicitly denoted, we have

$$\frac{d}{dt}\Psi_{q}(\rho_{t})\Big|_{t=t_{0}} = \left\langle \frac{\partial p_{t}(\boldsymbol{\alpha})}{\partial t}\Big|_{t=t_{0}}, v_{1}(\boldsymbol{\alpha})\right\rangle_{\boldsymbol{\alpha}} \\
= \left\langle L_{\boldsymbol{\alpha}}p_{t_{0}}(\boldsymbol{\alpha}), v_{1}(\boldsymbol{\alpha})\right\rangle_{\boldsymbol{\alpha}} \\
= \left\langle p_{t_{0}}(\boldsymbol{\alpha}), L_{\boldsymbol{\alpha}}^{*}v_{1}(\boldsymbol{\alpha})\right\rangle_{\boldsymbol{\alpha}} \\
= \left\langle p_{t_{0}}(\boldsymbol{\alpha}), \overline{\Lambda_{1}}v_{1}(\boldsymbol{\alpha})\right\rangle_{\boldsymbol{\alpha}} \\
= \overline{\Lambda_{1}}\Psi_{q}(\rho_{t_{0}}), \qquad (37)$$

or, equivalently,

$$\frac{d}{dt}\Psi_{q}(\rho_{t})\Big|_{t=t_{0}} = \langle \dot{\rho}_{t}|_{t=t_{0}}, V_{1} \rangle_{tr}$$

$$= \langle \mathcal{L}\rho_{t_{0}}, V_{1} \rangle_{tr} = \langle \rho_{t_{0}}, \mathcal{L}^{*}V_{1} \rangle_{tr}$$

$$= \langle \rho_{t_{0}}, \overline{\Lambda_{1}}V_{1} \rangle_{tr}$$

$$= \overline{\Lambda_{1}}\langle \rho_{t_{0}}, V_{1} \rangle_{tr}$$

$$= \overline{\Lambda_{1}}\Psi_{q}(\rho_{t_{0}}).$$
(38)

Integrating by time, we obtain

$$\Psi_q(\rho_t) = \exp(\Lambda_1 t) \Psi_q(\rho_0) = \exp[(\mu_q + i\Omega_q)t] \Psi_q(\rho_0), \quad (39)$$

where ρ_0 is the initial state at t=0 and, hence, the asymptotic phase is given by

$$\Phi_q(\rho_t) = \arg \Psi_q(\rho_t) = \Omega_q t + \arg \Psi_q(\rho_0).$$
(40)

Differentiating by *t*, we obtain

$$\frac{d}{dt}\Phi_q(\rho_t) = \Omega_q,\tag{41}$$

namely, the asymptotic phase $\Phi_q(\rho_t)$ increases with a constant frequency Ω_q with the evolution of the quantum state ρ_t .

Thus, by using the eigenfunction $v_1(\alpha)$ of the adjoint linear operator L^*_{α} or equivalently the eigenoperator V_1 of the adjoint operator \mathcal{L}^* associated with the eigenvalue $\overline{\Lambda_1}$, we can define the asymptotic phase $\Phi_q(\rho)$ of the quantum state ρ . The quantum master equation (15), adjoint Liouville operator \mathcal{L}^* (or adjoint differential operator L^*_{α} in the *P* representation), and eigenoperator V_1 [or the eigenfunction $v_1(\alpha)$ in the *P* representation] with the eigenvalue $\overline{\Lambda_1}$ correspond to the forward Fokker–Planck equation (5), backward Fokker–Planck operator L^*_X , and eigenfunction $Q_1(X)$ with the eigenvalue $\overline{\lambda_1}$ in the classical stochastic system discussed in Sec. II, respectively.

We stress, however, that the system state is generally represented by the density operator ρ or quasiprobability distribution $p(\alpha)$ and individual trajectories cannot be considered in the quantum case. As in the classical stochastic case, we may also choose the eigenvalue Λ_1 , eigenfunction $\overline{v_1(\alpha)}$, and eigenoperator V_1^{\dagger} to define the quantum asymptotic phase, which reverses its direction of increase.

IV. EXAMPLE: QUANTUM VAN DER POL OSCILLATOR

In this section, using the quantum van der Pol model with the quantum Kerr effect as an example, we illustrate the validity of the quantum asymptotic phase defined in Sec. III. We also analyze a damped harmonic oscillator in Appendix D.

A. Quantum van der Pol model with the quantum Kerr effect

As an example of quantum oscillatory systems, we consider the quantum van der Pol model with the quantum Kerr effect. The system's density operator ρ obeys the master equation

$$\dot{\rho} = \mathcal{L}\rho = -i[H,\rho] + \gamma_1 \mathcal{D}[a^{\dagger}]\rho + \gamma_2 \mathcal{D}[a^2]\rho, \qquad (42)$$

where *a* and a^{\dagger} are annihilation and creation operators, $H = \omega_0 a^{\dagger} a + K a^{\dagger 2} a^2$ is the Hamiltonian, ω_0 is the frequency parameter of the oscillator, *K* is the Kerr parameter, and γ_1 and γ_2 are the decay rates for negative damping and nonlinear damping, respectively.^{21,31} By using the standard rule of calculus,^{51,52,55} we can derive the differential operator L_{α} in Eq. (26) describing the evolution of the *P* representation $p(\alpha)$ of ρ from the Liouville operator \mathcal{L} , which consists of third- and higher-order derivative with respect to α .²¹

To evaluate the fundamental frequency Ω_q and the asymptotic phase $\Phi_q(\rho)$ of the system, we numerically calculate the eigenvalues and eigenfunctions of the Liouville and adjoint Liouville operators \mathcal{L} and \mathcal{L}^* . To this end, we approximately truncate the number representation of the density operator as a large $N \times N$ matrix and map it to a N^2 -dimensional vector of the double-ket notation.⁵⁶ We can then approximately represent the Liouville operators \mathcal{L} and \mathcal{L}^* by $N^2 \times N^2$ matrices and calculate their eigensystem to obtain the asymptotic phase in Eq. (34).

B. Semiclassical regime

We first consider the semiclassical regime where γ_2 and *K* are sufficiently small. In this case, as explained in Appendix B, we can approximate Eq. (26) by a Fokker–Planck equation for $p(\alpha)$, namely, the system state is approximately equivalent to a classical stochastic system. Furthermore, in the classical limit where the quantum noise vanishes, the system is described by a single complex variable $\alpha \in \mathbb{C}$ obeying a deterministic ordinary differential equation

$$\dot{\alpha} = \left(\frac{\gamma_1}{2} - i\omega_0\right)\alpha - (\gamma_2 + 2Ki)\overline{\alpha}\alpha^2.$$
(43)

This equation represents the Stuart–Landau oscillator (normal form of the supercritical Hopf bifurcation)⁴ and possesses a stable limitcycle solution $\alpha_0(\phi) = R e^{i\phi}$, which is represented as a function of the phase $\phi = \Omega_c t + \text{const.}$ with a natural frequency $\Omega_c = -\omega_0$ $- K\gamma_1/\gamma_2$ and radius $R = \sqrt{\gamma_1/2\gamma_2}$. The basin *B* of this limit cycle is the whole complex plane except the origin. The classical asymptotic phase Φ_c of this system is given by¹⁶

$$\Phi_c(\boldsymbol{\alpha}) = \arg \alpha - \frac{2K}{\gamma_2} \ln \frac{|\boldsymbol{\alpha}|}{R} + \text{const.}$$
(44)

and satisfies $\dot{\Phi}_c(\alpha) = \Omega_c$ as α evolves in *B* under Eq. (43).⁶ As stated in Sec. II, this $\Phi_c(\alpha)$ is the argument of the Koopman eigenfunction $\Psi_c(\alpha)$ of the system associated with the eigenvalue $i\Omega_c$. In Ref. 16, we



FIG. 1. Quantum asymptotic phase in the semiclassical regime. The parameters are $\gamma_1 = 1$ and $(\omega_0, \gamma_2, K)/\gamma_1 = (0.1, 0.05, 0.025)$. (a) Eigenvalues of \mathcal{L}_0 near the imaginary axis. The red dot represents the principal eigenvalue $\overline{\Lambda_1}$ with the slowest decay rate. (b) Quantum asymptotic phase Φ_q with $\Omega_q = -0.605$. (c) Classical asymptotic phase Φ_c with $\Omega_c = -0.6$. (d)–(g) Evolution of the expectation values of V_1 and *a* and their arguments from a pure coherent state: (d) $\Psi(\rho_t) = \langle V_1 \rangle_t$, (e) $\Phi(\rho_t) = \arg \langle V_1 \rangle_t$, (f) $\langle a \rangle_t$, and (g) $\arg \langle a \rangle_t$. In (a), individual branches of eigenvalues are shown with different colors. In (b) and (c), (x, p) = (2.5, 0) is chosen as the phase origin. In (c), the red-thin line represents the limit cycle in the classical limit.

used this Φ_c for the phase-reduction analysis of quantum synchronization in the semiclassical regime with weak quantum noise. It is expected that the quantum asymptotic phase $\Phi_q(\alpha)$ of the coherent state α is close to the classical asymptotic phase $\Phi_c(\alpha)$ when the quantum noise is sufficiently small.

Figure 1(a) shows the eigenvalues of \mathcal{L} near the imaginary axis obtained numerically, where the principal eigenvalue $\overline{\Lambda_1} = \mu_q$ $+ i\Omega_q$ is shown by a red dot ($\mu_q < 0$), and Figs. 1(b) and 1(c) compare the quantum-mechanical phase $\Phi_q(\alpha)$ with the corresponding classical phase $\Phi_c(\alpha)$. Here, we adopt a negative value for Ω_q so that the resulting phase Φ_q increases in the counterclockwise direction from 0 to 2π on the complex plane, i.e., Φ_q satisfies $\oint_c \nabla \Phi_q(\mathbf{x}) \cdot d\mathbf{x}$ $= 2\pi$ where $\mathbf{x} = (x, p) = (\text{Re } \alpha, \text{Im } \alpha)$ and *C* is a circle around 0.

As the quantum noise is small, the quantum frequency Ω_q and the asymptotic phase $\Phi_q(\boldsymbol{\alpha})$ obtained numerically from \mathcal{L} are close to the classical frequency Ω_c and asymptotic phase $\Phi_c(\boldsymbol{\alpha})$, respectively. The difference between Ω_q and Ω_c arises from the small quantum noise; in the limit of vanishing quantum noise, the eigenfunction $v_1(\boldsymbol{\alpha})$ of $L_{\boldsymbol{\alpha}}$ in the *P* representation coincides with the Koopman eigenfunction of the deterministic system Eq. (43) with the eigenvalue $i\Omega_c$ and, therefore, Φ_q reproduces the classical phase Φ_c (see Appendix C). Here, we note that the principal eigenvalues $i\Omega_q$ and $i\Omega_c$ are well separated from other branches of eigenvalues with faster decay rates and the corresponding oscillatory modes become quickly dominant.

To demonstrate that the present definition of the quantum asymptotic phase yields appropriate values, we consider free oscillatory relaxation of ρ_t from a coherent initial state $\rho^{\alpha_0} = |\alpha_0\rangle\langle\alpha_0|$ with $\alpha_0 = 1$ at t = 0 and measure $\Phi_q(\rho_t)$. For comparison, we also measure the argument arg $\langle a \rangle_t$ of the expectation $\langle a \rangle_t = \langle \rho_t, a \rangle_t$, of the annihilation operator a, which simply gives the polar angle of $\langle a \rangle_t$ on the complex plane. We note that, although the system state ρ starts from a pure coherent state, it soon becomes a mixed state due to the coupling with the reservoirs and eventually relaxes to the stationary state ρ^s .

Figures 1(d) and 1(e) plot the evolution of the expectation $\Psi_q(\rho_t) = \langle \rho_t, V_1 \rangle_{tr}$ of the eigenoperator V_1 and the quantum asymptotic phase $\Phi_q(\rho_t) = \arg \Psi_q(\rho_t)$, respectively, and Figs. 1(f) and 1(g) show the evolution of the expectation $\langle a \rangle$ and its polar angle $\arg\langle a \rangle$, respectively. The asymptotic phase $\Phi_q(\rho_t)$ increases with a constant frequency Ω_q and appropriately yields isochronous phase values. In contrast, the polar angle $\arg\langle a \rangle$ does not increase constantly with time, in particular, in the transient process before t = 10, as shown in Fig. 1(g); as the limit cycle in the classical limit is rotationally symmetric in this model, the polar angle also yields almost constantly increasing phase values after relaxation. Thus, the quantum asymptotic phase Φ_q increases with a constant frequency Ω_q in the semiclassical regime of a quantum van der Pol model with the quantum Kerr effect.

C. Strong quantum regime

Next, we consider a strong quantum regime with relatively large γ_2 and *K*, where only a small number of energy states participates in the system dynamics and the semiclassical description is not valid. The eigenvalues of \mathcal{L} obtained numerically are shown in Fig. 2(a). Figures 2(b) and 2(c) show the quantum-mechanical phase Φ_q and the corresponding classical phase Φ_c . Because the system is in the strong quantum regime, Φ_c is distinctly different from Φ_q and the classical frequency Ω_c also differs largely from the true quantum frequency Ω_q . Here, we again note that the principal eigenvalues have a much smaller (less than half of the second largest) decay rate than the eigenvalues in the other branches.

We consider free oscillatory relaxation of ρ from a coherent initial state $\rho = |\alpha_0\rangle \langle \alpha_0|$ with $\alpha_0 = 1$ at t = 0 and measure the evolution of the asymptotic phase $\Phi_q(\rho_t) = \langle V_1 \rangle_t$ of the system state ρ_t . For comparison, we also measure the polar angle $\arg \langle a \rangle_t$ of $\langle a \rangle_t$.

Figures 2(d) and 2(e) plot the evolution of the expectation $\Psi_q(\rho_t) = \langle \rho_t, V_1 \rangle_{tr}$ of the eigenoperator V_1 and the quantum asymptotic phase $\Phi_q(\rho_t) = \arg \Psi_q(\rho_t)$, respectively, and Figs. 2(f) and 2(g) show the evolution of the expectation $\langle a \rangle$ and its polar angle $\arg\langle a \rangle$, respectively. As expected, the asymptotic phase $\Phi_q(\rho_t)$ appropriately gives constantly varying phase values with the frequency Ω_q . In contrast, the polar angle $\arg\langle a \rangle_t$ does not vary constantly with time and is not isochronous. This is because the transition between relatively a small number of energy levels takes part in the system dynamics and the discreteness of the energy spectra can play important roles in this strong quantum regime.

Thus, the quantum asymptotic phase Φ_q gives appropriate phase values that increase with a constant frequency Ω_q even in the strong quantum regime of a quantum van der Pol model with the quantum Kerr effect, even though the strong quantum effect strongly alters the dynamics of the system from the classical limit.

D. Fundamental difference between the classical and quantum systems

Although we introduced the definition of the quantum asymptotic phase Φ_q by analogy with the classical deterministic phase Φ_c and classical stochastic asymptotic phase Φ_s , a fundamental difference exists between the quantum and classical cases. Specifically, in the quantum case, the system state is described by the density operator ρ and the asymptotic phase Φ_q assigns a phase value on each ρ , while in the classical case, the phase function Φ_c or Φ_s assigns a phase value on each individual state X, although the probability density function p(X) is used in defining the stochastic asymptotic phase Φ_s . This difference arises because the system state can be described by a SDE representing a single stochastic trajectory of the noisy oscillator in the classical stochastic case, whereas the system state can only be characterized by the density operator ρ representing the statistical state of the system in the quantum case.

Though we have taken the viewpoint that the asymptotic phase is defined for a single stochastic trajectory in the classical stochastic case in this study, we may also consider that the asymptotic phase is defined for the probability density $p(\mathbf{X})$ rather than for the state \mathbf{X} in the classical stochastic case. Then, the quantum asymptotic phase Φ_q as a function of the density matrix ρ corresponds to the stochastic asymptotic phase $\Phi_s(p(\mathbf{X})) = \arg \int p(\mathbf{X})Q_1(\mathbf{X}) d\mathbf{X}$ as a function of the probability density $p(\mathbf{X})$ in the classical stochastic case in Sec. II. We regarded this quantity as the exponential average of the asymptotic phase values defined for individual states in Eq. (13).



FIG. 2. Quantum asymptotic phase in the strong quantum regime. The parameters are $\gamma_1 = 0.1$ and $(\omega_0, \gamma_2, K)/\gamma_1 = (300, 4, 100)$. (a) Eigenvalues of \mathcal{L} near the imaginary axis. The red dot represents the principal eigenvalue $\overline{\Lambda_1}$ with the slowest decay rate. (b) Quantum asymptotic phase Φ_q with $\Omega_q = -30$. (c) Classical asymptotic phase Φ_c with $\Omega_c = -32.5$. (d)–(g) Evolution of the expectation values of V_1 and *a* and their arguments from a pure coherent state: (d) $\Psi(\rho_t) = \langle V_1 \rangle_t$, (e) $\Phi(\rho_t) = \arg\langle V_1 \rangle_t$, (f) $\langle a \rangle_t$, and (g) $\arg\langle a \rangle_t$. In (a), individual branches of eigenvalues are shown with different colors. In (b) and (c), (x, p) = (2.5, 0) is chosen as the phase origin. In (c), the red-thin line represents the limit cycle in the classical limit.

In the present approach, it is always possible to formally introduce a "phase function" for any oscillatory system as long as it has the decaying mode with a non-zero imaginary part by using the associated eigenoperator. However, this phase function does not necessarily play the role of a quantum asymptotic phase unless the system exhibits limit-cycle oscillations and synchronization. As an example, in Appendix D, we introduce a phase function of a damped quantum harmonic oscillator, which cannot be considered the quantum asymptotic phase since the system does not exhibit synchronization. Only when the system is nonlinear and exhibits synchronization, the asymptotic phase captures the synchronization dynamics of the system. The asymptotic phase is the basis for developing phase reduction theory for classical nonlinear oscillators. For quantum oscillatory systems, however, even if we can introduce the asymptotic phase as described in this study, it does not necessarily mean that we can develop the phase reduction theory. This is because the quantum state ρ may not be appropriately localized in the phasespace representation and, hence, it may not be reconstructed from the phase value even approximately. Still, as we showed for the case of the quantum vdP oscillator in the semiclassical regime,¹⁶ we may approximately describe the dynamics of the quantum state by using the reduced phase equation and perform a detailed synchronization analysis in appropriate physical situations. It should be noted that we also encounter a similar problem in developing phase reduction theory for classical oscillators under strong noise. We also note that, even if we cannot derive a reduced phase description for quantum nonlinear oscillators, the asymptotic phase can be used to characterize the peculiar properties of quantum synchronization.⁵⁷

V. SUMMARY

In this study, we proposed a definition of the asymptotic phase for quantum oscillatory systems by generalizing the asymptotic phase for the classical stochastic oscillatory system proposed by Thomas and Lindner¹ from the Koopman operator viewpoint.² The proposed asymptotic phase is defined by using the eigenoperator of the adjoint Liouville operator describing the evolution of the quantum-mechanical observable, in close analogy to the asymptotic phase for classical limit-cycle oscillators that can be interpreted as the argument of the Koopman eigenfunction associated with the fundamental frequency. By using the quantum van der Pol model with the quantum Kerr effect as an example, we demonstrated that the proposed asymptotic phase appropriately yields isochronous phase values even in the strong quantum regime where the semiclassical approximation is not valid.

Though quantum synchronization has attracted much attention recently, compared with classical synchronization, systematic analysis of the quantum synchronization has been restricted, partly due to the lack of the clear definition of the phase. The proposed definition of the asymptotic phase valid in strong quantum regimes may be used for systematic and quantitative analysis of synchronization phenomena in quantum nonlinear oscillators.⁵⁷ Moreover, we may be able to develop a phase reduction theory for strongly quantum nonlinear oscillators by using the proposed definition, which would allow us to reduce the system dynamics to a simple phase equation and facilitate detailed analysis, control, and optimization of quantum nonlinear oscillators. It will also be interesting to extend the definition of the amplitude functions for classical stochastic oscillators^{2,58} to strongly quantum oscillatory systems on the basis of the Koopman operator theory.

ACKNOWLEDGMENTS

The authors thank the anonymous reviewers for their enlightning comments. Numerical simulations have been performed by using QuTiP numerical toolbox.^{59,60} This research was funded by Japan Society for the Promotion of Science KAKENHI (Nos. JP17H03279, JP18H03287, JPJSBP120202201, JP20J13778, JP22K14274, JP22K11919, and JP22H00516) and Japan Science and Technology Agency Core Research for Evolutional Science and Technology (No. JP-MJCR1913).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflicts to disclose.

DATA AVAILABILITY

The data that support the findings of this study are available within the article.

APPENDIX A: KOOPMAN OPERATOR FOR CLASSICAL STOCHASTIC PROCESSES

In this section, we briefly summarize the definition of the Koopman operator for classical stochastic processes described by the Ito SDE (4). The evolution of an observable $g : \mathbb{R}^N \to \mathbb{C}$ for this Ito diffusion process from time t_0 to $t + t_0$ is expressed as^{47,61,62}

$$g_{t+t_0}(\mathbf{X}) = (U^t g_{t_0})(\mathbf{X}),$$
 (A1)

where U^t ($t \ge 0$) is the stochastic Koopman operator defined as

$$\begin{aligned} (U^{t}g)(Y) &= \mathbb{E}^{T}[g(X(t+t_{0}))] \\ &= \int g(X)p(X,t+t_{0}|Y,t_{0})dX = \langle p(X,t+t_{0}|Y,t_{0}),g(X)\rangle_{X}. \end{aligned}$$
(A2)

Here, $\mathbb{E}^{Y}[\cdot]$ represents the expectation over the stochastic realizations of X(t) started from $X(t_0) = Y$ at $t = t_0$, p(X, t|Y, s) ($t \ge s$) is the transition probability density, and $\langle \cdot, \cdot \rangle_X$ represents the inner product defined in Sec. II. Defining a time evolution operator S^t $= e^{tL_X}$ ($t \ge 0$), the evolution of the probability density p(X) is expressed as

$$p_{t+t_0}(\mathbf{X}) = (S^t p_{t_0})(\mathbf{X}),$$
 (A3)

where the operation of S^t on a function $f : \mathbb{R} \to \mathbb{C}$ is given by

$$(S^{t}f)(\boldsymbol{X}) = \int p(\boldsymbol{X}, t + t_{0} | \boldsymbol{Y}, t_{0}) f(\boldsymbol{Y}) \, d\boldsymbol{Y}.$$
 (A4)

The Koopman operator U^t is the adjoint of S^t , i.e.,

$$\langle f(\mathbf{X}), (U^{t}g)(\mathbf{X}) \rangle_{\mathbf{X}} = \int \overline{f(\mathbf{Y})} \left[\int g(\mathbf{X}) p(\mathbf{X}, t+t_{0}|\mathbf{Y}, t_{0}) \, d\mathbf{X} \right] d\mathbf{Y}$$

$$= \int \overline{\left[\int p(\mathbf{X}, t+t_{0}|\mathbf{Y}, t_{0}) f(\mathbf{Y}) \, d\mathbf{Y} \right]} g(\mathbf{X}) \, d\mathbf{X}$$

$$= \int \overline{(S^{t}f)(\mathbf{X})} g(\mathbf{X}) \, d\mathbf{X} = \langle (S^{t}f)(\mathbf{X}), g(\mathbf{X}) \rangle_{\mathbf{X}}$$
 (A5)

for two functions $f, g : \mathbb{R}^N \to \mathbb{C}$. It is noted that the expectation of the observable *g* at time $t + t_0$ can be expressed as

$$\int p_{t_0}(\mathbf{X})(U^t g_{t_0})(\mathbf{X}) d\mathbf{X} = \langle p_{t_0}(\mathbf{X}), (U^t g_{t_0})(\mathbf{X}) \rangle_{\mathbf{X}}$$
$$= \langle (S^t p_{t_0})(\mathbf{X}), g_{t_0}(\mathbf{X}) \rangle_{\mathbf{X}}$$
$$= \int (S^t p_{t_0})(\mathbf{X}) g_{t_0}(\mathbf{X}) d\mathbf{X}.$$
(A6)

It can be shown that the infinitesimal generator of U^t is given by the backward operator L_X^* in Eq. (8) (see, e.g., Ref. 62 for a rigorous treatment). From the Ito formula,^{49,62} the SDE for a function $g: \mathbb{R}^N \to \mathbb{R}$ is given by

$$dg(\mathbf{X}) = \left[\mathbf{A}(\mathbf{X}) \cdot \frac{\partial}{\partial \mathbf{X}} g(\mathbf{X}) + \frac{1}{2} \mathbf{D}(\mathbf{X}) \frac{\partial^2}{\partial \mathbf{X}^2} g(\mathbf{X}) \right] dt + \mathbf{B}(\mathbf{X}) \frac{\partial}{\partial \mathbf{X}} g(\mathbf{X}) d\mathbf{W},$$
(A7)

Chaos **32**, 063133 (2022); doi: 10.1063/5.0088559 Published under an exclusive license by AIP Publishing

ARTICLE

which gives

$$\mathbb{E}^{\mathbf{Y}}[g(\mathbf{X}(t+t_0))] = \int_{t_0}^{t+t_0} \left[\mathbf{A}(\mathbf{X}) \cdot \frac{\partial}{\partial \mathbf{X}} g(\mathbf{X}) + \frac{1}{2} \mathbf{D}(\mathbf{X}) \frac{\partial^2}{\partial \mathbf{X}^2} g(\mathbf{X}) \right] dt$$
$$= \int_{t_0}^{t+t_0} L_{\mathbf{X}}^* g(\mathbf{X}) dt \tag{A8}$$

by integration. Assuming the function *g* to be the observable g_{t_0} at $t = t_0$, the infinitesimal evolution of g_t at $t = t_0$ can be represented as

$$\frac{d}{dt}g_{t}(\mathbf{Y})\Big|_{t=t_{0}} = \lim_{t \to +0} \frac{(U^{t}g_{t_{0}})(\mathbf{Y}) - g_{t_{0}}(\mathbf{Y})}{t}$$
$$= \lim_{t \to +0} \frac{\mathbb{E}^{\mathbf{Y}}[g_{t_{0}}(\mathbf{X}(t+t_{0}))] - g_{t_{0}}(\mathbf{Y})}{t} = L_{\mathbf{X}}^{*}g_{t_{0}}(\mathbf{Y}).$$
(A9)

Thus, we can express the Koopman operator as $U^t = e^{tL_X^*}$.

From the adjoint relation for S^t and U^t , L_X and L_X^* are also adjoint to each other, i.e., $\langle L_X f(X), g(X) \rangle_X = \langle f(X), L_X^* g(X) \rangle_X$, and the evolution of the expectation of the observable *g* at time $t = t_0$ can be expressed as in Eq. (9).

APPENDIX B: QUANTUM VAN DER POL OSCILLATOR WITH KERR EFFECT IN THE SEMICLASSICAL REGIME

In this section, we briefly explain the classical limit of the quantum van der Pol oscillator. As shown in Ref. 16, in the semiclassical regime, the linear operator L_{α} in Eq. (26) describing the evolution of the quasiprobability distribution $p(\alpha)$ in the *P* representation of the quantum van der Pol oscillator can be approximated by a Fokker–Planck operator,

$$\tilde{L}_{\boldsymbol{\alpha}} = \left[-\sum_{j=1}^{2} \partial_{j} \{A_{j}(\boldsymbol{\alpha})\} + \frac{1}{2} \sum_{j=1}^{2} \sum_{k=1}^{2} \partial_{j} \partial_{k} \{D_{jk}(\boldsymbol{\alpha})\} \right], \quad (B1)$$

by neglecting the third- and higher-order derivatives, where $\partial_1 = \partial/\partial \alpha$ and $\partial_2 = \partial/\partial \bar{\alpha}$. The drift vector $A(\alpha) = (A_1(\alpha), A_2(\alpha)) \in \mathbb{C}^2$ and the matrix $D(\alpha) = (D_{jk}(\alpha)) \in \mathbb{C}^{2\times 2}$ are given by

$$A(\boldsymbol{\alpha}) = \begin{pmatrix} \left(\frac{\gamma_1}{2} - i\omega_0\right) \boldsymbol{\alpha} - (\gamma_2 + 2Ki)\overline{\boldsymbol{\alpha}} \boldsymbol{\alpha}^2\\ \left(\frac{\gamma_1}{2} + i\omega_0\right) \overline{\boldsymbol{\alpha}} - (\gamma_2 - 2Ki)\boldsymbol{\alpha} \overline{\boldsymbol{\alpha}}^2 \end{pmatrix},$$
(B2)

$$\boldsymbol{D}(\boldsymbol{\alpha}) = \begin{pmatrix} -(\gamma_2 + 2Ki)\alpha^2 & \gamma_1 \\ \gamma_1 & -(\gamma_2 - 2Ki)\bar{\alpha}^2 \end{pmatrix}.$$
(B3)

The corresponding stochastic differential equation is, thus, given by

$$d\begin{pmatrix}\alpha\\\overline{\alpha}\end{pmatrix} = \begin{pmatrix} \left(\frac{\gamma_1}{2} - i\omega_0\right)\alpha - (\gamma_2 + 2Ki)\overline{\alpha}\alpha^2\\ \left(\frac{\gamma_1}{2} + i\omega_0\right)\overline{\alpha} - (\gamma_2 - 2Ki)\alpha\overline{\alpha}^2 \end{pmatrix} dt + \boldsymbol{\beta}(\boldsymbol{\alpha}) \begin{pmatrix} dW_1\\ dW_2 \end{pmatrix},$$
(B4)

where W_1 and W_2 are independent Wiener processes and the matrix $\boldsymbol{\beta}(\boldsymbol{\alpha})$ is given by

$$\boldsymbol{\beta}(\boldsymbol{\alpha}) = \begin{pmatrix} \sqrt{\frac{(\gamma_1 + R_{11}(\boldsymbol{\alpha}))}{2}} e^{i\chi(\boldsymbol{\alpha})/2} & -i\sqrt{\frac{(\gamma_1 - R_{11}(\boldsymbol{\alpha}))}{2}} e^{i\chi(\boldsymbol{\alpha})/2} \\ \sqrt{\frac{(\gamma_1 - R_{11}(\boldsymbol{\alpha}))}{2}} e^{-i\chi(\boldsymbol{\alpha})/2} & i\sqrt{\frac{(\gamma_1 - R_{11}(\boldsymbol{\alpha}))}{2}} e^{-i\chi(\boldsymbol{\alpha})/2} \end{pmatrix}, \quad (B5)$$

where $R_{11}(\alpha) e^{i\chi(\alpha)} = -(\gamma_2 + 2Ki)\alpha^2$. It is noted that the two equations for α and $\overline{\alpha}$ in Eq. (B4) are mutually complex conjugate and represent the same dynamics.

In the classical limit, the deterministic part of Eq. (B4) gives the Stuart–Landau equation for the complex variable α given in Eq. (43), which is analytically solvable and the asymptotic phase $\Phi_c(\alpha)$ can be explicitly obtained as given in Eq. (44).^{4,6}

APPENDIX C: CLASSICAL LIMIT OF THE QUANTUM ASYMPTOTIC PHASE

In this section, we explain that the quantum asymptotic phase formally reproduces the deterministic asymptotic phase in the classical limit. In the semiclassical regime, the linear operator L_{α} of Eq. (26) for the quasiprobability distribution $p(\alpha)$ in the *P* representation can be approximated by a Fokker–Planck operator \tilde{L}_{α} of the form Eq. (B1). By introducing a real vector $\mathbf{X} = (\text{Re } \alpha, \text{Im } \alpha)$ and the corresponding probability density function $p(\mathbf{X})$, the Fokker–Planck operator \tilde{L}_{α} for $p(\alpha)$ can be cast into a real Fokker–Planck operator L_X for $p(\mathbf{X})$ in Eq. (7).

In the classical limit, the quantum noise vanishes and the diffusion term in L_X disappears. Thus, L_X formally becomes a classical Liouville operator, i.e., $L_X \rightarrow -(\partial/\partial X)A(X)$, and the corresponding backward Liouville operator formally becomes the infinitesimal generator of the deterministic Koopman operator, i.e., $L_X^* \rightarrow A = A(X) \cdot (\partial/\partial X) = A(X) \cdot \nabla$. Also, the decay rate μ_q approaches 0 and the eigenvalue Λ_q approaches $i\Omega_c$, where Ω_c is the frequency of the limiting classical deterministic system. As discussed in Sec. II A, the classical asymptotic phase Φ_c is obtained as the argument of the eigenfunction Ψ_c of A associated with the eigenvalue $\overline{\Lambda_1} = i\Omega_q$. Thus, the quantum asymptotic phase Φ_c in the classical limit with vanishing quantum noise.

Figure 3 schematically shows the behavior of the eigenvalues of L_X^* approaching those of the deterministic system in the classical limit with a stable limit-cycle solution. The eigenvalues in the classical limit are given in the form $\lambda_c = m\kappa_1 + in\omega_1$ (m = 0, 1, 2, ... and $n = 0, \pm 1, \pm 2$), where κ_1 is the real part of the largest non-zero eigenvalue and ω_1 is the imaginary part of the pure-imaginary eigenvalue with the smallest absolute imaginary part. In the example of a quantum van der Pol oscillator with the quantum Kerr effect in





Chaos **32**, 063133 (2022); doi: 10.1063/5.0088559 Published under an exclusive license by AIP Publishing Sec. IV, $\kappa_1 = -\gamma_1$ and $\omega_1 = \Omega_c$. As the system approaches the classical limit, each curved branch of eigenvalues Λ_q of L_X in Fig. 3(a) approaches the corresponding straight branch of eigenvalues λ_c in Fig. 3(b) in the classical limit.

The above formal correspondence with the conventional definition of the asymptotic phase in the classical limit supports the validity of our definition of the quantum asymptotic phase.

APPENDIX D: PHASE FUNCTION OF A QUANTUM DAMPED HARMONIC OSCILLATOR

In Ref. 63, Thomas and Lindner considered the stochastic phase function for a classical damped harmonic oscillator described by a multi-dimensional Ornstein-Uhlenbeck process. In this section, generalizing their result, we consider a simple quantum harmonic oscillator with damping and formally calculate the phase function. This system is linear and does not possess a limit cycle in the classical limit, but the isochronous phase function as defined in Sec. III can still be introduced (as the system lacks an asymptotic periodic orbit in this case, we do not use the term "asymptotic").

The eigenoperator V_1 of the adjoint Liouville operator \mathcal{L}^* can be analytically obtained in this case. The evolution of a damped harmonic oscillator is described by a quantum master equation

$$\dot{\rho} = \mathcal{L}\rho = -i[\omega a^{\dagger}a, \rho] + \gamma \mathcal{D}[a]\rho, \tag{D1}$$

where ω is the natural frequency of the system, γ denotes the decay rate for the linear damping, and \mathcal{D} is the Lindblad form.⁵¹ The eigenoperator associated with the slowest non-vanishing decay rate of the adjoint Liouville operator \mathcal{L}^* of \mathcal{L} is simply given by $V_1 = a$, i.e., $\mathcal{L}^* a = \overline{\Lambda_1} a$, where $\overline{\Lambda_1} = -\gamma/2 - i\omega$.^{64,65} Therefore, the phase function $\Phi_q(\boldsymbol{\alpha})$ of the coherent state $\boldsymbol{\alpha}$ is given by

$$\Phi_q(\boldsymbol{\alpha}) = \arg\langle \rho_t, a \rangle_{tr} = \arg\langle \alpha | a | \alpha \rangle = \arg \alpha = \arg \left(r e^{i\theta} \right) = \theta,$$
 (D2)

where $\alpha = re^{i\theta}$, and the phase function $\Phi_q(\rho_t)$ of the density operator ρ_t at time t is given by

$$\Phi_q(\rho_t) = \arg\langle a \rangle_t = \arg\langle \rho_t, a \rangle_{tr}.$$
 (D3)

For the initial condition $\rho_0 = |\alpha_0\rangle \langle \alpha_0|$ with $\alpha_0 = r_0 e^{i\theta_0}$, the expectation of a evolves as

$$\begin{aligned} \frac{d}{dt} \langle a \rangle_t \Big|_{t=t_0} &= \langle \dot{\rho}_t |_{t=t_0}, a \rangle_{tr} = \langle \mathcal{L} \rho_{t_0}, a \rangle_{tr} = \langle \rho_{t_0}, \mathcal{L}^* a \rangle_{tr} \\ &= \overline{\Lambda_1} \langle \rho_{t_0}, a \rangle_{tr} = \overline{\Lambda_1} \langle a \rangle_{t}, \end{aligned}$$
(D4)

which gives $\langle a \rangle_t = e^{\overline{\Lambda_1}t} \langle a \rangle_0 = e^{\overline{\Lambda_1}t} \langle \alpha_0 | a | \alpha_0 \rangle = e^{\overline{\Lambda_1}t} \alpha_0$. Thus, the phase of the state ρ_t is given by

$$\Phi_q(\rho_t) = \arg(e^{(-\gamma/2 - i\omega)t}\alpha_0) = -\omega t + \theta_0, \tag{D5}$$

which decreases with a constant frequency ω .

As shown in this example of a quantum damped harmonic oscillator, we can formally introduce the phase function for a wide class of oscillators, even if the system does not exhibit limit-cycle dynamics. In a special case where the system exhibits limit-cycle dynamics, our definition of the phase function plays the role of the quantum asymptotic phase.

REFERENCES

¹P. J. Thomas and B. Lindner, "Asymptotic phase for stochastic oscillators," Phys. Rev. Lett. 113(25), 254101 (2014).

²Y. Kato, J. Zhu, W. Kurebayashi, and H. Nakao, "Asymptotic phase and amplitude for classical and semiclassical stochastic oscillators via Koopman operator theory," Mathematics 9(18), 2188 (2021).

³A. T. Winfree, The Geometry of Biological Time (Springer, New York, 2001).

⁴Y. Kuramoto, Chemical Oscillations, Waves, and Turbulence (Springer, Berlin, 1984).

⁵A. Pikovsky, M. Rosenblum, and J. Kurths, Synchronization: A Universal Concept in Nonlinear Sciences (Cambridge University Press, Cambridge, 2001).

⁶H. Nakao, "Phase reduction approach to synchronisation of nonlinear oscillators," Contemp. Phys. 57(2), 188-214 (2016).

⁷G. B. Ermentrout and D. H. Terman, Mathematical Foundations of Neuroscience (Springer, New York, 2010).

⁸S. H. Strogatz, Nonlinear Dynamics and Chaos (Westview Press, 1994).

⁹M. H. Matheny, J. Emenheiser, W. Fon, A. Chapman, A. Salova, M. Rohden, J. Li, M. Hudoba de Badyn, M. Pósfai, L. Duenas-Osorio, M. Mesbahi, J. P. Crutchfield, M. C. Cross, R. M. Dsouza, and M. L. Roukes, "Exotic states in a simple network of nanoelectromechanical oscillators," Science 363(6431), eaav7932 (2019).

10S. Kreinberg, X. Porte, D. Schicke, B. Lingnau, C. Schneider, S. Höfling, I. Kanter, K. Lüdge, and S. Reitzenstein, "Mutual coupling and synchronization of optically coupled quantum-dot micropillar lasers at ultra-low light levels," Nat. Commun. 10(1), 137 (2019).

¹¹H. Singh, S. Bhuktare, A. Bose, A. Fukushima, K. Yakushiji, S. Yuasa, H. Kubota, and A. A. Tulapurkar, "Mutual synchronization of spin-torque nano-oscillators via Oersted magnetic fields created by waveguides," Phys. Rev. Appl. 11(5), 054028 (2019).

¹²M. F. Colombano, G. Arregui, N. E. Capuj, A. Pitanti, J. Maire, A. Griol, B. Garrido, A. Martinez, C. M. Sotomayor-Torres, and D. Navarro-Urrios, "Synchronization of optomechanical nanobeams by mechanical interaction," Phys. Rev. Lett. 123(1), 017402 (2019).

¹³T. E. Lee and H. R. Sadeghpour, "Quantum synchronization of quantum van der Pol oscillators with trapped ions," Phys. Rev. Lett. 111(23), 234101 (2013).

¹⁴S. Walter, A. Nunnenkamp, and C. Bruder, "Quantum synchronization of a driven self-sustained oscillator," Phys. Rev. Lett. 112(9), 094102 (2014).

¹⁵S. Sonar, M. Hajdušek, M. Mukherjee, R. Fazio, V. Vedral, S. Vinjanampathy, and L.-C. Kwek, "Squeezing enhances quantum synchronization," Phys. Rev. Le 120(16), 163601 (2018).

¹⁶Y. Kato, N. Yamamoto, and H. Nakao, "Semiclassical phase reduction theory for quantum synchronization," Phys. Rev. Res. 1, 033012 (2019).

17 W.-K. Mok, L.-C. Kwek, and H. Heimonen, "Synchronization boost with singlephoton dissipation in the deep quantum regime," Phys. Rev. Res. 2(3), 033422 (2020).

¹⁸T. E. Lee, C.-K. Chan, and S. Wang, "Entanglement tongue and quantum synchronization of disordered oscillators," Phys. Rev. E 89(2), 022913 (2014). ¹⁹D. Witthaut, S. Wimberger, R. Burioni, and M. Timme, "Classical synchro-

nization indicates persistent entanglement in isolated quantum systems," Nat. Commun. 8, 14829 (2017).

²⁰A. Roulet and C. Bruder, "Quantum synchronization and entanglement generation," Phys. Rev. Lett. 121(6), 063601 (2018).

²¹N. Lörch, E. Amitai, A. Nunnenkamp, and C. Bruder, "Genuine quantum signatures in synchronization of anharmonic self-oscillators," Phys. Rev. Lett. 117(7), 073601 (2016).

22 N. Lörch, S. E. Nigg, A. Nunnenkamp, R. P. Tiwari, and C. Bruder, "Quantum synchronization blockade: Energy quantization hinders synchronization of identical oscillators," Phys. Rev. Lett. 118(24), 243602 (2017).

²³S. E. Nigg, "Observing quantum synchronization blockade in circuit quantum electrodynamics," Phys. Rev. A 97(1), 013811 (2018).

²⁴T. Weiss, A. Kronwald, and F. Marquardt, "Noise-induced transitions in optomechanical synchronization," New J. Phys. **18**(1), 013043 (2016). ²⁵N. Es' haqi Sani, G. Manzano, R. Zambrini, and R. Fazio, "Synchronization

along quantum trajectories," Phys. Rev. Res. 2(2), 023101 (2020).

²⁶Y. Kato and H. Nakao, "Enhancement of quantum synchronization via continuous measurement and feedback control," New J. Phys. 23(1), 013007 (2021).

²⁷Y. Kato and H. Nakao, "Instantaneous phase synchronization of two decoupled quantum limit-cycle oscillators induced by conditional photon detection," Phys. Rev. Res. 3(1), 013085 (2021).

²⁸W. Li, N. Es'haqi Sani, W.-Z. Zhang, and D. Vitali, "Quantum zeno effect in self-sustaining systems: Suppressing phase diffusion via repeated measurements," Phys. Rev. A **103**(4), 043715 (2021).

²⁹A. W. Laskar, P. Adhikary, S. Mondal, P. Katiyar, S. Vinjanampathy, and S. Ghosh, "Observation of quantum phase synchronization in spin-1 atoms," Phys. Rev. Lett. **125**(1), 013601 (2020).

³⁰M. Koppenhöfer, C. Bruder, and A. Roulet, "Quantum synchronization on the IBM Q system," Phys. Rev. Res. **2**(2), 023026 (2020).

³¹Y. Kato and H. Nakao, "Semiclassical optimization of entrainment stability and phase coherence in weakly forced quantum limit-cycle oscillators," Phys. Rev. E **101**(1), 012210 (2020).

³² M. R. Hush, W. Li, S. Genway, I. Lesanovsky, and A. D. Armour, "Spin correlations as a probe of quantum synchronization in trapped-ion phonon lasers," Phys. Rev. A **91**(6), 061401 (2015).

³³T. Weiss, S. Walter, and F. Marquardt, "Quantum-coherent phase oscillations in synchronization," Phys. Rev. A **95**(4), 041802 (2017).

³⁴A. Mari, A. Farace, N. Didier, V. Giovannetti, and R. Fazio, "Measures of quantum synchronization in continuous variable systems," Phys. Rev. Lett. **111**(10), 103605 (2013).

³⁵M. Xu, D. A. Tieri, E. C. Fine, J. K. Thompson, and M. J. Holland, "Synchronization of two ensembles of atoms," Phys. Rev. Lett. **113**(15), 154101 (2014).

³⁶A. Roulet and C. Bruder, "Synchronizing the smallest possible system," Phys. Rev. Lett. **121**(6), 053601 (2018).

³⁷A. Chia, L. C. Kwek, and C. Noh, "Relaxation oscillations and frequency entrainment in quantum mechanics," Phys. Rev. E **102**(4), 042213 (2020).

³⁸L. Ben Arosh, M. C. Cross, and R. Lifshitz, "Quantum limit cycles and the Rayleigh and van der Pol oscillators," Phys. Rev. Res. 3(1), 013130 (2021).

³⁰N. Jaseem, M. Hajdušek, P. Solanki, L.-C. Kwek, R. Fazio, and S. Vinjanampathy, "Generalized measure of quantum synchronization," Phys. Rev. Res. 2(4), 043287 (2020).

⁴⁰N. Jaseem, M. Hajdušek, V. Vedral, R. Fazio, L.-C. Kwek, and S. Vinjanampathy, "Quantum synchronization in nanoscale heat engines," Phys. Rev. E 101(2), 020201 (2020).

⁴¹A. Cabot, G. Luca Giorgi, and R. Zambrini, "Metastable quantum entrainment," New J. Phys. 23(10), 103017 (2021).

⁴²A. Mauroy, I. Mezić, and J. Moehlis, "Isostables, isochrons, and Koopman spectrum for the action–angle representation of stable fixed point dynamics," Physica D 261, 19–30 (2013).

⁴³Y. S. Mauroy and I. Mezic, *The Koopman Operator in Systems and Control* (Springer, 2020).

⁴⁴S. Shirasaka, W. Kurebayashi, and H. Nakao, "Phase-amplitude reduction of transient dynamics far from attractors for limit-cycling systems," Chaos **27**(2), 023119 (2017).

⁴⁵Y. Kuramoto and H. Nakao, "On the concept of dynamical reduction: The case of coupled oscillators," Philos. Trans. R. Soc. A 377(2160), 20190041 (2019).

⁴⁶O. Junge, J. E. Marsden, and I. Mezic, "Uncertainty in the dynamics of conservative maps," in 2004 43rd IEEE Conference on Decision and Control (CDC) (IEEE Cat. No. 04CH37601) (IEEE, 2004), Vol. 2, pp. 2225–2230.

⁴⁷N. Črnjarić-Žic, S. Maćešić, and I. Mezić, "Koopman operator spectrum for random dynamical systems," J. Nonlinear Sci. 30(5), 2007–2056 (2020).

⁴⁸M. T. Wanner, *Robust Approximation of the Stochastic Koopman Operator* (University of California, Santa Barbara, 2020).

49 C. Gardiner, Stochastic Methods (Springer, Berlin, 2009), Vol. 4.

⁵⁰D. A. Lidar, I. L. Chuang, and K. B. Whaley, "Decoherence-free subspaces for quantum computation," Phys. Rev. Lett. 81(12), 2594 (1998).
 ⁵¹H. J. Carmichael, Statistical Methods in Quantum Optics 1, 2 (Springer, New

⁵¹ H. J. Carmichael, *Statistical Methods in Quantum Optics 1, 2* (Springer, New York, 2007).

52 C. W. Gardiner, Quantum Noise (Springer, New York, 1991).

⁵³H.-P. Breuer *et al.*, *The Theory of Open Quantum Systems* (Oxford University Press on Demand, 2002).

⁵⁴A. C. Y. Li, F. Petruccione, and J. Koch, "Perturbative approach to Markovian open quantum systems," Sci. Rep. 4, 1301 (2015).
 ⁵⁵K. E. Cahill and R. J. Glauber, "Density operators and quasiprobability distribu-

⁵⁵K. E. Cahill and R. J. Glauber, "Density operators and quasiprobability distributions," Phys. Rev. 177(5), 1882 (1969).

⁵⁶V. V. Albert, "Lindbladians with multiple steady states: Theory and applications," arXiv:1802.00010 (2018).

⁵⁷Y. Kato and H. Nakao, "Quantum asymptotic phase reveals signatures of quantum synchronization," arXiv:2006.00760 (2020).
 ⁵⁸A. Pérez-Cervera, B. Lindner, and P. J. Thomas, "Isostables for stochastic

⁵⁸A. Pérez-Cervera, B. Lindner, and P. J. Thomas, "Isostables for stochastic oscillators," Phys. Rev. Lett. **127**(25), 254101 (2021).

⁵⁹J. R. Johansson, P. D. Nation, and F. Nori, "QuTiP: An open-source Python framework for the dynamics of open quantum systems," Comput. Phys. Commun. 183(8), 1760–1772 (2012).

⁶⁰ J. R. Johansson, P. D. Nation, and F. Nori, "QuTiP 2: A Python framework for the dynamics of open quantum systems," Comput. Phys. Commun. 184, 1234–1240 (2013).

⁶¹I. Mezić, "Spectral properties of dynamical systems, model reduction and decompositions," Nonlinear Dyn. **41**(1-3), 309–325 (2005).

⁶²B. Oksendal, Stochastic Differential Equations: An Introduction with Applications (Springer Science & Business Media, 2013).

⁶³P. J. Thomas and B. Lindner, "Phase descriptions of a multidimensional Ornstein-Uhlenbeck process," Phys. Rev. E **99**(6), 062221 (2019).

⁶⁴S. M. Barnett and S. Stenholm, "Spectral decomposition of the Lindblad operator," J. Mod. Opt. **47**(14-15), 2869–2882 (2000).

⁶⁵H.-J. Briegel and B.-G. Englert, "Quantum optical master equations: The use of damping bases," Phys. Rev. A **47**(4), 3311 (1993).