

論文 / 著書情報
Article / Book Information

題目(和文)	
Title(English)	Elucidating the effect of Valproic Acid on the differentiation of iPSC-derived enterocyte-like cells
著者(和文)	Leo Han Yun Sylvia
Author(English)	Leo Han Yun Sylvia
出典(和文)	学位:博士(学術), 学位授与機関:東京工業大学, 報告番号:甲第12595号, 授与年月日:2023年9月22日, 学位の種別:課程博士, 審査員:糸 昭苑,白木 伸明,山本 直之,山田 拓司,田川 陽一
Citation(English)	Degree:Doctor (Academic), Conferring organization: Tokyo Institute of Technology, Report number:甲第12595号, Conferred date:2023/9/22, Degree Type:Course doctor, Examiner:,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	論文要旨
Type(English)	Summary

論文要旨

THESIS SUMMARY

系・コース： Department of, Graduate major in	Life Science and Technology	系 コース	申請学位（専攻分野）： Academic Degree Requested	博士 Doctor of	(Philosophy)
学生氏名： Student's Name	Leo Han Yun Sylvia		指導教員（主）： Academic Supervisor(main)	Kume Shoen	
			指導教員（副）： Academic Supervisor(sub)	Shiraki Nobuaki	

要旨（和文 2000 字程度）

Thesis Summary (approx.2000 Japanese Characters)

<u>Introduction</u> Despite being highly regenerative in vivo, primary human intestine epithelial cells undergo rapid apoptosis in vitro and cannot persist in long-term culture, resulting in the need for a suitable substitute for drug development and disease modelling. Highly proliferative human induced pluripotent stem cells (iPSCs) have been successfully differentiated into all major intestine epithelial cell types and provide an alternative to primary cells. In Ogaki et al., 2013, Kume-Shiraki laboratory reported the differentiation of human embryonic stem cells (hESCs) into the four major intestine epithelial cell types induced by BIO and DAPT treatment on the M15 feeder layer, confirming that Wnt and Notch signals governed the differentiation of hESC into intestinal lineages. This protocol was then adapted into a directed differentiation system for enterocytes with the addition of maturation medium M3 with BIO, Dexamethasone, DAPT and activated Vitamin D3 (VD3) to improve drug transporter and metabolic enzyme gene expression and functionality on the novel Collagen Vitrigel Membrane (CVM) in a culture system specialized for pharmacokinetic studies. However, aside from drug transporter and xenobiotic-focused metabolic enzyme functionality, other intestine-related functions were not looked at, nor were the specific changes of inducer VD3 on our cells investigated. In this dissertation, bulk RNA-sequencing methods were employed to investigate the changes in gene expression induced by metabolic enzyme Cytochrome P40 (CYP) inducer VD3, as well as the effects of Notch activator and known CYP modulator Valproic Acid (VPA) in promoting function-related gene expression and maturity of our iPSC-derived enterocytes. <u>Results and Discussion</u> 1. Notch, Wnt and BMP signaling regulate the differentiation of iPSCs into absorptive enterocyte-like cells To analyse if the differentiation process of iPSC-derived enterocytes in our established protocol echoes that of in vivo processes, qPCR and bulk RNA sequencing analysis was carried out on samples at undifferentiated (D0), precursor (D10, D15) and mature (D23) samples. We confirmed that the in vitro differentiation of iPSCs to iPSC-derived enterocyte-like cells on CVM demonstrated the expected functions as reported in developmental studies, suggesting that iPSCs provide a suitable platform to study the differentiation process of enterocytes in vivo. In summary, genes regulating general cellular processes are downregulated upon differentiation, whilst intestine-specific gene expressions are upregulated upon differentiation, and heatmaps illustrate the obvious changes in Notch, BMP and Wnt pathways throughout the differentiation timeline. Whilst BMP signaling promotes enterocyte maturation, Notch signaling could also promote enterocyte differentiation by inhibiting transcription factors for secretory fates.

備考：論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note：Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1copy of 800 Words (English).

注意：論文要旨は、東工大リサーチリポジトリ(T2R2)にてインターネット公表されますので、公表可能な範囲の内容で作成してください。

Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).

論文要旨

THESIS SUMMARY

系・コース： Department of, Graduate major in	Life Science and Technology	系 コース	申請学位 (専攻分野)： Academic Degree Requested	博士 Doctor of	(Philosophy)
学生氏名： Student's Name	Leo Han Yun Sylvia		指導教員 (主)： Academic Supervisor(main)	Kume Shoen	
			指導教員 (副)： Academic Supervisor(sub)	Shiraki Nobuaki	

要旨 (英文 300 語程度)

Thesis Summary (approx.300 English Words)

2. VD3 induces CYP3A4 and functional specification in iPSC-derived enterocyte-like cells
The inclusion of VD3 in M3 maturation media not only induces the expression and activity of CYP3A4, but also has the effect of promoting enterocyte functional specification to that characteristic of villus-tip enterocytes, likely due to BMP-signaling mediated by VD3. BMP signaling directly regulates the functionality of mouse enterocytes, upregulating fatty acid metabolism-related genes apolipoproteins and downregulating and sugar metabolizing genes (Buemer et al., 2022). These reports were in-line with our results, where VD3 enriches Gene Ontology (GO) categories related to fatty acid metabolism, such as bile transporter activity, digestive processes, and lipid translocation. We also noted a significant upregulation of *BMP2* transcript levels, a potent modulator of the apolipoprotein-coding genes, as well as modulator *SMAD4* and BMP receptor *BMPR2* with VD3 treatment.

3. VPA activates Notch signaling and upregulates expression of intestine function-related genes
We next attempted to elucidate the effect of VPA, an HDAC inhibitor, Notch activator and known CYP3A4 regulator on the maturation of iPSC-derived enterocytes in M3 medium. VPA was found not to exert much significant effect in the absence of VD3 but worked synergistically with VD3 to significantly improve gene expression of intestine functional genes such as those in GO categories aquaporins, fatty-acid degradation and vitamin digestion and absorption. VPA treatment in tandem with VD3 also upregulated key regulators of enterohepatic circulation such as *FGF19* and basolateral bile acid transporters *SLC51A* and *SLC51B*. Interestingly, VPA treatment instead downregulated expression of apical bile acid uptake transporter *SLC10A2*, which could be attributed to increased sensitivity of iPSC-derived enterocytes to bile acid levels in culture medium as a result of enriched levels of regulator *FGF19*. To sum up, we were able to ascertain the genetic profile of our D23 iPSC-derived enterocyte-like cells to be specialized towards fatty acid absorption and metabolism instead of absorption and metabolism of carbohydrates and other nutrients. These results suggest a possibility for our VD3VPA-treated iPSC-derived enterocyte-like cells for use in a co-culture model with hepatocytes to model enterohepatic circulation *in vivo*.

備考：論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note：Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1copy of 800 Words (English).

注意：論文要旨は、東工大リサーチリポジトリ(T2R2)にてインターネット公表されますので、公表可能な範囲の内容で作成してください。
Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).