

論文 / 著書情報  
Article / Book Information

題目(和文)	開放量子系におけるパリティ・時間対称性由来 の散逸連続時間結晶
Title(English)	Dissipative continuous time crystals originating from parity-time symmetry in open quantum systems
著者(和文)	中西優馬
Author(English)	Yuma Nakanishi
出典(和文)	学位:博士(理学), 学位授与機関:東京工業大学, 報告番号:甲第12640号, 授与年月日:2024年3月26日, 学位の種別:課程博士, 審査員:笹本 智弘,村上 修一,西田 祐介,石塚 大晃,納富 雅也
Citation(English)	Degree:Doctor (Science), Conferring organization: Tokyo Institute of Technology, Report number:甲第12640号, Conferred date:2024/3/26, Degree Type:Course doctor, Examiner:,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	要約
Type(English)	Outline

(博士課程)  
Doctoral Program

## 論文要約

THESIS OUTLINE

系・コース： Department of, Graduate major in	物理学 物理学	系 コース	申請学位 (専攻分野)： Academic Degree Requested	博士 Doctor of	(理学)
学生氏名： Student's Name	中西 優馬		審査員主査： Chief Examiner	笹本 智弘	

### 要約

Exploration of novel non-equilibrium phases of matter or phase transitions and clarification of their mechanisms are important problems for non-equilibrium statistical physics. In particular, (continuous) time crystals that spontaneously break continuous time-translation symmetry into discrete time-translation symmetry are a typical example, and these have been observed in open quantum systems but proved to be impossible to exist in closed quantum systems. Here, continuous time crystals in open quantum systems are called dissipative continuous time crystals. In this thesis, we provide novel non-equilibrium phase transitions with anti-unitary symmetry breaking in Markovian open quantum systems and show that they can explain the mechanism of a class of dissipative continuous time crystals in open collective spin systems.

First, we give a novel phase transition similar to the parity-time (PT) phase transitions, which are studied in non-Hermitian systems well, for investigating a two-collective spin system with balanced gain and loss described by the GKSL (Gorini-Kossakowski-Sudarshan-Lindblad) equation. Here, a generator of the GKSL equation is called the Lindbladian. In the non-Hermitian PT phase transitions, for the Hamiltonian with the PT symmetry, the eigenvalues change from real to complex eigenvalues, thereby changing the dynamics from periodic oscillations to diverging or decaying dynamics, and the PT symmetry of the eigenstate is broken at the same parameter point. On the other hand, for the Lindbladian case, we show that when the Lindbladian has a PT symmetry, the Lindbladian spectrum transition, dynamics transition, and PT symmetry breaking of the stationary state occur at the same parameter point. Here, we use the Holstein-Promakoff approximation, which linearly maps the collective spin operators to bosonic operators, and the third quantization, which is the prescription to diagonalize the Lindbladian with quadratic Hamiltonian and linear Lindblad operators. Also, in the non-Hermitian PT phase transition, the transition point is known as the exceptional point where more than two eigenvectors coalesce. We will explain that the transition point is expected to be an exceptional point in the Lindbladian case as well. These properties are similar to those of non-Hermitian PT phase transitions. This dissipative phase transition is called the Lindbladian PT phase transition. Then, we can define the Lindbladian PT phase as satisfying the following three properties: (i) the model has a strong Lindbladian PT symmetry, (ii) purely imaginary eigenvalues exist in the thermodynamic limit, and (iii) the PT symmetry of a stationary state holds in the thermodynamic limit.

Next, we show the physical origin of the boundary time crystal (BTC) for specific models, which is a dissipative time crystal only in the thermodynamic limit. In particular, we focus on the dissipative one-collective spin model with transverse magnetic field and excitation loss (hereafter, we refer to this model as the Driven Dicke model). In the Driven Dicke model, the exact solution of the stationary state had already been found in 1979, and the static dissipative continuous phase transition was known to occur. However, the mechanism of this phase transition is unclear. Notably, this continuous phase transition has been recently observed in a cold atom gas system. Also, the persistent oscillation appearing in this model was reaffirmed as the BTC in 2018, and this mechanism is also unclear.

In this thesis, we analytically show that the Driven Dicke model has the Lindbladian PT symmetry when we choose the parity operator to reflect the basis of Sz. Physically speaking, it means that the Driven Dicke model returns to itself after transforming the parity and time reversal operations. Moreover, we analytically show that the PT symmetry breaking of the stationary state occurs at the point where BTC breaks down. Also, we numerically show a similar statement for another BTC (generalized Driven Dicke) model. Here, we introduce the PT symmetry parameter that characterizes the PT symmetry of the density matrix. Finally, using perturbation theory with degeneracy, we prove that BTC arises under weak dissipation for a class of one-collective spin systems only when gain and loss are balanced. These results indicate that there exist BTCs in collective spin systems originating from parity-time symmetry.

Furthermore, we give a simple and clear description of BTCs and the Lindbladian PT phase transitions. Our strategy is that the Schwinger boson transformation makes it easier to discuss the effect of PT symmetry, even at the mean-field level. In particular, we show that when the Lindbladian has PT symmetry at the microscopic level, the Schwinger boson mean-field equation becomes a nonlinear Schrödinger equation with U(1) symmetry. Moreover, we prove that if a Lindbladian has PT symmetry at the microscopic level, the effective nonlinear Hamiltonian also has PT symmetry for the general

collective spin system. Also, the continuous and discontinuous phase transitions can occur with spontaneous non-linear PT symmetry breaking. Then, we demonstrate that all collective excitation modes are shown to be destabilized at the transition point using the linear stability analysis. As a result, the time-dependent phases appear, such as the swap or chiral phases, in which the amplitude or phase periodically oscillates respectively. In particular, the phase where amplitudes are time-dependent (e.g., swap phase) corresponds to the time-crystalline (Lindbladian PT) phase in the notation of collective spin. Moreover, we show that critical exceptional points (CEPs), in which two excitation modes coalesce, emerge in the case of continuous phase transitions. Finally, we confirm the validity of these analyses by comparing them with numerical calculations. As a result, we can expect that the exceptional point also emerges for the Lindbladian.

Lastly, we summarize and mention the outlook on the BTCs and Lindbladian PT phase transitions. In particular, we mention the application of Lindbladian exceptional points. The PT phase transitions in non-Hermitian systems have been used for many applications, such as sensors and lasers, by exploiting the properties of exceptional points. In particular, in order to be used as a sensor, both (i) the response coefficient of the output quantity to the input quantity of interest and (ii) the signal-to-noise ratio must be high. Around exceptional points, the property (i) satisfies. However, it has been shown that the signal-to-noise ratio for classical and quantum noise cannot improve even around exceptional points. On the other hand, by including nonlinearity to the Hamiltonian, it has recently been shown that the signal-to-noise ratio for classical noise can be improved around the exceptional point. Since the Driven Dicke model has nonlinearity at the mean-field level, it can be expected to improve the signal-to-noise ratio for classical noise around the transition point. In addition, since the Driven Dicke model shows exceptional points at the Lindbladian level, it would be very interesting to investigate whether the signal-to-noise ratio for quantum noise is also improved around the transition point. My thesis will not only further our understanding of non-equilibrium phase transitions or phases of matter, but also help us develop applied techniques using exceptional points in open quantum systems.