

論文 / 著書情報
Article / Book Information

論題	
Title	Possible anti-correlations between pulsation amplitudes and the disk growth of Be stars in giant-outbursting Be X-ray binaries
著者	庭野 聖史, 笹田 真人, 谷津 陽一, 河合 誠之
Authors	Masafumi Niwano, Ichiro Takahashi, Mahito Sasada, Yoichi Yatsu, Nobuyuki Kawai, Michael M. Fausnaugh, George R. Ricker, Roland Vanderspek, Kishalay De, Ryan M. Lau, Michael C. B. Ashley, Nicholas Earley, Mansi M. Kasliwal, Matthew J. Hankins, Anna M. Moore, Jamie Soon, Tony Trvouillon, Roberto Soria
出典	日本天文学会2024年秋季年会講演予稿集, , , N15a
会議名称	日本天文学会2024年秋季年会
発行日 / Pub. date	2024, 9
権利情報	本著作物の著作権は公益社団法人 日本天文学会に帰属します。

N15a Possible anti-correlations between pulsation amplitudes and the disk growth of Be stars in giant-outbursting Be X-ray binaries

Masafumi Niwano, Ichiro Takahashi, Mahito Sasada, Yoichi Yatsu, Nobuyuki Kawai (Tokyo Tech), Michael M. Fausnaugh (TTU), George R. Ricker, Roland Vanderspek, Kishalay De (Kavli-MIT), Ryan M. Lau (NOIR-Lab), Michael C. B. Ashley (UNSW), Nicholas Earley, Mansi M. Kasliwal (Caltech), Matthew J. Hankins (ATU), Anna M. Moore, Jamie Soon, Tony Travoignon (ANU), Roberto Soria (UCAS, INAF, USYD), MAXI team

The mechanism of X-ray outbursts in Be X-ray binaries remains a mystery, and understanding their circumstellar disks is crucial for a solution of the mass-transfer problem. In particular, it is important to identify the Be star activities (e.g., pulsations) that cause mass ejection and, hence, disk formation. Therefore, we investigated the relationship between optical flux oscillations and the infrared (IR) excess in a sample of five Be X-ray binaries. Applying the Lomb-Scargle technique to high-cadence optical light curves from the Transiting Exoplanet Survey Satellite (TESS), we detected several significant oscillation modes in the 3 to 24 hour period range for each source. We also measured the IR excess (a proxy for disk growth) of those five sources, using J-band light curves from Palomar Gattini-IR. In four of the five sources, we found anti-correlations between the IR excess and the amplitude of the main flux oscillation modes. This result is inconsistent with the conventional idea that non-radial pulsations drive mass ejections. We propose an alternative scenario where internal temperature variations in the Be star cause transitions between pulsation-active and mass-ejection-active states.