

論文 / 著書情報  
Article / Book Information

論題	Lower boundary of the Caledonian Barrovian metamorphic belt at Loch Leven, Scotland:
Title	Lower boundary of the Caledonian Barrovian metamorphic belt at Loch Leven, Scotland:
著者	青木一勝, ブライアン ウィンドレー, 佐藤佳子, 澤木佑介, 澁谷岳造, 熊谷英憲, 鈴木勝彦, 丸山茂徳
Author	Kazumasa Aoki, Brian Windley, Keiko Sato, Yusuke Sawaki, takazou shibuya, Hidenori Kumagai, Katsuhiko Suzuki, SHIGENORI MARUYAMA
掲載誌/書名	, , ,
Journal/Book name	, , ,
発行日 / Issue date	2012, 5

## Lower boundary of the Caledonian Barrovian metamorphic belt at Loch Leven, Scotland: Phengite K-Ar ages of metapelites

AOKI, Kazumasa<sup>1\*</sup>, Brian Windley<sup>2</sup>, Keiko Sato<sup>3</sup>, Yusuke Sawaki<sup>3</sup>, Takazo Shibuya<sup>3</sup>, Hidenori Kumagai<sup>3</sup>, Katsuhiko Suzuki<sup>3</sup>, Shigenori Maruyama<sup>4</sup>

<sup>1</sup>Dept of Earth and Astronomy, The Univ. Tokyo, <sup>2</sup>Dept of Geology, Leicester Univ, <sup>3</sup>JAMSTEC, <sup>4</sup>Earth and Planetary Sci., Tokyo Tech

Since the Barrow's report (1893), the Barrovian metamorphic belt in the Caledonides of Scotland has become one of the best-studied orogenic belts in the world. Based on many geological and isotopic studies, this belt has long been considered to be formed by arc-continental collision in the Ordovician-Silurian (the Grampian orogeny: ca. 480-430 Ma) (Dewey and Bird, 1999; Oliver, 2008). However, many controversies and fundamental uncertainties remain. For example, in terms of the polarity of subduction, Ryan and Dewey (1991) and Oliver (2008) assumed that the polarity of subduction was toward the NW, but Cliff et al. (2004) proposed the opposite. Also, the main reason for the "orogeny" and metamorphism for several decades has been just burial at depth caused by thrusting, which we consider to be a dated, unlikely concept.

The Barrovian metamorphic rocks in Loch Leven belong to the biotite zone, have been folded and thrust northwards (e.g., Treagus, 1974; Piasecki, 1980), and they overlie weakly metamorphosed rocks (Eilde Flags). Inferring from our understandings of the occurrence and exhumation of HP rocks in Japan (Aoki et al., 2008; 2009), and in many Barrovian orogenic belts worldwide (Agard et al., 2009), we predicted that the lower tectonic boundary of the Barrovian metamorphic rocks should have occurred at Loch Leven. In order to examine this model, we determined K-Ar ages of phengite-rich mineral separates from 6 metapelites in the area.

Our new phengite K-Ar ages are 398.2 +/- 10.4 (sample no. LL46), 406.4 +/- 10.6 (LL20), 405 +/- 10.5 (LL19), 399.7 +/- 10.5 (LL24) and 445.0 +/- 11.6 Ma (LL16). One sample (LL13) has an "anomalously young" age of 340 +/- 8.9 Ma, probably because of its low K<sub>2</sub>O content (1.3 wt%). The main results indicate a K-Ar age-gap between ca. 445 Ma and ca. 400 Ma, which, when integrated with previous metamorphic-age (e.g., Oliver, 2008) and structural data from Loch Leven (Rovers, 1976; Atherton, 1977) suggest that the lower boundary of the Barrovian metamorphic belt formed at 445-400 Ma.

The lower boundary of the E-dipping, Barrovian metamorphic belt at Loch Leven has long been considered to be an extension of the 430-413 Ma Moine Thrust (MT) displaced by the Great Glen Fault. The shear sense of the E-dipping MT is top-to-the-WNW/NNW (Butler, 2004). Moreover, we have recently discovered that the upper boundary of the originally E-dipping Barrovian metamorphic belt crops out near Portsoy in NE Scotland, the shear sense of which was top-to-the-SW (Kawai et al., in prep.). In terms of the internal thermobaric Barrovian zones of Scotland, the grade of metamorphism decreases symmetrically upwards and downwards from a central highest-grade zone (Kennedy, 1948), which contains relicts of retrogressed HP rocks. Integration of all multi-disciplinary data provides robust evidence that the Barrovian metamorphic zones were exhumed from a HP depth by N-directed wedge extrusion. The lower wedge boundary was the Moine thrust-Loch Leven thrust, and the upper extensional boundary at Portsoy enabled the downward emplacement of the overlying, lower pressure Buchan metamorphic zonal belt. Thus, formation of the whole Caledonian orogenic belt of Scotland was controlled by wedge extrusion.