

論文 / 著書情報
Article / Book Information

題目(和文)	サンゴ礁域における水理ならびに幼生分散過程に関する数値モデル解析と現地観測
Title(English)	Computational model analyses and field observations of hydrodynamic and larval dispersal processes in coral reef areas
著者(和文)	LBERNARDO
Author(English)	Lawrence Patrick Cases Bernardo
出典(和文)	学位:博士(学術), 学位授与機関:東京工業大学, 報告番号:甲第10893号, 授与年月日:2018年3月26日, 学位の種別:課程博士, 審査員:灘岡 和夫,中村 隆志,鼎 信次郎,中村 恭志,吉村 千洋
Citation(English)	Degree:Doctor (Academic), Conferring organization: Tokyo Institute of Technology, Report number:甲第10893号, Conferred date:2018/3/26, Degree Type:Course doctor, Examiner:,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	論文要旨
Type(English)	Summary

論文要旨

THESIS SUMMARY

専攻 : Department of	Civil Engineering	専攻	申請学位 (専攻分野) : Academic Degree Requested	博士 Doctor of	(Philosophy)
学生氏名 : Student's Name	Lawrence Patrick Cases Bernardo		指導教員 (主) : Academic Supervisor(main)	Kazuo Nadaoka	
			指導教員 (副) : Academic Supervisor(sub)	Takashi Nakamura	

要旨 (英文 800 語程度)

Thesis Summary (approx.800 English Words)

Coral reef ecosystems are marine ecosystems of great significance. They have been generally known to be sites of high marine biodiversity and provide a wide variety of ecosystem services, such as food security in the form of fishery resources, coastal protection through ocean wave attenuation, and tourism due to their natural attractiveness. Given their importance, there is a continuing need to study coral reef ecosystems especially during this recent period of alarming environmental changes. This research was conducted with the aim of addressing this need and contribute to the body of work on coral reefs by putting together data collected from various field survey activities, analyzing the data using different techniques, and the development of modeling approaches to supplement the observations and provide further insights. A nested modeling configuration was developed and used to investigate hydrodynamics around the Yaeyama Islands and in the Sekisei Lagoon area. The Regional Ocean Modelling System (ROMS) was adapted for the study site and the configuration used in this study consisted of a 1.5 km grid resolution Yaeyamal model which included the eastern side of Taiwan and ocean waters surrounding the Yaeyama Island. A 300 m grid resolution Yaeyama2 was nested on the Yaeyamal model and was used to more closely investigate hydrodynamics in Sekisei Lagoon. Oceanographic sensor deployments provided a means to validate results generated by the model, and model performance was found to be satisfactory upon evaluation using a widely used skill index.

While widely known to be destructive, typhoon events can benefit coral reef ecosystems through typhoon-induced cooling, which can mitigate against thermally stressful conditions causing coral bleaching. Sensor deployments in Sekisei Lagoon, Japan's largest coral reef area, during the summer months of 2013, 2014, and 2015 were able to capture local hydrodynamic features of numerous typhoon passages. Through an analysis of the field data and jointly investigated using Regional Ocean Modeling System (ROMS) numerical simulations conducted for the summer of 2015, the presence of an island-enhanced cooling mechanism in typhoon events was discovered. Local cooling events may have been initiated by the Yaeyama Islands acting as an obstacle to a strong typhoon-generated flow which was modulated and led to prominent cooling of waters on the leeward sides. Such cooled waters could then enter the inner Sekisei Lagoon area likely through density-driven flow.

The modeling configuration was also used to simulate hydrodynamics in Sekisei Lagoon for April to November 2016, during which massive coral bleaching occurred. Temperature sensor deployments and coral bleaching surveys were also meticulously conducted by the Ministry of the Environment during this period. The model exhibited relatively good skill at reproducing the general temperature trends in the sensor data. Model results were also used to calculate accumulated heating and cooling based on specified threshold temperatures, and the resulting outputs may aid in the interpretation of spot check field survey results which looked at the spatial trend in the percentages of bleached and dead corals. Unlike the observed temperature data which were at specific points and relatively shallow depths, the model was able to show a more complete spatial picture of which areas potentially experienced the most heat stress as well as those which were cooled effectively by the passage of typhoons.

Water temperature at different locations within and just offshore of Shiraho Reef, Okinawa, Japan, as well as offshore bottom current and wave height have been continuously monitored since 2010 using standalone sensors. Trends in the long-term dataset were examined to gain more insight into the mechanisms that may be causing them through various analyses methods. While inner reef temperatures were strongly affected by the diurnal cycle, offshore temperatures were shown to be significantly affected by regional circulation trends. A particularly notable transition between lower and higher temperatures generally occurring in the month April, characterized by pulses of cooling and warming over a span of 2 to 3 weeks. This feature was investigated using Regional Ocean Modeling System (ROMS) numerical simulations and was found to be possibly linked to regional flow patterns, particularly changes in the Kuroshio flow

characteristics and subsequent eddy formation in the vicinity of the Yaeyama Islands. Simulations for the winter season were also conducted and further supported the potential influence of regional eddy activity on the local temperatures in Shiraho Reef.

The nested hydrodynamic model was also used to run particle tracking scenarios to simulate potential larval dispersal within the Sekisei Lagoon domain and the coastal areas of the Yaeyama Islands. Particle tracking experiments were focused on attempting to characterize the differences in larval dispersal trajectories between relatively “normal” hydrodynamic conditions and hydrodynamics during the passage of a strong typhoon. In 2015, 3 strong typhoons with relatively distinct tracks struck the area, and model-generated hydrodynamics during the passage of these typhoons was used as the basis to run typhoon track dispersal cases. Particle release locations were then organized into more distinct numbered sites around the coastal areas and parts of Sekisei Lagoon to generate connectivity matrices which quantitatively described the marine interconnectivity between the different locations.

備考：論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note : Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1copy of 800 Words (English).

注意：論文要旨は、東工大リサーチリポジトリ(T2R2)にてインターネット公表されますので、公表可能な範囲の内容で作成してください。

Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).