

論文 / 著書情報  
Article / Book Information

題目(和文)	トンレサップ湖の洪水氾濫原における土砂動態
Title(English)	Sediment Dynamics in the Floodplain of Tonle Sap Lake
著者(和文)	SievSokly
Author(English)	Sokly Siev
出典(和文)	学位:博士(工学), 学位授与機関:東京工業大学, 報告番号:甲第10913号, 授与年月日:2018年6月30日, 学位の種別:課程博士, 審査員:吉村 千洋,藤井 学,鼎 信次郎,竹村 次朗,中村 恭志,藤井 秀人
Citation(English)	Degree:Doctor (Engineering), Conferring organization: Tokyo Institute of Technology, Report number:甲第10913号, Conferred date:2018/6/30, Degree Type:Course doctor, Examiner:,,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	論文要旨
Type(English)	Summary

(博士課程)  
Doctoral Program

## 論文要旨

THESIS SUMMARY

専攻:	Civil Engineering	専攻
Department of		
学生氏名:	Siev Sokly	
Student's Name		

申請学位 (専攻分野):	博士 (Engineering)
Academic Degree Requested	Doctor of
指導教員 (主):	Yoshimura Chihiro
Academic Supervisor(main)	
指導教員 (副):	
Academic Supervisor(sub)	

要旨 (英文 800 語程度)

Thesis Summary (approx.800 English Words )

Generally, resuspension is a dominant process of sediment dynamics in a stable shallow lake. However, resuspension might not be the dominant process in a shallow lake influenced by flood pulse because floodplain vegetation most likely plays as a significant role in lessening the intensity of resuspension and promoting sedimentation process. Little attention, however, has been paid to this type of lake because investigations of sediment dynamics have typically focused on relatively stable shallow lakes. Therefore, this research aimed to investigate the sediment dynamics in a large shallow lake and its floodplain characterized by flood pulse using field observation and modelling techniques. The targeted shallow lake was Tonle Sap Lake (TSL), the largest freshwater lake in Southeast Asia, located in Cambodia. This important study not only improved our understanding of sediment dynamics in shallow lake, but also contributed to the ecological and environmental management of ecosystems in such lakes driven by the flood pulse such as TSL.

A major objective in this research has been the investigation of sediment dynamics in a large shallow lake and its floodplain characterized by flood pulse using field observation. An extensive and seasonal sampling survey was conducted to measure total suspended solid (TSS) concentrations, sedimentation and resuspension rates in TSL and its 4 floodplain areas. As the results, TSS concentrations ranged from 1.3 to 751.8 mg/L during the sampling period (September 2016 to June 2017). The concentration was highest in March, followed by June, September, and December. The average resuspension rates were  $116.6 \pm 146.4$  g/m<sup>2</sup>/day,  $178.2 \pm 205.2$  g/m<sup>2</sup>/day,  $215.4 \pm 365.3$  g/m<sup>2</sup>/day, and  $306.7 \pm 369.6$  g/m<sup>2</sup>/day in grassland, shrub, flooded forest, and open water areas, respectively. These values corresponded to 81.2%, 72.8%, 59%, and 85.3% of the average gross sedimentation rates, respectively. This study, for the first time, conducts a quantitative measurement of resuspension rate contributing to not only the knowledge of sediment dynamics in TSL but also the large shallow lake studies around the globe. The measured resuspension rate was meaningful to explain the resuspension process and the reason why the sedimentation rate in TSL is very low, which is a substantial improvement in our knowledge from the previous studies. The study revealed that sedimentation process was dominant in the high water period (September-December) while resuspension process was dominant only in the low water period (March-June). In addition, floodplain vegetation reduced the resuspension of sediment (up to 26.3%) in water.

Another major objective of this research that stands out was the model application and improvement of a land landscape evolution model (LEM). Most of LEMs considered only a simplification of vegetation growth of a grass surface, not a spatial variability of vegetation community. Therefore, this study aims to improve a landscape evolution model by integrating spatial variability of vegetation and to apply the integrated model for investigation of sediment dynamics. Boundary conditions (e.g. water discharge and sediment load) from surrounding tributaries of TSL and Tonle Sap River (TSR) were estimated using a physical process-based sediment model and semi distributed sediment model, respectively. The Nash-Sutcliffe Efficiency (NSE) of both models ranged from satisfactory to good performance for the discharge (NSE ranged 0.65-0.85) and sediment load (NSE ranged 0.44-0.79) which were good for the subsequent investigation. A landscape evolution model, Caesar-Lisflood (CL) model was successfully applied in TSL and TSR, having a good and acceptable simulation performance for the water level (NSE: 0.93) and suspended sediment concentration ( $R^2=0.73$  in September), respectively. Moreover, the study integrated and tested the spatial variability of vegetation (e.g. grass, shrub and forest) in the model. As the result, compared to the single type of vegetation (e.g. CL-G, CL-S and CL-F cases), the integrated spatial variability vegetation CL model (CL-G-S-F case) showed spatial difference of flow velocity resulting in different erosion and sedimentation patterns in the floodplain vegetation areas. The simulated sedimentation rate of CL-G-S-F case also showed a relative fit with the observed data at Chhnok Tru area in each type vegetation zones (e.g., grass, shrub and forest). The improved CL model was also a useful tool for evaluating the impact of some scenarios (e.g. shift in water level and suspended sediment load of boundary conditions) and supporting environment management. Hence, the overall research title is deemed appropriate, as the study indeed has in various ways attempted to generate insight into some importance processes related to sediment dynamics in a large shallow lake influenced by flood pulse, mainly through an analysis of field observations and modelling techniques.

備考 : 論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note : Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1copy of 800 Words (English).

注意 : 論文要旨は、東工大リサーチリポジトリ (T2R2) にてインターネット公表されますので、公表可能な範囲の内容で作成してください。

Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).