

論文 / 著書情報
Article / Book Information

題目(和文)	高圧二酸化炭素中での生体触媒反応：リパーゼの触媒活性の向上
Title(English)	Pressurized carbon dioxide in biocatalysis: Enhanced catalytic activity of lipase
著者(和文)	HOANGNAM HAI
Author(English)	Hai Hoang
出典(和文)	学位:博士(工学), 学位授与機関:東京工業大学, 報告番号:甲第10971号, 授与年月日:2018年9月20日, 学位の種別:課程博士, 審査員:松田 知子,中村 聡,小林 雄一,占部 弘和,福居 俊昭
Citation(English)	Degree:Doctor (Engineering), Conferring organization: Tokyo Institute of Technology, Report number:甲第10971号, Conferred date:2018/9/20, Degree Type:Course doctor, Examiner:,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	要約
Type(English)	Outline

Pressurized Carbon Dioxide in Biocatalysis: Enhanced Catalytic Activity of Lipase

Hoang Nam Hai

[Introduction]

The scope of biocatalysis, particularly in enantioselective synthesis of target molecules for pharmaceuticals, has been extended through the use of enzymes in organic solvents and in “green” nonaqueous media. Carbon dioxide (CO_2), which has numerous positive impacts on green chemistry, has been utilized as novel and sustainable reaction platform for lipase catalysis in this study (Figure 1).

[Results and Discussion]

1. Liquid carbon dioxide: Broadening substrate scope of lipase catalysis

The kinetic resolution of typical substrate *rac*-1-phenylethanol by *Candida antarctica* lipase B (CALB, Novozym 435) using vinyl acetate as an acyl donor was investigated as a model reaction in liquid CO_2 and compared with that in conventional organic solvents. The enzyme exhibits the highest activity in liquid CO_2 with excellent enantioselectivity. The potential applicability of liquid CO_2 fluid for large-scale enantioselective synthesis was also examined with continuous flow bioreactors. The continuous kinetic resolution of *rac*-1-phenylethanol by the immobilized lipase in a packed-column reactor stably afforded corresponding enantiopure products with minimum waste generated (Figure 1). Interestingly, by using in liquid CO_2 . The substrate scope CALB was found to be expanded towards sterically bulky *sec*-alcohols such as 1-phenyldodecanol, of which the lipase has been known to have no reactivity in standard conventional solvent.




Fig. 1 Biocatalytic synthesis using liquid CO_2

2. CO_2 -expanded bio-based liquids: Modulating lipase activity by controlling physicochemical properties of solvent

As emerging alternative solvents, bio-based liquids provide environmentally advantages, such as sustainable production, better biodegradability and lower toxicity. However, some bio-based liquids with high polarity are not favourable solvents for biocatalysis, such as 2-methyltetrahydrofuran (MeTHF), as it often witnessed in decreased lipase activity. With tunability, pressurized CO_2 was employed to manipulate the physicochemical properties




Fig. 2 Tuning CALB activity in CO_2 -expanded MeTHF (unfilled legends) and solvent property (dipolarity π^* , filled legends) as a function of CO_2 mole fraction X_{CO_2} . Reaction conditions: 0.10 mmol *rac*-1-adamantylethanol, 0.53 mmol vinyl acetate, 10 mg Novozym 435, 1.0 mL MeTHF, pressures ranging from 0.1 MPa to 9 MPa. Reaction rate (μmol substrate/mg enzyme/h) was normalized to 1 for the reaction at $X_{\text{CO}_2}=0$ of each temperature (round 20 °C, triangle 40 °C, diamond 60 °C).

such as dipolarity π^* of many bio-based liquids into favorable platforms for lipase catalysis (herein called as CO₂-expanded bio-based liquids, Figure 2).

3. Molecular behaviour of *Candida antarctica* lipase B in pressurized carbon dioxide

The underlying mechanism of crucial role of CO₂ to induce increased activity of lipase was studied. Molecular dynamics simulation of CALB showed that pressurized CO₂ can modify the structure flexibility of the active-site entrance, which allows bulkier substrates to reach the active site cavity of CALB. Moreover, the volume of main cavity of CALB was found to be enlarged in pressurized CO₂, so that the enzyme can comfortably accommodate bulkier substrates. On the basis of the experimental and computational findings, the structure of CALB in pressurized CO₂ were proposed as in Figure 3.




Fig. 3 Proposed model of CALB structures

[Conclusion]

Herein, pressurized CO₂ (liquid CO₂ and CO₂-expanded bio-based liquids) was proved as superior media for lipase catalysis. In pressurized CO₂ not only much higher activity were possible, also CALB retained its originally high enantioselectivity towards various sterically bulky *sec*-alcohols. The molecular dynamics simulation of CALB in pressurized CO₂ showed important changes in the lipase conformation that the lipase became more tolerant to bulkier substrates. With these positive findings, I expected that pressurized CO₂ is promising non-aqueous media for other biocatalysis.

[Publications]

- (1) Hoang, H. N.; Are, K. R. A.; Matsuda, T. Biocatalysis in supercritical and liquid carbon dioxide and carbon dioxide expanded liquids. In *Supercritical and Other High-pressure Solvent Systems: For Extraction, Reaction and Material Processing*; Hunt, A. J., Attard, T. M., Eds.; Green Chemistry Series; The Royal Society of Chemistry: London, in press.
- (2) Hoang, H. N.; Granero-Fernandez, E.; Yamada, S.; Mori, S.; Kagechika, H.; Medina-Gonzalez, Y.; Matsuda, T. Modulating biocatalytic activity toward sterically bulky substrates in CO₂-expanded biobased liquids by tuning the physicochemical properties. *ACS Sustain. Chem. Eng.* **2017**, 5 (11), 11051–11059.
- (3) Hoang, H. N.; Matsuda, T. Biotransformation using liquid and supercritical CO₂. In *Future Directions in Biocatalysis*; Matsuda, T., Ed.; Elsevier: Amsterdam, 2017; pp 3–25.
- (4) Hoang, H. N.; Nagashima, Y.; Mori, S.; Kagechika, H.; Matsuda, T. CO₂-expanded bio-based liquids as novel solvents for enantioselective biocatalysis. *Tetrahedron* **2017**, 73 (20), 2984–2989.
- (5) Hoang, H. N.; Matsuda, T. Expanding substrate scope of lipase-catalyzed transesterification by the utilization of liquid carbon dioxide. *Tetrahedron* **2016**, 72 (46), 7229–7234.
- (6) Hoang, H. N.; Matsuda, T. Liquid carbon dioxide as an effective solvent for immobilized *Candida antarctica* lipase B catalyzed transesterification. *Tetrahedron Lett.* **2015**, 56 (4), 639–641.