

論文 / 著書情報
Article / Book Information

題目(和文)	微衛星の形成における困難とガリレオ衛星の起源についての新しいシナリオ
Title(English)	Difficulties in Formation of Satellitesimals and a New Scenario for the Origin of the Galilean Satellites
著者(和文)	芝池諭人
Author(English)	Yuhito Shibaike
出典(和文)	学位:博士(理学), 学位授与機関:東京工業大学, 報告番号:甲第11055号, 授与年月日:2019年3月26日, 学位の種別:課程博士, 審査員:井田 茂,綱川 秀夫,関根 康人,奥住 聡,玄田 英典
Citation(English)	Degree:Doctor (Science), Conferring organization: Tokyo Institute of Technology, Report number:甲第11055号, Conferred date:2019/3/26, Degree Type:Course doctor, Examiner:,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	論文要旨
Type(English)	Summary

論文要旨

THESIS SUMMARY

系・コース：	地球惑星科学	系
Department of, Graduate major in	地球惑星科学	コース
学生氏名：	芝池論人	
Student's Name		

申請学位 (専攻分野)：	博士	(理学)
Academic Degree Requested	Doctor of	
指導教員 (主)：	井田茂	
Academic Supervisor(main)		
指導教員 (副)：		
Academic Supervisor(sub)		

要旨 (英文 800 語程度)

Thesis Summary (approx.800 English Words)

It is generally accepted that the large satellites around gas planets formed out of the gas disks, the circumplanetary disks (CPDs), which accompanied the planets' formation. Previous works showed that if enough satellitoids (km-sized bodies) exist in the disk, satellites with the current satellites' mass can form by two-body collisions. A problem with this scenario, however, is the formation of satellitoids. Unless CPDs features pressure reversals, the progenitor dust grains will not have had the time to conglomerate, because of the strong radial drift. Moreover, gas gap structures formed by the planets must prevent the dust from flowing into CPDs. An alternative idea to get the material of satellites is planetesimal-capture. However, the growing planets and the gas gaps push out the planetesimals from the feeding-zones and the capture rates decrease. Therefore, how to get the material is a big issue in satellite formation research. Another problem is difficulty in reproducing the characteristics of the satellites simultaneously and consistently. In the case of the Galilean (Jovian) satellites, there are a lot of studies about the characteristics but they can only explain a part of them and contradict each other in part. First, we investigated the possibility that satellitoids form in-situ in CPDs. We calculated the radial distribution of the surface density and representative size of dust particles. They grow by mutual collisions and drift toward the planet in with a continuous supply of gas and dust from the parent PPD. In order for the particles to form satellitoids, the growth must take place faster than they fall onto the planet. We found that this radial drift barrier is overcome if the particles can grow larger than the size which their Stokes number (stopping time normalized by the Kepler-time) is greater than unity. This condition is the same with that of planetesimal formation in PPDs. We also found that the satellitoids can form (i.e. the above condition is achieved) if the ratio of the dust to gas accretion rates into CPDs is larger than unity. The dust-to-gas ratio must be lower than unity in general, our results suggest that the in-situ satellitoid formation via particle sticking is viable only under an extreme condition. However, satellitoids may form episodically or locally in special conditions, for example, the dust vertical diffusion is weak and the gas surface density is low, predicted in some recent hydrodynamical simulations. We then constructed a new alternative scenario for the origin of the Galilean satellites which can form the satellites even in the material scarcity. In our new scenario, two kinds of small amount of solid material are supplied to the CPD; mm-sized dust particles and km-sized planetesimals. The captured planetesimals accrete the particles drifting toward Jupiter (i.e. pebbles) so effectively that the planetesimals can grow to satellites before the disk disappears thanks to strong gravitational focusing and aerodynamic drag. However, the growth timescale is longer than that of classical formation scenarios, we invoked an inner disk truncation radius to halt their migration. Other parameters were tuned for the model to match physical, dynamical, compositional, and structural constraints of the Galilean satellites. In contrast to the previous scenarios, our new "slow-pebble-accretion" scenario then reproduced most of the important characteristics simultaneously and consistently; (1) the mass of all the Galilean satellites, (2) the orbits of Io, Europa, and Ganymede captured in mutual 2:1 mean motion resonances, (3) the ice mass fractions of all the Galilean satellites, and (4) the unique ice-rock undifferentiated Callisto and the other three differentiated satellites. However, only Callisto's orbit could not be reproduced but it was also captured in resonance with Ganymede's one. We found that the in-situ satellitoid formation in CPDs is very difficult. Therefore, we built a new formation scenario especially for the Galilean satellites, using drifting pebbles instead of satellitoids. However, our simulations are simplified and, in reality, satellitoids originated from special conditions may contribute to satellite formation. Therefore, we should investigate satellite formation processes including both satellitoids and pebbles as a next step. Future measurements of the D/H ratio in H₂O ice of the Galilean satellites by JUICE mission may be able to find out which accretion mechanism was actually dominant. We are now also working on 2-D orbital simulations of the satellites to investigate the resonance capture to solve the problem of Callisto's orbit because we have only carried out simple 1-D simulations. The origin of the large Saturnian satellites should also be investigated. Our discussion about satellitoid formation can be applied to the Saturnian system. However, more explanations for the long orbital periods of the satellites will be needed for applying the new Galilean satellite formation scenario to the Saturnian system. General satellite formation will be an attractive research area in future. In particular, satellite formation associated with the evolution of the planets' envelope will be important. Finally, observations of exomoons will open the way to statistical general satellite formation studies.

備考：論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note：Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1 copy of 800 Words (English).

注意：論文要旨は、東工大リサーチリポジトリ(T2R2)にてインターネット公表されますので、公表可能な範囲の内容で作成してください。

Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).