

論文 / 著書情報
Article / Book Information

題目(和文)	深層自己符号化器によるクエーサー可視光光度変動の特徴量抽出
Title(English)	Deep Autoencoder Applied for Feature Extraction of Temporal Variations in Quasar Optical Flux
著者(和文)	橘優太郎
Author(English)	Yutaro Tachibana
出典(和文)	学位:博士(理学), 学位授与機関:東京工業大学, 報告番号:甲第11042号, 授与年月日:2019年3月26日, 学位の種別:課程博士, 審査員:河合 誠之,堂谷 忠靖,中村 隆司,西森 秀稔,宗宮 健太郎,篠田 浩一
Citation(English)	Degree:Doctor (Science), Conferring organization: Tokyo Institute of Technology, Report number:甲第11042号, Conferred date:2019/3/26, Degree Type:Course doctor, Examiner:,,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	要約
Type(English)	Outline

Tile

Deep Autoencoder Applied for Feature Extraction of Temporal Variations in Quasar Optical Flux

Author

Yutaro Tachibana

Affiliation

Department of Physics, Tokyo Institute of Technology, 2-12-1
Ookayama, Meguro-ku, Tokyo 152-8551, Japan

Abstract

Quasars have long been known as intrinsically variable sources, but the physical mechanism underlying the temporal optical/UV variability is still under discussion from the first identification of the optical variability of quasars in the middle of the twentieth century. In this paper, we proposed a nonparametric method for modeling and forecasting the optical temporal variation of quasars utilizing an autoencoder neural network, to gain insight of the physical mechanism on the temporal flux variation of quasars. The autoencoder is trained and validated by ~ 10 yrs long $\sim 15,000$ quasar light curves obtained by the Catalina Real-Time Transient Survey, where the quasars are selected to be the flux contamination from the host galaxy is negligible in the temporal variation. Its performance in forecasting the temporal flux variation is superior to that of the Ornstein–Uhlenbeck process. In addition, a novel relation, the magnitude of the temporal variability asymmetry decreases as luminosity and/or black hole mass increases, is suggested with the help of autoencoded features. We conclude that the characteristics of the variability asymmetry are in agreement with those resulted in from simulation with the self-organized disk instability model, which predicts that the magnitude of the variability asymmetry decrease as the ratio of the diffusion mass to inflow mass in the accretion disk increases.