

論文 / 著書情報  
Article / Book Information

題目(和文)	M型矮星の特性決定
Title(English)	Characterization of M-type Dwarfs
著者(和文)	小泉陽平
Author(English)	Yohei Koizumi
出典(和文)	学位:博士(理学), 学位授与機関:東京工業大学, 報告番号:甲第11885号, 授与年月日:2021年3月26日, 学位の種別:課程博士, 審査員:佐藤 文衛,井田 茂,奥住 聡,玄田 英典,中島 淳一
Citation(English)	Degree:Doctor (Science), Conferring organization: Tokyo Institute of Technology, Report number:甲第11885号, Conferred date:2021/3/26, Degree Type:Course doctor, Examiner:,,,,,
学位種別(和文)	博士論文
Category(English)	Doctoral Thesis
種別(和文)	論文要旨
Type(English)	Summary

# 論文要旨

## THESIS SUMMARY

系・コース： Department of, Graduate major in	地球惑星科学 地球惑星科学	系 コース	申請学位(専攻分野)： Academic Degree Requested	博士 Doctor of	(理学)
学生氏名： Student's Name	小泉 陽平		指導教員(主)： Academic Supervisor(main)	佐藤 文衛	
			指導教員(副)： Academic Supervisor(sub)		

### 要旨 (英文 800 語程度)

Thesis Summary (approx.800 English Words)

An M-type dwarf is a cool and small star in the main sequence and it has been gathering attention as a promising target for detecting Earth-like small planets. Since the estimation of planetary properties strongly depends on the properties of the host star, understanding its stellar properties such as temperature, radius, mass, and metallicity is vital for planet search projects and for understanding planetary systems. Although the characterization of a Solar-like star is generally performed via comparisons of a high-resolution spectrum with a synthetic model atmosphere, that for the M-type dwarf is significantly more complicated than that for the Sun-like FGK type star. The spectrum of an M type dwarf has strong and complex atomic and molecular absorption bands, which result in significantly blended spectral lines and unclear continuum regions. The current model atmosphere of low mass stars suffers from incomplete knowledge of stellar physics, line opacity, and other issues. Therefore, the estimation based on a synthetic model atmosphere may fail to yield accurate stellar properties. Recently, several studies have suggested that machine learning tasks can provide model-independent stellar properties for M-type dwarfs. However, in order to apply these data-driven methods, the spectra must be obtained with the same spectrograph. This work presents the methods to estimate the stellar parameters of low mass stars based on various observations.

In Chapter 2, the temperatures and radii of M-type dwarfs are revised based on the optical spectra and photometric observations with the latest parallaxes. The optical spectra are obtained in the target selection observation of the planet search program. This observation is conducted using several instruments. Therefore, the temperatures are thus calculated by comparing the optical spectra with the synthetic spectra. To estimate the precise temperatures, the fitting procedure is calibrated by using the M-type dwarfs whose temperatures are well-determined. For spectra with limited or narrow wavelength coverages, it was found that the temperature estimation was refined by utilizing only VO absorption features if available. Combining near infrared spectra with the optical spectra in this work will enable us to obtain more robust temperature estimations. Even if the synthetic model atmosphere is poorly constrained, selecting the suitable wavelength region and fitting the synthetic model spectra are still useful in order to estimate temperatures. The temperatures and radii of M-type dwarfs are also calculated via spectral energy distributions with the

latest parallaxes. The Comparison of the resulting radii with the empirical ones and those from the stellar evolution model suggests that the radii are larger than the empirical and theoretical predictions.

The stellar properties of M-type dwarfs are determined using the data-driven method of optical low-resolution spectra in Chapter 3. Sparse modeling, which is a machine learning task, can be used to determine model-independent temperatures and metallicities with a precision of  $\sim 50$  K and 0.09 dex, respectively. The sparse method of optical spectra can provide accurate metallicity for metal-poor stars, and the metallicity trend is observed in higher metallicity regions. This suggests that the low-resolution spectroscopy does not contain sufficient resolution power to estimate higher metallicity values.

Further, the sparse approximation is also applied to M-type dwarfs that have near infrared high-resolution spectra and provides temperatures, radii, masses, and metallicities in Chapter 4. The proposed method can obtain the properties of M-type dwarfs with the precision of 109 K in temperature, 6% in radius, 6% in mass, and 0.07 dex in metallicity. Similar to the sparse modeling of low-resolution spectroscopy, the presented method also does not utilize the synthetic spectra of low mass stars that suffer from the incomplete knowledge of the stellar atmosphere. The estimation of metallicity with high-resolution spectra yields better results than that with low-resolution spectra and that with the synthetic model spectra. The proposed sparse method can derive the metallicity values of the planet-host M-type dwarfs, which confirm a correlation between planetary mass and stellar metallicity. The result supports the scenario that a metal-rich star tends to have dust-rich disks and it results in less Earth-like planets around those stars.

The characterization of M-type dwarfs remains challenging. However, combining various types of observations enable us to provide accurate stellar properties. This study emphasizes that the data-driven methods can help achieve breakthroughs in stellar characterization because sparse modeling does not make use of the synthetic model atmosphere which suffers from the incomplete knowledge of stellar physics. Currently, M-type dwarfs are one of the primary targets not only in radial velocity surveys but also in the transit survey. Several planet search projects are planning or conducting their observations. Without accurate stellar properties, understanding planetary systems and constraining planet formation theory cannot be achieved. This research is expected to make a significant contribution to such planet survey projects and to understanding planetary systems.

備考：論文要旨は、和文 2000 字と英文 300 語を 1 部ずつ提出するか、もしくは英文 800 語を 1 部提出してください。

Note : Thesis Summary should be submitted in either a copy of 2000 Japanese Characters and 300 Words (English) or 1 copy of 800 Words (English).

注意：論文要旨は、東工大リサーチリポジトリ (T2R2) にてインターネット公表されますので、公表可能な範囲の内容で作成してください。

Attention: Thesis Summary will be published on Tokyo Tech Research Repository Website (T2R2).